

Jean T Greenberg

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5848069/jean-t-greenberg-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

70
papers

8,858
citations

45
h-index

74
g-index

74
ext. papers

9,944
ext. citations

8.1
avg, IF

6.05
L-index

#	Paper	IF	Citations
70	"How Do We Do This at a Distance?!" A Descriptive Study of Remote Undergraduate Research Programs during COVID-19.. <i>CBE Life Sciences Education</i> , 2022 , 21, ar1	3.4	4
69	<i>Pseudomonas syringae</i> effector HopZ3 suppresses the bacterial AvrPto1-tomato PTO immune complex via acetylation. <i>PLoS Pathogens</i> , 2021 , 17, e1010017	7.6	3
68	Kinases and protein motifs required for AZI1 plastid localization and trafficking during plant defense induction. <i>Plant Journal</i> , 2021 , 105, 1615-1629	6.9	1
67	ALD1 accumulation in Arabidopsis epidermal plastids confers local and non-autonomous disease resistance. <i>Journal of Experimental Botany</i> , 2021 , 72, 2710-2726	7	2
66	An Improved Bioassay to Study Induced Systemic Resistance (ISR) Against Bacterial Pathogens and Insect Pests. <i>Bio-protocol</i> , 2019 , 9, e3236	0.9	5
65	Underground Azelaic Acid-Conferred Resistance to <i>Pseudomonas syringae</i> in Arabidopsis. <i>Molecular Plant-Microbe Interactions</i> , 2019 , 32, 86-94	3.6	14
64	PROHIBITIN3 Forms Complexes with ISOCHORISMATE SYNTHASE1 to Regulate Stress-Induced Salicylic Acid Biosynthesis in Arabidopsis. <i>Plant Physiology</i> , 2018 , 176, 2515-2531	6.6	31
63	Simple strategies to enhance discovery of acetylation post-translational modifications by quadrupole-orbitrap LC-MS/MS. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2018 , 1866, 224-229	4	3
62	A Suite of Receptor-Like Kinases and a Putative Mechano-Sensitive Channel Are Involved in Autoimmunity and Plasma Membrane-Based Defenses in Arabidopsis. <i>Molecular Plant-Microbe Interactions</i> , 2017 , 30, 150-160	3.6	13
61	Flagellin peptide flg22 gains access to long-distance trafficking in Arabidopsis via its receptor, FLS2. <i>Journal of Experimental Botany</i> , 2017 , 68, 1769-1783	7	20
60	Carbon Nanofiber Arrays: A Novel Tool for Microdelivery of Biomolecules to Plants. <i>PLoS ONE</i> , 2016 , 11, e0153621	3.7	6
59	Arabidopsis AZI1 family proteins mediate signal mobilization for systemic defence priming. <i>Nature Communications</i> , 2015 , 6, 7658	17.4	70
58	The Plant Cell Introduces Breakthrough Reports: A New Forum for Cutting-Edge Plant Research. <i>Plant Cell</i> , 2015 , tpc.15.00862	11.6	78
57	Linking pattern recognition and salicylic acid responses in Arabidopsis through ACCELERATED CELL DEATH6 and receptors. <i>Plant Signaling and Behavior</i> , 2015 , 10, e1010912	2.5	12
56	Acetylation of an NB-LRR Plant Immune-Effector Complex Suppresses Immunity. <i>Cell Reports</i> , 2015 , 13, 1670-82	10.6	46
55	ALD1 Regulates Basal Immune Components and Early Inducible Defense Responses in Arabidopsis. <i>Molecular Plant-Microbe Interactions</i> , 2015 , 28, 455-66	3.6	40
54	Loss of ceramide kinase in Arabidopsis impairs defenses and promotes ceramide accumulation and mitochondrial H ₂ O ₂ bursts. <i>Plant Cell</i> , 2014 , 26, 3449-67	11.6	58

53	Plant pathogenic bacteria target the actin microfilament network involved in the trafficking of disease defense components. <i>Bioarchitecture</i> , 2014 , 4, 149-53		8
52	Salicylic acid regulates Arabidopsis microbial pattern receptor kinase levels and signaling. <i>Plant Cell</i> , 2014 , 26, 4171-87	11.6	94
51	HopW1 from <i>Pseudomonas syringae</i> disrupts the actin cytoskeleton to promote virulence in Arabidopsis. <i>PLoS Pathogens</i> , 2014 , 10, e1004232	7.6	60
50	Salicylic acid signaling controls the maturation and localization of the arabidopsis defense protein ACCELERATED CELL DEATH6. <i>Molecular Plant</i> , 2014 , 7, 1365-1383	14.4	33
49	Accelerated cell death 2 suppresses mitochondrial oxidative bursts and modulates cell death in Arabidopsis. <i>Plant Journal</i> , 2012 , 69, 589-600	6.9	40
48	Type III secretion and effectors shape the survival and growth pattern of <i>Pseudomonas syringae</i> on leaf surfaces. <i>Plant Physiology</i> , 2012 , 158, 1803-18	6.6	48
47	SGT1b is required for HopZ3-mediated suppression of the epiphytic growth of <i>Pseudomonas syringae</i> on <i>N. benthamiana</i> . <i>Plant Signaling and Behavior</i> , 2012 , 7, 1129-31	2.5	2
46	A conserved cysteine motif is critical for rice ceramide kinase activity and function. <i>PLoS ONE</i> , 2011 , 6, e18079	3.7	12
45	<i>Pseudomonas syringae</i> hijacks plant stress chaperone machinery for virulence. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 13177-82	11.5	131
44	Genetic analysis of <i>acd6-1</i> reveals complex defense networks and leads to identification of novel defense genes in Arabidopsis. <i>Plant Journal</i> , 2009 , 58, 401-12	6.9	49
43	Priming in systemic plant immunity. <i>Science</i> , 2009 , 324, 89-91	33.3	611
42	Comparative large-scale analysis of interactions between several crop species and the effector repertoires from multiple pathovars of <i>Pseudomonas</i> and <i>Ralstonia</i> . <i>Plant Physiology</i> , 2009 , 150, 1733-49	6.6	81
41	Arabidopsis proteins important for modulating defense responses to <i>Pseudomonas syringae</i> that secrete HopW1-1. <i>Plant Journal</i> , 2008 , 54, 452-65	6.9	73
40	Signaling pathways that regulate the enhanced disease resistance of Arabidopsis "defense, no death" mutants. <i>Molecular Plant-Microbe Interactions</i> , 2008 , 21, 1285-96	3.6	66
39	Evolutionary dynamics of <i>Ralstonia solanacearum</i> . <i>Applied and Environmental Microbiology</i> , 2007 , 73, 1225-38	4.8	104
38	A J domain virulence effector of <i>Pseudomonas syringae</i> remodels host chloroplasts and suppresses defenses. <i>Current Biology</i> , 2007 , 17, 499-508	6.3	216
37	Whole-genome analysis to identify type III-secreted effectors. <i>Methods in Molecular Biology</i> , 2007 , 354, 19-34	1.4	5
36	A key role for the Arabidopsis WIN3 protein in disease resistance triggered by <i>Pseudomonas syringae</i> that secrete AvrRpt2. <i>Molecular Plant-Microbe Interactions</i> , 2007 , 20, 1192-200	3.6	56

35	Arabidopsis ACCELERATED CELL DEATH2 modulates programmed cell death. <i>Plant Cell</i> , 2006 , 18, 397-411	11.6	197
34	Identification of open reading frames unique to a select agent: <i>Ralstonia solanacearum</i> race 3 biovar 2. <i>Molecular Plant-Microbe Interactions</i> , 2006 , 19, 69-79	3.6	98
33	The type III effector repertoire of <i>Pseudomonas syringae</i> pv. <i>syringae</i> B728a and its role in survival and disease on host and non-host plants. <i>Molecular Microbiology</i> , 2006 , 62, 26-44	4.1	146
32	Bioinformatics correctly identifies many type III secretion substrates in the plant pathogen <i>Pseudomonas syringae</i> and the biocontrol isolate <i>P. fluorescens</i> SBW25. <i>Molecular Plant-Microbe Interactions</i> , 2005 , 18, 877-88	3.6	54
31	Degrade or die: a dual function for autophagy in the plant immune response. <i>Developmental Cell</i> , 2005 , 8, 799-801	10.2	11
30	Proposed guidelines for a unified nomenclature and phylogenetic analysis of type III Hop effector proteins in the plant pathogen <i>Pseudomonas syringae</i> . <i>Molecular Plant-Microbe Interactions</i> , 2005 , 18, 275-82	3.6	119
29	Structure-function analysis of the plasma membrane- localized Arabidopsis defense component ACD6. <i>Plant Journal</i> , 2005 , 44, 798-809	6.9	44
28	Divergent roles in <i>Arabidopsis thaliana</i> development and defense of two homologous genes, aberrant growth and death2 and AGD2-LIKE DEFENSE RESPONSE PROTEIN1, encoding novel aminotransferases. <i>Plant Cell</i> , 2004 , 16, 353-66	11.6	105
27	Free Radicals and Oxidative Stress 2004 , 203-214		2
26	The role and regulation of programmed cell death in plant-pathogen interactions. <i>Cellular Microbiology</i> , 2004 , 6, 201-11	3.9	582
25	A key role for ALD1 in activation of local and systemic defenses in Arabidopsis. <i>Plant Journal</i> , 2004 , 40, 200-12	6.9	141
24	The mitochondrion--an organelle commonly involved in programmed cell death in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2004 , 40, 596-610	6.9	221
23	ACD6, a novel ankyrin protein, is a regulator and an effector of salicylic acid signaling in the Arabidopsis defense response. <i>Plant Cell</i> , 2003 , 15, 2408-20	11.6	147
22	Identifying type III effectors of plant pathogens and analyzing their interaction with plant cells. <i>Current Opinion in Microbiology</i> , 2003 , 6, 20-8	7.9	157
21	Ceramides modulate programmed cell death in plants. <i>Genes and Development</i> , 2003 , 17, 2636-41	12.6	275
20	A functional screen for the type III (Hrp) secretome of the plant pathogen <i>Pseudomonas syringae</i> . <i>Science</i> , 2002 , 295, 1722-6	33.3	326
19	Functional analysis of the type III effectors AvrRpt2 and AvrRpm1 of <i>Pseudomonas syringae</i> with the use of a single-copy genomic integration system. <i>Molecular Plant-Microbe Interactions</i> , 2001 , 14, 145-55	3.6	126
18	The Arabidopsis aberrant growth and death2 mutant shows resistance to <i>Pseudomonas syringae</i> and reveals a role for NPR1 in suppressing hypersensitive cell death. <i>Plant Journal</i> , 2001 , 27, 203-11	6.9	115

17	A role for salicylic acid and NPR1 in regulating cell growth in Arabidopsis. <i>Plant Journal</i> , 2001 , 28, 209-166.9	126
16	The Arabidopsis-accelerated cell death gene ACD2 encodes red chlorophyll catabolite reductase and suppresses the spread of disease symptoms. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001 , 98, 771-776	11.5 256
15	Positive and negative regulation of salicylic acid-dependent cell death and pathogen resistance in Arabidopsis lsd6 and ssi1 mutants. <i>Molecular Plant-Microbe Interactions</i> , 2000 , 13, 877-81	3.6 16
14	Uncoupling salicylic acid-dependent cell death and defense-related responses from disease resistance in the Arabidopsis mutant acd5. <i>Genetics</i> , 2000 , 156, 341-50	4 174
13	The Gain-of-Function Arabidopsis acd6 Mutant Reveals Novel Regulation and Function of the Salicylic Acid Signaling Pathway in Controlling Cell Death, Defenses, and Cell Growth. <i>Plant Cell</i> , 1999 , 11, 1695	11.6 2
12	The gain-of-function Arabidopsis acd6 mutant reveals novel regulation and function of the salicylic acid signaling pathway in controlling cell death, defenses, and cell growth. <i>Plant Cell</i> , 1999 , 11, 1695-708	11.6 294
11	Differential expression of a senescence-enhanced metallothionein gene in Arabidopsis in response to isolates of <i>Peronospora parasitica</i> and <i>Pseudomonas syringae</i> . <i>Plant Journal</i> , 1998 , 16, 209-21	6.9 116
10	PROGRAMMED CELL DEATH IN PLANT-PATHOGEN INTERACTIONS. <i>Annual Review of Plant Biology</i> , 1997 , 48, 525-545	390
9	Programmed cell death: a way of life for plants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 12094-7	11.5 488
8	Programmed cell death in plants: a pathogen-triggered response activated coordinately with multiple defense functions. <i>Cell</i> , 1994 , 77, 551-63	56.2 612
7	Posttranscriptional repression of Escherichia coli OmpF protein in response to redox stress: positive control of the micF antisense RNA by the soxRS locus. <i>Journal of Bacteriology</i> , 1993 , 175, 1026-31	3.5 142
6	Arabidopsis mutants compromised for the control of cellular damage during pathogenesis and aging. <i>Plant Journal</i> , 1993 , 4, 327-41	6.9 255
5	Activation of oxidative stress genes by mutations at the soxQ/cfxB/marA locus of Escherichia coli. <i>Journal of Bacteriology</i> , 1991 , 173, 4433-9	3.5 114
4	Positive control of a global antioxidant defense regulon activated by superoxide-generating agents in Escherichia coli. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990 , 87, 6181-5	11.5 494
3	A global response induced in Escherichia coli by redox-cycling agents overlaps with that induced by peroxide stress. <i>Journal of Bacteriology</i> , 1989 , 171, 3933-9	3.5 218
2	Glutathione in Escherichia coli is dispensable for resistance to H ₂ O ₂ and gamma radiation. <i>Journal of Bacteriology</i> , 1986 , 168, 1026-9	3.5 134
1	Alkylation and oxidative damages to DNA: constitutive and inducible repair systems. <i>Basic Life Sciences</i> , 1986 , 39, 205-17	10