

Arnab Banerjee

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5848252/arnab-banerjee-publications-by-citations.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

46 papers	1,800 citations	24 h-index	42 g-index
47 ext. papers	1,906 ext. citations	5.1 avg, IF	4.28 L-index

#	Paper	IF	Citations
46	A naphthalene-based Al ³⁺ selective fluorescent sensor for living cell imaging. <i>Organic and Biomolecular Chemistry</i> , 2011 , 9, 5523-9	3.9	176
45	Rhodamine-based fluorescent probe for Al ³⁺ through time-dependent PET-CHEF-FRET processes and its cell staining application. <i>Inorganic Chemistry</i> , 2013 , 52, 3627-33	5.1	157
44	A naphthalene exciplex based Al ³⁺ selective on-type fluorescent probe for living cells at the physiological pH range: experimental and computational studies. <i>Analyst, The</i> , 2012 , 137, 2166-75	5	97
43	FRET based tri-color emissive rhodamine-pyrene conjugate as an Al ³⁺ selective colorimetric and fluorescence sensor for living cell imaging. <i>Dalton Transactions</i> , 2013 , 42, 13311-4	4.3	89
42	Ratiometric fluorescence sensing and intracellular imaging of Al ³⁺ ions driven by an intramolecular excimer formation of a pyrimidine-pyrene scaffold. <i>Dalton Transactions</i> , 2013 , 42, 4757-63	4.3	84
41	A coumarin-based "turn-on" fluorescent sensor for the determination of Al ³⁺ : single crystal X-ray structure and cell staining properties. <i>Dalton Transactions</i> , 2013 , 42, 10198-207	4.3	71
40	Thiophene anchored coumarin derivative as a turn-on fluorescent probe for Cr ³⁺ : cell imaging and speciation studies. <i>Talanta</i> , 2012 , 91, 18-25	6.2	71
39	An Al ³⁺ induced green luminescent fluorescent probe for cell imaging and naked eye detection. <i>Analytical Methods</i> , 2012 , 4, 1906	3.2	70
38	Nickel(II)-induced excimer formation of a naphthalene-based fluorescent probe for living cell imaging. <i>Inorganic Chemistry</i> , 2012 , 51, 5699-704	5.1	67
37	Exploiting Copper Redox for (19)F Magnetic Resonance-Based Detection of Cellular Hypoxia. <i>Journal of the American Chemical Society</i> , 2016 , 138, 2937-40	16.4	63
36	Antipyrine based arsenate selective fluorescent probe for living cell imaging. <i>Analytical Chemistry</i> , 2013 , 85, 1778-83	7.8	58
35	9-Acridone-4-carboxylic acid as an efficient Cr(III) fluorescent sensor: trace level detection, estimation and speciation studies. <i>Journal of Hazardous Materials</i> , 2011 , 188, 274-80	12.8	53
34	Fluorescence sensing of arsenate at nanomolar level in a greener way: naphthalene based probe for living cell imaging. <i>Chemical Communications</i> , 2013 , 49, 7231-3	5.8	49
33	Selective sensing of Hg ²⁺ using rhodamine-thiophene conjugate: red light emission and visual detection of intracellular Hg ²⁺ at nanomolar level. <i>Journal of Hazardous Materials</i> , 2013 , 261, 198-205	12.8	48
32	Hypoxia-Responsive F MRI Probes with Improved Redox Properties and Biocompatibility. <i>Inorganic Chemistry</i> , 2017 , 56, 6429-6437	5.1	47
31	Highly selective organic fluorescent probe for azide ion: formation of a "molecular ring". <i>Analyst, The</i> , 2012 , 137, 1544-6	5	47
30	A rhodamine-naphthalene conjugate as a FRET based sensor for Cr ³⁺ and Fe ³⁺ with cell staining application. <i>Analytical Methods</i> , 2013 , 5, 442-445	3.2	45

29	Interaction of soft donor sites with a hard metal ion: crystallographically characterized blue emitting fluorescent probe for Al(III) with cell staining studies. <i>RSC Advances</i> , 2012 , 2, 12447	3.7	44
28	2-(2-Pyridyl) benzimidazole based Co(II) complex as an efficient fluorescent probe for trace level determination of aspartic and glutamic acid in aqueous solution: a displacement approach. <i>Organic and Biomolecular Chemistry</i> , 2011 , 9, 7097-104	3.9	43
27	Methionine-pyrene hybrid based fluorescent probe for trace level detection and estimation of Hg(II) in aqueous environmental samples: experimental and computational studies. <i>Journal of Hazardous Materials</i> , 2011 , 186, 738-44	12.8	40
26	A rhodamine derivative as a "lock" and SCN ⁻ as a "key": visible light excitable SCN ⁻ sensing in living cells. <i>Chemical Communications</i> , 2013 , 49, 2527-9	5.8	38
25	Naphthalene based highly selective OFF-ON type fluorescent probe for Al ³⁺ and NO ₂ ⁻ ions for living cell imaging at physiological pH. <i>Inorganica Chimica Acta</i> , 2013 , 398, 64-71	2.7	30
24	Cd(II)-triggered excimer-monomer conversion of a pyrene derivative: time dependent red-shift of monomer emission with cell staining application. <i>Analyst</i> , 2012 , 137, 3910-3	5	29
23	A naphthalene-thiophene hybrid molecule as a fluorescent AND logic gate with Zn ²⁺ and OAc ⁻ ions as inputs: cell imaging and computational studies. <i>Dalton Transactions</i> , 2013 , 42, 6708-15	4.3	29
22	Interaction of a naphthalene based fluorescent probe with Al ³⁺ : experimental and computational studies. <i>Analytical Methods</i> , 2012 , 4, 3620	3.2	21
21	Thiophene anchored naphthalene derivative: Cr ³⁺ selective turn-on fluorescent probe for living cell imaging. <i>Analytical Methods</i> , 2012 , 4, 2254	3.2	21
20	Pyridine appended L-methionine: a novel chelating resin for pH dependent Cr speciation with scanning electron microscopic evidence and monitoring of yeast mediated green bio-reduction of Cr(VI) to Cr(III) in environmental samples. <i>Journal of Hazardous Materials</i> , 2011 , 185, 1448-57	12.8	20
19	A positron-emission tomography (PET)/magnetic resonance imaging (MRI) platform to track in vivo small extracellular vesicles. <i>Nanoscale</i> , 2019 , 11, 13243-13248	7.7	19
18	A FRET operated sensor for intracellular pH mapping: strategically improved efficiency on moving from an anthracene to a naphthalene derivative. <i>RSC Advances</i> , 2013 , 3, 14397	3.7	19
17	Structure, magnetism and catecholase activity of the first dicopper(II) complex having a single alkoxo bridge. <i>RSC Advances</i> , 2015 , 5, 10987-10993	3.7	18
16	An anion induced multisignaling probe for Hg(2+) and its application for fish kidney and liver tissue imaging studies. <i>Dalton Transactions</i> , 2015 , 44, 13186-95	4.3	18
15	Visible light excitable fluorescence probe and its functionalized Merrifield polymer: selective sensing and removal of arsenate from real samples. <i>RSC Advances</i> , 2014 , 4, 3887-3892	3.7	18
14	Xanthone based Pb ²⁺ selective turn on fluorescent probe for living cell staining. <i>Analytical Methods</i> , 2013 , 5, 169-172	3.2	13
13	Anthracene appended coumarin derivative as a Cr(III) selective turn-on fluorescent probe for living cell imaging: a green approach towards speciation studies. <i>Analytical Methods</i> , 2012 , 4, 3163	3.2	13
12	An INHIBIT logic gate from a thiophene derivative using iron and zinc ions as the input: tuning the efficiency on moving from naphthalene to anthracene to pyrene for the green luminescent detection of the intracellular iron. <i>Dalton Transactions</i> , 2013 , 42, 16387-95	4.3	12

11	Selective fluorescence and naked eye detection of histidine in aqueous medium via hydrogen bonding assisted Schiff base condensation. <i>Tetrahedron Letters</i> , 2014 , 55, 174-176	2	10
10	2-(2-Pyridyl) benzimidazole-based ternary Mn(II) complex as an arsenate selective turn-on fluorescence probe: ppb level determination and cell imaging studies. <i>New Journal of Chemistry</i> , 2014 , 38, 2744	3.6	9
9	A smart rhodamine-pyridine conjugate for bioimaging of thiocyanate in living cells. <i>RSC Advances</i> , 2015 , 5, 103350-103357	3.7	8
8	pH dependent separation of uranium by chelation chromatography using pyridine 2,6-dimethanol as a chelator: single crystal X-ray structural confirmation of the chelated uranium complex. <i>Journal of Hazardous Materials</i> , 2010 , 181, 154-60	12.8	8
7	A visible light excitable pyrene-naphthalene conjugate for ON fluorescence sensing of histidine in living cells. <i>RSC Advances</i> , 2014 , 4, 7495	3.7	7
6	Lighting of a rhodamine-based fluorescent lamp using ClO ₄ ⁻ as a connector: detection by the naked eye and cell imaging studies of trace amounts of ClO ₄ ⁻ ions. <i>RSC Advances</i> , 2013 , 3, 14044	3.7	7
5	Aluminum(III) induced green luminescence for naked eye detection: Experimental and computational studies. <i>Inorganica Chimica Acta</i> , 2014 , 412, 67-72	2.7	6
4	Spectroscopic Studies of a New Multi-Element Sensitive Fluorescent Probe Derived from 2-(2-pyridyl)benzimidazole: Selective Discrimination of Zn ²⁺ from Its Congeners. <i>Spectroscopy Letters</i> , 2013 , 46, 28-35	1.1	3
3	Tuning FRET efficiency as a novel approach for improved detection of naphthalene: application to environmental samples. <i>Journal of Molecular Recognition</i> , 2016 , 29, 303-7	2.6	3
2	pH-controlled solid-phase enrichment of Mn(II): confirmation of the structure of the extracted ternary Mn(II) complex by single crystal X-ray structure analysis. <i>Desalination and Water Treatment</i> , 2014 , 52, 6069-6078		1
1	THIOCYANATE-ASSISTED SOLID PHASE ENRICHMENT OF Cu(II) USING PYRIDINE DERIVATIVE WITH X-RAY CRYSTALLOGRAPHIC EVIDENCE. <i>Chemical Engineering Communications</i> , 2013 , 200, 638-654	2.2	1