Roberta Montisci

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/5850375/roberta-montisci-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

28 700 13 26 g-index

40 848 4.6 2.99 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
28	Multiparametric analysis of coronary flow in psoriasis using a coronary flow reserve companion. <i>European Journal of Clinical Investigation</i> , 2021 , e13711	4.6	O
27	Role of coronary microvascular dysfunction in heart failure with preserved ejection fraction. <i>Reviews in Cardiovascular Medicine</i> , 2021 , 22, 97-104	3.9	2
26	Surviving to Acute Myocardial Infarction: The Role of Psychological Factors and Alexithymia in Delayed Time to Searching Care: A Systematic Review. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	1
25	New insights to the potential mechanisms driving coronary flow reserve impairment in Cushingls syndrome: A pilot noninvasive study by transthoracic Doppler echocardiography. <i>Microvascular Research</i> , 2020 , 128, 103940	3.7	1
24	Long-term prognostic value of coronary flow reserve in psoriasis patients. <i>Atherosclerosis</i> , 2019 , 289, 57-63	3.1	7
23	Coronary flow reserve is related to the extension and transmurality of myocardial necrosis and predicts functional recovery after acute myocardial infarction. <i>Echocardiography</i> , 2019 , 36, 844-853	1.5	2
22	Assessment of the German and Italian Stress Cardiomyopathy Score for Risk Stratification for In-hospital Complications in Patients With Takotsubo Syndrome. <i>JAMA Cardiology</i> , 2019 , 4, 892-899	16.2	27
21	The Aim was About the Association with Psychiatric Disorders not on the Pathogenesis of Takotsubo - Authorts Reply. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2019 , 15, 5-7	3.2	1
20	Nailfold avascular score and coronary microvascular dysfunction in systemic sclerosis: A newsworthy association. <i>Autoimmunity Reviews</i> , 2019 , 18, 177-183	13.6	17
19	Coronary flow reserve in patients with primary biliary cholangitis. <i>Digestive and Liver Disease</i> , 2019 , 51, 542-548	3.3	
18	Coronary microvascular dysfunction may be related to IGF-1 in acromegalic patients and can be restored by therapy. <i>Atherosclerosis</i> , 2018 , 269, 100-105	3.1	6
17	Takotsubo Syndrome is Associated with Mood Disorders and Antidepressants Use, not with Anxiety and Impairment of Quality of Life Due to the Psychiatric Disorder. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2018 , 14, 26-32	3.2	13
16	Prognosis and first diagnostic ECG in STEMI patients referred to the emergency medical system for primary PCI. <i>Journal of Electrocardiology</i> , 2018 , 51, 1131-1134	1.4	3
15	Feasibility, symptoms, adverse effects, and complications associated with noninvasive assessment of coronary flow velocity reserve in women with suspected or known coronary artery disease. <i>Echocardiography</i> , 2017 , 34, 1324-1331	1.5	5
14	Role of cardiac dyssynchrony and resynchronization therapy in functional mitral regurgitation. <i>European Heart Journal Cardiovascular Imaging</i> , 2016 , 17, 471-80	4.1	31
13	Alexithymia affects the time from symptom onset to calling the emergency system in STEMI patients referred for primary PCI. <i>International Journal of Cardiology</i> , 2016 , 219, 428-32	3.2	8
12	Comment on Shore et al. Association between hyperglycemia at admission during hospitalization for acute myocardial infarction and subsequent diabetes: insights from the Veterans Administration Cardiac Care Follow-up Clinical Study. Diabetes Care 2014;37:409-418. <i>Diabetes Care</i>	14.6	1

LIST OF PUBLICATIONS

11	Prognostic impact of coronary microcirculation abnormalities in systemic sclerosis: a prospective study to evaluate the role of non-invasive tests. <i>Arthritis Research and Therapy</i> , 2013 , 15, R8	5.7	20
10	Impaired coronary flow reserve in young patients affected by severe psoriasis. <i>Atherosclerosis</i> , 2012 , 221, 113-7	3.1	42
9	Early noninvasive evaluation of coronary flow reserve after angioplasty in the left anterior descending coronary artery identifies patients at high risk of restenosis at follow-up. <i>Journal of the American Society of Echocardiography</i> , 2012 , 25, 902-10	5.8	8
8	Coronary microvascular dysfunction induced by primary hyperparathyroidism is restored after parathyroidectomy. <i>Circulation</i> , 2012 , 126, 1031-9	16.7	61
7	Early myocardial dysfunction after chronic use of anabolic androgenic steroids: combined pulsed-wave tissue Doppler imaging and ultrasonic integrated backscatter cyclic variations analysis. <i>Journal of the American Society of Echocardiography</i> , 2010 , 23, 516-22	5.8	15
6	Effect of a single IV administration of L-propionylcarnitine on myocardial microcirculation assessed by coronary flow velocity reserve measurement in patients with systemic sclerosis: a pilot study. <i>Clinical Therapeutics</i> , 2007 , 29, 163-71	3.5	5
5	Coronary flow velocity pattern and coronary flow reserve by contrast-enhanced transthoracic echocardiography predict long-term outcome in heart transplantation. <i>Circulation</i> , 2006 , 114, I49-55	16.7	64
4	Detection of coronary restenosis after coronary angioplasty by contrast-enhanced transthoracic echocardiographic Doppler assessment of coronary flow velocity reserve. <i>Journal of the American College of Cardiology</i> , 2002 , 40, 896-903	15.1	39
3	Reduced microvascular and myocardial damage in patients with acute myocardial infarction and preinfarction angina. <i>American Heart Journal</i> , 2002 , 144, 796-803	4.9	32
2	Contrast-enhanced transthoracic second harmonic echo Doppler with adenosine: a noninvasive, rapid and effective method for coronary flow reserve assessment. <i>Journal of the American College of Cardiology</i> , 1999 , 34, 122-30	15.1	131
1	Validation of a new noninvasive method (contrast-enhanced transthoracic second harmonic echo Doppler) for the evaluation of coronary flow reserve: comparison with intracoronary Doppler flow wire. <i>Journal of the American College of Cardiology</i> , 1999 , 34, 1193-200	15.1	158