

Signe Wiese

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5855083/publications.pdf>

Version: 2024-02-01

18
papers

522
citations

933447

10
h-index

888059

17
g-index

18
all docs

18
docs citations

18
times ranked

719
citing authors

#	ARTICLE	IF	CITATIONS
1	Cirrhotic cardiomyopathy: pathogenesis and clinical relevance. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2014, 11, 177-186.	17.8	205
2	Cardiac and proinflammatory markers predict prognosis in cirrhosis. <i>Liver International</i> , 2014, 34, e19-30.	3.9	65
3	Bile acids and cardiovascular function in cirrhosis. <i>Liver International</i> , 2017, 37, 1420-1430.	3.9	44
4	An update on cirrhotic cardiomyopathy. <i>Expert Review of Gastroenterology and Hepatology</i> , 2019, 13, 497-505.	3.0	33
5	Myocardial extracellular volume quantified by magnetic resonance is increased in cirrhosis and related to poor outcome. <i>Liver International</i> , 2018, 38, 1614-1623.	3.9	30
6	Diastolic dysfunction in cirrhosis. <i>Heart Failure Reviews</i> , 2016, 21, 599-610.	3.9	28
7	Cardiac dysfunction in cirrhosis: a 2-yr longitudinal follow-up study using advanced cardiac imaging. <i>American Journal of Physiology - Renal Physiology</i> , 2019, 317, G253-G263.	3.4	19
8	Cardiac imaging in patients with chronic liver disease. <i>Clinical Physiology and Functional Imaging</i> , 2017, 37, 347-356.	1.2	16
9	Fibrogenesis and inflammation contribute to the pathogenesis of cirrhotic cardiomyopathy. <i>Alimentary Pharmacology and Therapeutics</i> , 2020, 52, 340-350.	3.7	16
10	Total bile acid levels are associated with left atrial volume and cardiac output in patients with cirrhosis. <i>European Journal of Gastroenterology and Hepatology</i> , 2018, 30, 392-397.	1.6	13
11	Dobutamine reverses the cardio-suppressive effects of terlipressin without improving renal function in cirrhosis and ascites: a randomized controlled trial. <i>American Journal of Physiology - Renal Physiology</i> , 2020, 318, G313-G321.	3.4	10
12	Left atrial volume changes assessed by real time 3-dimensional echocardiography in relation to liver function and prognosis in patients with cirrhosis. <i>International Journal of Cardiovascular Imaging</i> , 2020, 36, 2121-2127.	1.5	10
13	Pronounced Coronary Arteriosclerosis in Cirrhosis: Influence on Cardiac Function and Survival?. <i>Digestive Diseases and Sciences</i> , 2018, 63, 1355-1362.	2.3	8
14	Cardiac Biomarkers in Cirrhosis and Portal Hypertension: Relation to Circulatory and Cardiac Dysfunction. , 2016, , 573-599.		7
15	Plasma ADAMTS-13 protein is not associated with portal hypertension or hemodynamic changes in patients with cirrhosis. <i>Digestive and Liver Disease</i> , 2016, 48, 404-408.	0.9	6
16	Low ascitic fluid total protein levels is not associated to the development of spontaneous bacterial peritonitis in a cohort of 274 patients with cirrhosis. <i>Scandinavian Journal of Gastroenterology</i> , 2018, 53, 200-205.	1.5	5
17	Few complications after paracentesis in patients with cirrhosis and refractory ascites. <i>Danish Medical Bulletin</i> , 2011, 58, A4212.	0.3	4
18	Dipeptidyl peptidase-3 is associated with severity of liver disease and circulatory complications in patients with cirrhosis. <i>Biomarkers</i> , 2022, 27, 196-204.	1.9	3