Alessandra Piras

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5856872/publications.pdf

Version: 2024-02-01

121 2,841 31 46
papers citations h-index g-index

121 121 3894
all docs docs citations times ranked citing authors

| # | Article | IF | CITATIONS |
|----|--|--------------------|----------------------|
| 1 | Chemical composition and in vitro bioactivity of the volatile and fixed oils of Nigella sativa L. extracted by supercritical carbon dioxide. Industrial Crops and Products, 2013, 46, 317-323. | 5.2 | 108 |
| 2 | Dyeing polyester fibres with disperse dyes in supercritical CO2. Dyes and Pigments, 2000, 45, 75-79. | 3.7 | 99 |
| 3 | Chemical Composition of the Essential Oil and Supercritical CO2Extract ofCommiphora myrrha(Nees) Engl. and ofAcorus calamusL Journal of Agricultural and Food Chemistry, 2005, 53, 7939-7943. | 5.2 | 99 |
| 4 | Supercritical CO ₂ Extract of Cinnamomum zeylanicum: Chemical Characterization and Antityrosinase Activity. Journal of Agricultural and Food Chemistry, 2007, 55, 10022-10027. | 5.2 | 97 |
| 5 | Chemical characterization and biological activity of essential oils from Daucus carota L. subsp. carota growing wild on the Mediterranean coast and on the Atlantic coast. Fìtoterapìâ, 2009, 80, 57-61. | 2.2 | 88 |
| 6 | Antifungal and anti-inflammatory potential of Lavandula stoechas and Thymus herba-barona essential oils. Industrial Crops and Products, 2013, 44, 97-103. | 5.2 | 86 |
| 7 | Comparative Analysis of the Oil and Supercritical CO2Extract ofElettaria cardamomum(L.) Maton. Journal of Agricultural and Food Chemistry, 2004, 52, 6278-6282. | 5. 2 | 68 |
| 8 | Coaxial microwave assisted hydrodistillation of essential oils from five different herbs (lavender,) Tj ETQq0 0 0 rg Food Science and Emerging Technologies, 2016, 33, 308-318. | gBT /Overlo 5.6 | ock 10 Tf 50 4 66 |
| 9 | 13C-CPMAS and 1H-NMR Study of the Inclusion Complexes of \hat{l}^2 -Cyclodextrin with Carvacrol, Thymol, and Eugenol Prepared in Supercritical Carbon Dioxide. Chemistry and Biodiversity, 2004, 1, 1354-1366. | 2.1 | 62 |
| 10 | Microalgae from domestic wastewater facility's high rate algal pond: Lipids extraction, characterization and biodiesel production. Bioresource Technology, 2016, 206, 239-244. | 9.6 | 59 |
| 11 | Extraction and Separation of Volatile and Fixed Oils from Seeds of <i>Myristica fragrans</i> by Supercritical CO ₂ : Chemical Composition and Cytotoxic Activity on Cacoâ€2 Cancer Cells. Journal of Food Science, 2012, 77, C448-53. | 3.1 | 58 |
| 12 | Antibacterial, allelopathic and antioxidant activities of essential oil of <i>Salvia officinalis </i> L. growing wild in the Atlas Mountains of Morocco. Natural Product Research, 2013, 27, 1673-1676. | 1.8 | 55 |
| 13 | Chemical and biological comparisons on supercritical extracts of <i>Tanacetum cinerariifolium</i> (Trevir) Sch. Bip. with three related species of chrysanthemums of Sardinia (Italy). Natural Product Research, 2009, 23, 190-199. | 1.8 | 54 |
| 14 | Novel configurations for a citrus waste based biorefinery: from solventless to simultaneous ultrasound and microwave assisted extraction. Green Chemistry, 2016, 18, 6482-6492. | 9.0 | 51 |
| 15 | Extraction of Oil from Wheat Germ by Supercritical CO2. Molecules, 2009, 14, 2573-2581. | 3 . 8 | 50 |
| 16 | Antioxidant activity of supercritical extract of Melissa of incinalissubsp. of incinalisand Melissa of incinalissubsp. inodora. Phytotherapy Research, 2004, 18, 789-792. | 5.8 | 49 |
| 17 | Extraction and Separation of Volatile and Fixed Oils from Berries of Laurus nobilis L. by Supercritical CO2. Molecules, 2008, 13, 1702-1711. | 3 . 8 | 45 |
| 18 | Chemical Composition and Antifungal Activity of Essential Oils and Supercritical CO2 Extracts of Apium nodiflorum (L.) Lag Mycopathologia, 2012, 174, 61-67. | 3.1 | 44 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Extraction and isolation of Pistacia lentiscus L. essential oil by supercritical CO2. Flavour and Fragrance Journal, 2002, 17, 239-244. | 2.6 | 43 |
| 20 | Ocimum tenuiflorum L. and Ocimum basilicum L., two spices of Lamiaceae family with bioactive essential oils. Industrial Crops and Products, 2018, 113, 89-97. | 5.2 | 43 |
| 21 | Extraction, separation and isolation of essential oils from natural matrices by supercritical CO2. Flavour and Fragrance Journal, 2003, 18, 505-509. | 2.6 | 42 |
| 22 | Antimicrobial activity of Inula helenium L. essential oil against Gram-positive and Gram-negative bacteria and Candida spp International Journal of Antimicrobial Agents, 2008, 31, 588-590. | 2.5 | 42 |
| 23 | Imazalil–cyclomaltoheptaose (β-cyclodextrin) inclusion complex: preparation by supercritical carbon dioxide and 13C CPMAS and 1H NMR characterization. Carbohydrate Research, 2003, 338, 2227-2232. | 2.3 | 38 |
| 24 | Antifungal activity of essential oil from <i>Mentha spicata</i> L. and <i>Mentha pulegium</i> L. growing wild in Sardinia island (Italy). Natural Product Research, 2021, 35, 993-999. | 1.8 | 38 |
| 25 | Chemical composition of Lycium europaeum fruit oil obtained by supercritical CO 2 extraction and evaluation of its antioxidant activity, cytotoxicity and cell absorption. Food Chemistry, 2017, 230, 82-90. | 8.2 | 37 |
| 26 | Chemical composition and biological assays of essential oils of <i>Calamintha nepeta </i> (L.) Savi subsp. <i>nepeta </i> (Lamiaceae). Natural Product Research, 2010, 24, 1734-1742. | 1.8 | 36 |
| 27 | Evaluation of Antioxidant Potential of "Maltese Mushroom―(Cynomorium coccineum) by Means of Multiple Chemical and Biological Assays. Nutrients, 2013, 5, 149-161. | 4.1 | 36 |
| 28 | Chemical composition and effect on intestinal Caco-2 cell viability and lipid profile of fixed oil from Cynomorium coccineum L Food and Chemical Toxicology, 2012, 50, 3799-3807. | 3.6 | 33 |
| 29 | Quality of wheat germ oil obtained by cold pressing and supercritical carbon dioxide extraction. Czech Journal of Food Sciences, 2013, 31, 236-240. | 1.2 | 32 |
| 30 | Extraction of Juniperus communis L. ssp.nana Willd. essential oil by supercritical carbon dioxide. Flavour and Fragrance Journal, 2006, 21, 148-154. | 2.6 | 31 |
| 31 | Extraction of Santalum album and Boswellia carterii Birdw. volatile oil by supercritical carbon dioxide: influence of some process parameters. Flavour and Fragrance Journal, 2006, 21, 718-724. | 2.6 | 31 |
| 32 | Supercritical CO ₂ Extraction of Waste Citrus Seeds: Chemical Composition, Nutritional and Biological Properties of Edible Fixed Oils. European Journal of Lipid Science and Technology, 2019, 121, 1800502. | 1.5 | 31 |
| 33 | Comparative analysis of the oil and supercritical CO2extract of Cymbopogon citratus Stapf Natural Product Research, 2006, 20, 455-459. | 1.8 | 30 |
| 34 | Excess enthalpy and excess volume for binary systems of two ionic liquidsÂ+Âwater. Journal of Thermal Analysis and Calorimetry, 2011, 103, 29-33. | 3.6 | 30 |
| 35 | Isolation of the volatile fraction from <i> Apium graveolens</i> L. (Apiaceae) by supercritical carbon dioxide extraction and hydrodistillation: Chemical composition and antifungal activity. Natural Product Research, 2013, 27, 1521-1527. | 1.8 | 30 |
| 36 | Extraction, Separation and Isolation of Volatiles and Dyes from <i>Calendula officinalis </i> L. and <i>Aloysia triphylla </i> (L'Her.) Britton by Supercritical CO ₂ . Journal of Essential Oil Research, 2003, 15, 350-355. | 2.7 | 28 |

| # | Article | IF | Citations |
|----|--|--------------|-----------|
| 37 | Comparative Analysis of Supercritical CO ₂ Extract and Oil of <i>Pimenta dioica</i> Leaves. Journal of Essential Oil Research, 2005, 17, 530-532. | 2.7 | 28 |
| 38 | Isolation of <i> Crithmum maritimum </i> L. volatile oil by supercritical carbon dioxide extraction and biological assays. Natural Product Research, 2007, 21, 1145-1150. | 1.8 | 28 |
| 39 | Biological activity evaluation of the oils from <i>Laurus nobilis</i> of Tunisia and Algeria extracted by supercritical carbon dioxide. Natural Product Research, 2009, 23, 230-237. | 1.8 | 28 |
| 40 | Essential oil composition and variability of <i>Laurus nobilis</i> L. growing in Tunisia, comparison and chemometric investigation of different plant organs. Natural Product Research, 2009, 23, 343-354. | 1.8 | 25 |
| 41 | Chemical composition and antioxidant activity of essential oil from aerial parts of <i>Teucrium flavum</i> L. subsp. <i>flavum</i> growing spontaneously in Tunisia. Natural Product Research, 2015, 29, 2336-2340. | 1.8 | 25 |
| 42 | Supercritical CO ₂ extract and essential oil of aerial part of <i>Ledum palustre</i> L. – Chemical composition and anti-inflammatory activity. Natural Product Research, 2015, 29, 999-1005. | 1.8 | 24 |
| 43 | In vitro antimicrobial, antioxidant and antiviral activities of the essential oil and various extracts of wild (Daucus virgatus (Poir.) Maire) from Tunisia. Industrial Crops and Products, 2017, 109, 109-115. | 5 . 2 | 24 |
| 44 | Analysis of the Volatile Concentrate of the Leaves and Flowers of <i>Helichrysum italicum </i> (Roth) Don ssp. <i>microphyllum </i> (Willd.) Nyman <i> (Asteraceae) </i> by Supercritical Fluid Extraction and Their Essential Oils. Journal of Essential Oil Research, 2003, 15, 120-126. | 2.7 | 23 |
| 45 | Excess enthalpies of aromatic ether or aromatic ketone(1)+n-heptane(2) mixtures DISQUAC analysis. Journal of Thermal Analysis and Calorimetry, 2008, 92, 137-144. | 3.6 | 21 |
| 46 | Maltese Mushroom (Cynomorium coccineum L.) as Source of Oil with Potential Anticancer Activity. Nutrients, 2015, 7, 849-864. | 4.1 | 21 |
| 47 | Anti-inflammatory activity of Pistacia lentiscus essential oil: involvement of IL-6 and TNF-alpha. Natural Product Communications, 2011, 6, 1543-4. | 0.5 | 21 |
| 48 | Comparative analysis of the oil and supercritical CO2extract of Artemisia arborescens L. and Helichrysum splendidum (Thunb.) Less Natural Product Research, 2006, 20, 421-428. | 1.8 | 19 |
| 49 | Comparative analysis of the oil and supercritical CO2extract ofRidolfia segetum(L.) Moris. Natural Product Research, 2007, 21, 412-417. | 1.8 | 19 |
| 50 | Phytochemical composition and the cholinesterase and xanthine oxidase inhibitory properties of seed extracts from the <i>Washingtonia filifera</i> palm fruit. RSC Advances, 2019, 9, 21278-21287. | 3.6 | 19 |
| 51 | A comparative study of thermodynamic properties of binary mixtures containing dimethylsulfoxide. Journal of Thermal Analysis and Calorimetry, 2007, 90, 909-922. | 3.6 | 18 |
| 52 | Chemical composition and antifungal activity of supercritical extract and essential oil of <i>Tanacetum vulgare</i> growing wild in Lithuania. Natural Product Research, 2014, 28, 1906-1909. | 1.8 | 18 |
| 53 | Effect of pressure variation on the efficiency of supercritical fluid extraction of wild carrot (Daucus carota subsp. maritimus) extracts. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2019, 1125, 121713. | 2.3 | 18 |
| 54 | Anti-Aging Potential of Extracts from Washingtonia filifera Seeds. Plants, 2021, 10, 151. | 3.5 | 18 |

| # | Article | IF | Citations |
|----|---|-----|-----------|
| 55 | Chemical composition and antioxidant activity of the essential oil of (i>Juniperus phoenicea L. berries. Natural Product Research, 2011, 25, 1695-1706. | 1.8 | 17 |
| 56 | Supercritical CO ₂ extraction of volatile oils from Sardinian <i>Foeniculum vulgare</i> ssp. <i>vulgare</i> 813-1825. Product Research, 2014, 28, 1819-1825. | 1.8 | 17 |
| 57 | Extraction, Separation and Isolation of Volatiles and Dyes from <i>Calendula officinalis</i> L. and <i>Aloysia triphylla</i> (L'Her.) Britton by Supercritical CO ₂ . Journal of Essential Oil Research, 2003, 15, 272-277. | 2.7 | 16 |
| 58 | Comparative Analysis of the Oil and Supercritical CO ₂ Extract of <i>Ferula communis</i> L Journal of Essential Oil Research, 2005, 17, 150-152. | 2.7 | 16 |
| 59 | Isolation of Seseli bocconi Guss., subsp. praecox Gamisans (Apiaceae) volatile oil by supercritical carbon dioxide extraction. Natural Product Research, 2006, 20, 820-826. | 1.8 | 16 |
| 60 | Essential oil composition of leaves of Stachys yemenensis obtained by supercritical CO2. Natural Product Research, 2010, 24, 1823-1829. | 1.8 | 16 |
| 61 | Antifungal activity of extracts fromCynomorium coccineumgrowing wild in Sardinia island (Italy). Natural Product Research, 2015, 29, 2247-2250. | 1.8 | 16 |
| 62 | Chemical variability in essential oils from <i>Ruta</i> species among seasons, and its taxonomic and ecological significance. Natural Product Research, 2017, 31, 2329-2334. | 1.8 | 16 |
| 63 | Chemical Composition and Antioxidant Potential Differences between Cynomorium coccineum L. Growing in Italy and in Tunisia: Effect of Environmental Stress. Diversity, 2018, 10, 53. | 1.7 | 16 |
| 64 | A comparative study of thermodynamic properties of binary mixtures containing perfluoroalkanes. Journal of Thermal Analysis and Calorimetry, 2008, 92, 145-154. | 3.6 | 15 |
| 65 | Comparison of the antimicrobial activity and the essential oil composition of Juniperus oxycedrus subsp. rufescens obtained by hydrodistillation and supercritical carbon dioxide extraction methods. Chemistry of Natural Compounds, 2009, 45, 739-741. | 0.8 | 15 |
| 66 | Antifungal activity and chemical composition of essential oils from <i>Smyrnium olusatrum</i> L. (Apiaceae) from Italy and Portugal. Natural Product Research, 2012, 26, 993-1003. | 1.8 | 15 |
| 67 | Chemical composition of the essential oils of the berries of <i>Juniperus oxycedrus </i> L. ssp. <i>rufescens </i> (L. K.) and <i>Juniperus oxycedrus </i> L. ssp. <i>macrocarpa </i> (S. & amp; m.) Ball. and their antioxidant activities. Natural Product Research, 2012, 26, 810-820. | 1.8 | 15 |
| 68 | Chemical composition and biological activity of Tanacetum audibertii (Req.) DC. (Asteraceae), an endemic species of Sardinia Island, Italy. Industrial Crops and Products, 2015, 65, 472-476. | 5.2 | 15 |
| 69 | Extraction, separation and isolation of volatiles from Vitex agnus-castus L. (Verbenaceae) wild species of Sardinia, Italy, by supercritical CO2. Natural Product Research, 2010, 24, 569-579. | 1.8 | 14 |
| 70 | Essential Oil Constituents and Antioxidant Activity of <i>Asplenium </i> Ferns. Journal of Chromatographic Science, 2016, 54, 1341-1345. | 1.4 | 14 |
| 71 | Antioxidant activity of supercritical carbon dioxide extracts of Salvia desoleana on two human endothelial cell models. Food Research International, 2012, 46, 354-359. | 6.2 | 13 |
| 72 | Fatty acid composition and antioxidant activity of Pistacia lentiscus L. fruit fatty oil from Algeria. Journal of Food Measurement and Characterization, 2018, 12, 1408-1412. | 3.2 | 13 |

| # | Article | IF | Citations |
|----|---|-----|-----------|
| 73 | Isolation of (i) Juniperus phoenicea (i) Volatiles by Supercritical Carbon Dioxide Extraction and Bioactivity Assays. Journal of Essential Oil Research, 2004, 16, 256-261. | 2.7 | 12 |
| 74 | Anti-inflammatory Activity of <i>Pistacia lentiscus</i> Essential Oil: Involvement of IL-6 and TNF-α. Natural Product Communications, 2011, 6, 1934578X1100601. | 0.5 | 12 |
| 75 | Extraction of the volatile oil from <i>Carum carvi</i> of Tunisia and Lithuania by supercritical carbon dioxide: chemical composition and antiulcerogenic activity. Natural Product Research, 2013, 27, 2132-2136. | 1.8 | 12 |
| 76 | Chemical composition, antibacterial and antioxidant activities of essential oils from flowers, leaves and aerial parts of Tunisian <i>Dittrichia Viscosa</i> . Journal of Essential Oil Research, 2019, 31, 582-589. | 2.7 | 12 |
| 77 | Excess enthalpies of chloroalkylbenzene+n-heptane or +cyclohexane mixtures. Journal of Thermal Analysis and Calorimetry, 2008, 91, 37-46. | 3.6 | 11 |
| 78 | Mugil cephalus roe oil obtained by supercritical fluid extraction affects the lipid profile and viability in cancer HeLa and B16F10 cells. Food and Function, 2016, 7, 4092-4103. | 4.6 | 11 |
| 79 | Chemical characterisation and biological activity of leaf essential oils obtained from Pistacia terebinthus growing wild in Tunisia and Sardinia Island. Natural Product Research, 2017, 31, 2684-2689. | 1.8 | 11 |
| 80 | Antifungal activity and chemical composition of the essential oil from the aerial parts of two new <i>Teucrium capitatum </i> L. chemotypes from Sardinia Island, Italy. Natural Product Research, 2021, 35, 6007-6013. | 1.8 | 10 |
| 81 | Supercritical extraction of volatile and fixed oils from <i>Petroselinum crispum</i> L. seeds: chemical composition and biological activity. Natural Product Research, 2022, 36, 1883-1888. | 1.8 | 10 |
| 82 | Essential Oils of Daucus carota subsp. carota of Tunisia Obtained by Supercritical Carbon Dioxide Extraction. Natural Product Communications, 2010, 5, 1934578X1000501. | 0.5 | 9 |
| 83 | Chemical Polymorphism of Essential Oils from Populations of <i>Laurus nobilis</i> Grown on Tunisia, Algeria and France. Natural Product Communications, 2011, 6, 1934578X1100601. | 0.5 | 9 |
| 84 | Ethylammonium alkanoate-based ionic liquidÂ+Âwater mixtures. Journal of Thermal Analysis and Calorimetry, 2015, 121, 1129-1137. | 3.6 | 9 |
| 85 | Isolation of the volatile oil from Satureja thymbra by supercritical carbon dioxide extraction: chemical composition and biological activity. Natural Product Communications, 2011, 6, 1523-6. | 0.5 | 9 |
| 86 | Thermodynamic properties of binary mixtures containing oxaalkanes. Journal of Thermal Analysis and Calorimetry, 2009, 95, 149-159. | 3.6 | 8 |
| 87 | Chemical and biomolecular analyses to discriminate three taxa of Pistacia genus from Sardinia Island (Italy) and their antifungal activity. Natural Product Research, 2018, 32, 2766-2774. | 1.8 | 8 |
| 88 | Bovine Viral Diarrhea Virus (BVDV): A Preliminary Study on Antiviral Properties of Some Aromatic and Medicinal Plants. Pathogens, 2021, 10, 403. | 2.8 | 8 |
| 89 | Chemical composition and biological activity of essential oil of <i>Teucrium scordium </i> L. subsp. <i>scordioides </i> (Schreb.) Arcang. (Lamiaceae) from Sardinia Island (Italy). Natural Product Research, 2021, , 1-8. | 1.8 | 8 |
| 90 | A comparative study of thermodynamic properties of binary mixtures containing alkynes. Thermochimica Acta, 2004, 418, 85-93. | 2.7 | 7 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 91 | Excess enthalpies of mixtures of mono-carboxylic acid with dibutylether. Journal of Thermal Analysis and Calorimetry, 2012, 108, 777-782. | 3.6 | 7 |
| 92 | Antiproliferative and antiviral activity of methanolic extracts from Sardinian Maltese Mushroom (Cynomorium coccineum L.). Natural Product Research, 2019, 35, 1-5. | 1.8 | 7 |
| 93 | Phytotoxic effects of Salvia rosmarinus essential oil on Acacia saligna seedling growth. Flora: Morphology, Distribution, Functional Ecology of Plants, 2020, 269, 151639. | 1.2 | 7 |
| 94 | Flavoring of sea salt with Mediterranean aromatic plants affects salty taste perception. Journal of the Science of Food and Agriculture, 2022, 102, 6005-6013. | 3.5 | 7 |
| 95 | Isolation of <i>Guaiacum Bulnesia </i> Volatile Oil by Supercritical Carbon Dioxide Extraction. Journal of Essential Oil-bearing Plants: JEOP, 2007, 10, 221-228. | 1.9 | 6 |
| 96 | Composition and Biological Activity of Supercritical CO2Extract of Some Lamiaceae Growing Wild in Sardinia (Italy). Journal of Essential Oil-bearing Plants: JEOP, 2010, 13, 625-632. | 1.9 | 6 |
| 97 | Comparative analysis of the oil and supercritical CO2extract of Schinus molleL. growing in Yemen. Natural Product Research, 2011, 25, 1366-1369. | 1.8 | 6 |
| 98 | Characterization of Essential Oils from Different Taxa Belonging to the Genus Teucrium in Sardinia Island, Italy. Plants, 2021, 10, 1359. | 3.5 | 6 |
| 99 | Extraction of Lantana camaraessential oil by supercritical carbon dioxide: influence of the grinding and biological activity. Natural Product Research, 2007, 21, 33-36. | 1.8 | 5 |
| 100 | Calorimetric study of nitro group/solvent interactions. Journal of Thermal Analysis and Calorimetry, 2010, 99, 1015-1023. | 3.6 | 5 |
| 101 | Calorimetric Study of Nitrile Groupâ^'Solvent Interactions and Comparison with Dispersive Quasi-Chemical (DISQUAC) Predictions. Journal of Chemical & Engineering Data, 2010, 55, 5406-5412. | 1.9 | 5 |
| 102 | Fatty acids from high rate algal pond's microalgal biomass and osmotic stress effects. Bioresource Technology, 2017, 244, 860-864. | 9.6 | 5 |
| 103 | Evaluation of the Antioxidant and Cytotoxic Activities on Cancer Cell Line of Extracts of Parasitic Plants Harvested in Tunisia. Polish Journal of Food and Nutrition Sciences, 0, , 253-263. | 1.7 | 5 |
| 104 | Supercritical CO2extract from needles ofPinus nigrassp.laricio: combined analysis by GC, GC–MS and13C NMR. Natural Product Research, 2007, 21, 834-837. | 1.8 | 4 |
| 105 | Morphological, Chemical and Antibacterial Characteristics of Laurus nobilis L. Growing in Tunisia. Asian Journal of Chemistry, 2015, 27, 3838-3842. | 0.3 | 4 |
| 106 | Excess enthalpies and excess volumes of binary mixtures containing a linear carboxylic acidÂ+Âdi-iso-propyl ether at 298.15ÂK and 0.1ÂMPa. Journal of Thermal Analysis and Calorimetry, 2016, 125, 607-615. | 3.6 | 4 |
| 107 | Evaluation of antioxidant and tyrosinase inhibitory activities ofÂthe extracts of <i>Sarcopoterium spinosum</i> (L.) Spach fruits. Natural Product Research, 2017, 31, 2900-2904. | 1.8 | 4 |
| 108 | Inhibitory effect of rosemary essential oil, loaded in liposomes, on seed germination of <i>Acacia saligna</i> , an invasive species in Mediterranean ecosystems. Botany, 2019, 97, 283-291. | 1.0 | 4 |

| # | Article | IF | CITATIONS |
|-----|--|------------|-------------|
| 109 | Waste salt from the manufacturing process of mullet bottarga as source of oil with nutritional and nutraceutical properties. Journal of the Science of Food and Agriculture, 2020, 100, 5363-5372. | 3.5 | 4 |
| 110 | Seasonal Variation in the Chemical Profile, Antifungal and Insecticidal Activities of Essential Oils from Daucus reboudii. Waste and Biomass Valorization, 2022, 13, 1859-1871. | 3.4 | 4 |
| 111 | Fatty Acid Composition, Essential Oil and Antibacterial Activity of Berries ofLaurus nobilisL Journal of Essential Oil-bearing Plants: JEOP, 2009, 12, 422-434. | 1.9 | 3 |
| 112 | Evaluation of the attractant effect and lipid profile modulation of natural fixed oils on the medfly <i>Ceratitis capitata</i> (Wiedemann). Archives of Insect Biochemistry and Physiology, 2018, 99, e21508. | 1.5 | 3 |
| 113 | Comparative evaluation of the composition of vegetable essential and fixed oils obtained by supercritical extraction and conventional techniques: a chemometric approach. International Journal of Food Science and Technology, 2021, 56, 4496-4505. | 2.7 | 3 |
| 114 | Extraction of volatile fractions and carotenoids from orange and kumquat peel by supercritical carbon dioxide. Journal of Essential Oil-bearing Plants: JEOP, 2003, 6, 86-96. | 1.9 | 2 |
| 115 | Chemical Characterization and Evaluation of Biological Activity of Essential Oil ofJuniperus phoeniceaof Tunisia. Journal of Essential Oil-bearing Plants: JEOP, 2008, 11, 233-241. | 1.9 | 2 |
| 116 | Thermodynamic properties of binary mixtures containing oxaalkanes. Journal of Thermal Analysis and Calorimetry, 2009, 97, 817-825. | 3.6 | 2 |
| 117 | Excess Enthalpies of Chloroalkylbenzene + Alkylbenzene Mixtures. Journal of Chemical & Engineering Data, 2007, 52, 1941-1945. | 1.9 | 1 |
| 118 | Calorimetric study of the selected nitroalkane+chloroalkane binary systems. Journal of Thermal Analysis and Calorimetry, 2014, 116, 119-127. | 3.6 | 1 |
| 119 | Chemical Composition of Essential Oils from Needles of Pinus pinaster from Italy and Tunisia. Asian Journal of Chemistry, 2015, 27, 2662-2664. | 0.3 | 1 |
| 120 | Composition, Insecticidal and Antifungal Activities of Tunisian Daucus setifolius Essential Oil. Waste and Biomass Valorization, $0, 1$. | 3.4 | 1 |
| 121 | Excess enthalpies of [CnMIM][NTf2] n = (2 or 10) + ethanol or + N-methyl-2-pyrrolido 298.15ÂK and 0.1ÂMPa. Journal of Thermal Analysis and Calorimetry, 0, , 1. | ong binary | mixtures at |