

# Hui Yuan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5862786/publications.pdf>

Version: 2024-02-01

24  
papers

645  
citations

932766

10  
h-index

610482

24  
g-index

24  
all docs

24  
docs citations

24  
times ranked

1202  
citing authors

#	ARTICLE	IF	CITATIONS
1	Galectin-3 as a novel biomarker for disease diagnosis and a target for therapy (Review). <i>International Journal of Molecular Medicine</i> , 2018, 41, 599-614.	1.8	210
2	Microarray profile of circular RNAs identifies hsa_circ_0014130 as a new circular RNA biomarker in non-small cell lung cancer. <i>Scientific Reports</i> , 2018, 8, 2878.	1.6	130
3	Magnitude of Soluble ST2 as a Novel Biomarker for Acute Aortic Dissection. <i>Circulation</i> , 2018, 137, 259-269.	1.6	80
4	Circulating microRNA signature for the diagnosis of childhood dilated cardiomyopathy. <i>Scientific Reports</i> , 2018, 8, 724.	1.6	37
5	Circular RNA as a biomarker for cancer: A systematic meta-analysis. <i>Oncology Letters</i> , 2018, 16, 4078-4084.	0.8	28
6	Ligustrazine Protects Homocysteine-Induced Apoptosis in Human Umbilical Vein Endothelial Cells by Modulating Mitochondrial Dysfunction. <i>Journal of Cardiovascular Translational Research</i> , 2019, 12, 591-599.	1.1	21
7	Elevated soluble ST2 concentration may involve in the progression of atrial fibrillation. <i>Clinica Chimica Acta</i> , 2018, 480, 138-142.	0.5	20
8	Epidemiology and antifungal susceptibilities of yeast isolates causing invasive infections across urban Beijing, China. <i>Future Microbiology</i> , 2017, 12, 1075-1086.	1.0	14
9	Relationship of HLA-B*51 and HLA-B*52 alleles and TNF- $\pm$ -308A/G polymorphism with susceptibility to Takayasu arteritis: a meta-analysis. <i>Clinical Rheumatology</i> , 2017, 36, 173-181.	1.0	14
10	Association of Soluble ST2 Serum Levels With Outcomes in Pediatric Dilated Cardiomyopathy. <i>Canadian Journal of Cardiology</i> , 2019, 35, 727-735.	0.8	14
11	Association of FCGR2A/FCGR3A variant rs2099684 with Takayasu arteritis in the Han Chinese population. <i>Oncotarget</i> , 2017, 8, 17239-17245.	0.8	11
12	Single nucleotide polymorphisms of IL12B are associated with Takayasu arteritis in Chinese Han population. <i>Rheumatology International</i> , 2017, 37, 547-555.	1.5	9
13	Platelets in Acute Coronary Syndrome Patients with High Platelet Reactivity after Dual Antiplatelet Therapy Exhibit Upregulation of miR-204-5p. <i>Annals of Clinical and Laboratory Science</i> , 2019, 49, 619-631.	0.2	7
14	Association between genetic variants in the human leukocyte antigen- $\beta$ -MICA and Takayasu arteritis in Chinese Han population. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 271-277.	0.9	6
15	Non-fasting lipids detection and their significance in pregnant women. <i>Lipids in Health and Disease</i> , 2019, 18, 96.	1.2	6
16	The relationships of serum homocysteine levels and traditional lipid indicators with disease activity and coronary artery involvement in Takayasu arteritis. <i>Immunologic Research</i> , 2020, 68, 405-413.	1.3	6
17	Single nucleotide polymorphisms in the ETS1 gene are associated with idiopathic inflammatory myopathies in a northern Chinese Han population. <i>Scientific Reports</i> , 2017, 7, 13128.	1.6	5
18	Serum C1q concentration is associated with disease activity in Chinese Takayasu arteritis patients: A case-control study. <i>Health Science Reports</i> , 2021, 4, e252.	0.6	5

#	ARTICLE	IF	CITATIONS
19	Serum concentrations of small dense low-density lipoprotein cholesterol and lipoprotein(a) are related to coronary arteriostenosis in Takayasu arteritis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e23966.	0.9	5
20	Association between the BANK1 rs3733197 polymorphism and polymyositis/dermatomyositis in a Chinese Han population. <i>Clinical Rheumatology</i> , 2019, 38, 431-436.	1.0	4
21	Abnormal brain activity in rats with sustained hypobaric hypoxia exposure. <i>Chinese Medical Journal</i> , 2019, 132, 2621-2627.	0.9	4
22	Homozygous familial hypercholesterolemia in China: Genetic and clinical characteristics from a real-world, multi-center, cohort study. <i>Journal of Clinical Lipidology</i> , 2022, 16, 306-314.	0.6	4
23	Identification of novel genes in Behcet's disease using integrated bioinformatic analysis. <i>Immunologic Research</i> , 2022, 70, 461-468.	1.3	3
24	Correlation Between Extended Leukocyte Differential Count and Coronary Artery Disease. <i>Journal of Cardiovascular Pharmacology</i> , 2018, 71, 359-366.	0.8	2