

# Qin Jin

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5863508/publications.pdf>

Version: 2024-02-01

21  
papers

455  
citations

687220

13  
h-index

713332

21  
g-index

21  
all docs

21  
docs citations

21  
times ranked

759  
citing authors

#	ARTICLE	IF	CITATIONS
1	Role of M2 Macrophages in Sepsis-Induced Acute Kidney Injury. <i>Shock</i> , 2018, 50, 233-239.	1.0	52
2	Decellularized breast matrix as bioactive microenvironment for in vitro three-dimensional cancer culture. <i>Journal of Cellular Physiology</i> , 2019, 234, 3425-3435.	2.0	49
3	Human breast cancer decellularized scaffolds promote epithelial-mesenchymal transitions and stemness of breast cancer cells in vitro. <i>Journal of Cellular Physiology</i> , 2019, 234, 9447-9456.	2.0	39
4	Effects of Rab27A and Rab27B on Invasion, Proliferation, Apoptosis, and Chemoresistance in Human Pancreatic Cancer Cells. <i>Pancreas</i> , 2017, 46, 1173-1179.	0.5	30
5	High Tmprss11d protein expression predicts poor overall survival in non-small cell lung cancer. <i>Oncotarget</i> , 2017, 8, 12812-12819.	0.8	29
6	High Eg5 expression predicts poor prognosis in breast cancer. <i>Oncotarget</i> , 2017, 8, 62208-62216.	0.8	29
7	Aldehyde dehydrogenase 1 expression is correlated with poor prognosis in breast cancer. <i>Medicine (United States)</i> , 2017, 96, e7171.	0.4	28
8	High kinesin family member 11 expression predicts poor prognosis in patients with clear cell renal cell carcinoma. <i>Journal of Clinical Pathology</i> , 2019, 72, 354-362.	1.0	28
9	LAMP1 expression is associated with poor prognosis in breast cancer. <i>Oncology Letters</i> , 2017, 14, 4729-4735.	0.8	27
10	Triple-amiRNA VEGFRs inhibition in pancreatic cancer improves the efficacy of chemotherapy through EMT regulation. <i>Journal of Controlled Release</i> , 2017, 245, 1-14.	4.8	27
11	EphA8 is a prognostic marker for epithelial ovarian cancer. <i>Oncotarget</i> , 2016, 7, 20801-20809.	0.8	20
12	Plasma miR-185 is decreased in patients with esophageal squamous cell carcinoma and might suppress tumor migration and invasion by targeting RAGE. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 309, G719-G729.	1.6	19
13	Ameliorative effect of recombinant human lactoferrin on the premature ovarian failure in rats after cyclophosphamide treatments. <i>Journal of Ovarian Research</i> , 2021, 14, 17.	1.3	16
14	Downregulated SASH1 expression indicates poor clinical prognosis in gastric cancer. <i>Human Pathology</i> , 2018, 74, 83-91.	1.1	14
15	Reduced expression of EphA5 is associated with lymph node metastasis, advanced TNM stage, and poor prognosis in colorectal carcinoma. <i>Histology and Histopathology</i> , 2017, 32, 491-497.	0.5	13
16	Overexpression of FIBCD1 Is Predictive of Poor Prognosis in Gastric Cancer. <i>American Journal of Clinical Pathology</i> , 2018, 149, 474-483.	0.4	10
17	High expression of PFTK1 in cancer cells predicts poor prognosis in colorectal cancer. <i>Molecular Medicine Reports</i> , 2017, 16, 224-230.	1.1	8
18	High methionyl-tRNA synthetase expression predicts poor prognosis in patients with breast cancer. <i>Journal of Clinical Pathology</i> , 2020, 73, 803-812.	1.0	8

#	ARTICLE	IF	CITATIONS
19	Association of High Expression of Gro $\beta$ 2 with Clinical and Pathological Characteristics of Unfavorable Prognosis in Gastrointestinal Stromal Tumors. <i>Disease Markers</i> , 2015, 2015, 1-8.	0.6	6
20	High Spy1 expression predicts poor prognosis in colorectal cancer. <i>Cancer Management and Research</i> , 2018, Volume 10, 2757-2765.	0.9	2
21	High expression of EphA4 is associated with invasion and lymph node metastasis in colorectal carcinomas. <i>International Journal of Clinical and Experimental Pathology</i> , 2017, 10, 9697-9703.	0.5	1