

Andres Redondo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5868718/publications.pdf>

Version: 2024-02-01

138
papers

7,763
citations

230014

27
h-index

62345

84
g-index

145
all docs

145
docs citations

145
times ranked

10535
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Prognostic implications of tumor-infiltrating T cells in early-stage endometrial cancer. <i>Modern Pathology</i> , 2022, 35, 256-265. | 2.9 | 12 |
| 2 | Olaparib as first line in BRCA-mutated advanced ovarian carcinoma: Is it cost-effective in Spain?. <i>Gynecologic Oncology</i> , 2022, 164, 406-414. | 0.6 | 4 |
| 3 | Dostarlimab for the treatment of advanced endometrial cancer. <i>Expert Review of Clinical Pharmacology</i> , 2022, 15, 1-9. | 1.3 | 14 |
| 4 | Phase 2 Trial (POLA Study) of Lurbinectedin plus Olaparib in Patients with Advanced Solid Tumors: Results of Efficacy, Tolerability, and the Translational Study. <i>Cancers</i> , 2022, 14, 915. | 1.7 | 4 |
| 5 | Prognosis Stratification Tools in Early-Stage Endometrial Cancer: Could We Improve Their Accuracy?. <i>Cancers</i> , 2022, 14, 912. | 1.7 | 4 |
| 6 | Antitumoral Effect of Plocabulin in High Grade Serous Ovarian Carcinoma Cell Line Models. <i>Frontiers in Oncology</i> , 2022, 12, 862321. | 1.3 | 1 |
| 7 | Bevacizumab in recurrent ovarian cancer: could it be particularly effective in patients with clear cell carcinoma?. <i>Clinical and Translational Oncology</i> , 2021, 23, 536-542. | 1.2 | 5 |
| 8 | Incidence of venous thromboembolic events in cancer patients receiving immunotherapy: a single-institution experience. <i>Clinical and Translational Oncology</i> , 2021, 23, 1245-1252. | 1.2 | 21 |
| 9 | SEOM clinical guideline in ovarian cancer (2020). <i>Clinical and Translational Oncology</i> , 2021, 23, 961-968. | 1.2 | 22 |
| 10 | Proteomic Analysis of Low-Grade, Early-Stage Endometrial Carcinoma Reveals New Dysregulated Pathways Associated with Cell Death and Cell Signaling. <i>Cancers</i> , 2021, 13, 794. | 1.7 | 31 |
| 11 | Inflammatory indexes neutrophils/lymphocytes, platelets/lymphocytes and red-cell distribution width (RDW) as prognostic biomarkers in advanced solitary fibrous tumors (SFT) treated with pazopanib: Correlative study of GEIS-32 trial.. <i>Journal of Clinical Oncology</i> , 2021, 39, 11511-11511. | 0.8 | 0 |
| 12 | A phase Ib/II study of selinexor in combination with imatinib in patients with advanced gastrointestinal stromal tumor (GIST): SeliGIST/GEIS-41 trial.. <i>Journal of Clinical Oncology</i> , 2021, 39, 11534-11534. | 0.8 | 1 |
| 13 | Efficacy and safety of tisotumab vedotin in previously treated recurrent or metastatic cervical cancer (innovaTV 204/GOG-3023/ENGOT-cx6): a multicentre, open-label, single-arm, phase 2 study. <i>Lancet Oncology</i> , The, 2021, 22, 609-619. | 5.1 | 186 |
| 14 | Management of advanced ovarian cancer in Spain: an expert Delphi consensus. <i>Journal of Ovarian Research</i> , 2021, 14, 72. | 1.3 | 2 |
| 15 | Retrospective world-wide registry on the efficacy of immune checkpoint inhibitors in alveolar soft part sarcoma: Updated results from sixty patients.. <i>Journal of Clinical Oncology</i> , 2021, 39, 11564-11564. | 0.8 | 4 |
| 16 | GEIS 39: Phase II trial of nabpaclitaxel for the treatment of patient with multiply relapsed/refractory desmoplastic small round cell tumor (DSRCT) and Ewing sarcoma (EwS).. <i>Journal of Clinical Oncology</i> , 2021, 39, 11529-11529. | 0.8 | 1 |
| 17 | Health-related quality of life (QoL) in platinum-resistant ovarian cancer patients treated with olaparib and pegylated liposomal doxorubicin (PLD), a multicenter single-arm phase II clinical trial (ROLANDO, GEICO-1601).. <i>Journal of Clinical Oncology</i> , 2021, 39, 5549-5549. | 0.8 | 1 |
| 18 | Olaparib in combination with pegylated liposomal doxorubicin for platinum-resistant ovarian cancer regardless of BRCA status: a GEICO phase III trial (ROLANDO study). <i>ESMO Open</i> , 2021, 6, 100212. | 2.0 | 15 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Clinicopathological features and prognostic significance of CTNNB1 mutation in low-grade, early-stage endometrial endometrioid carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2021, 479, 1167-1176. | 1.4 | 26 |
| 20 | Uncommon and peculiar soft tissue sarcomas: Multidisciplinary review and practical recommendations for diagnosis and treatment. Spanish group for Sarcoma research (GEIS â€“ GROUP). Part I. <i>Cancer Treatment Reviews</i> , 2021, 99, 102259. | 3.4 | 6 |
| 21 | Uncommon and peculiar soft tissue sarcomas: Multidisciplinary review and practical recommendations. Spanish Group for Sarcoma research (GEIS â€“GROUP). Part II. <i>Cancer Treatment Reviews</i> , 2021, 99, 102260. | 3.4 | 7 |
| 22 | Long-term response to olaparib in BRCA1-related ovarian cancer with brain metastases. <i>International Journal of Gynecological Cancer</i> , 2021, 31, 1292-1296. | 1.2 | 2 |
| 23 | Eribulin activity in soft tissue sarcoma monolayer and three-dimensional cell line models: could the combination with other drugs improve its antitumoral effect?. <i>Cancer Cell International</i> , 2021, 21, 646. | 1.8 | 6 |
| 24 | The hallmarks of ovarian cancer: Focus on angiogenesis and micro-environment and new models for their characterisation. <i>European Journal of Cancer, Supplement</i> , 2020, 15, 49-55. | 2.2 | 9 |
| 25 | Are antiangiogenics a good â€“partnerâ€™ for immunotherapy in ovarian cancer?. <i>Angiogenesis</i> , 2020, 23, 543-557. | 3.7 | 10 |
| 26 | Nivolumab and sunitinib combination in advanced soft tissue sarcomas: a multicenter, single-arm, phase Ib/II trial. , 2020, 8, e001561. | | 85 |
| 27 | Primary results from CECILIA, a global single-arm phase II study evaluating bevacizumab, carboplatin and paclitaxel for advanced cervical cancer. <i>Gynecologic Oncology</i> , 2020, 159, 142-149. | 0.6 | 30 |
| 28 | Prognostic value of neutrophil-to-lymphocyte ratio and other inflammatory markers in patients with high-risk soft tissue sarcomas. <i>Clinical and Translational Oncology</i> , 2020, 22, 1849-1856. | 1.2 | 12 |
| 29 | Atezolizumab in combination with bevacizumab and chemotherapy versus bevacizumab and chemotherapy in recurrent ovarian cancer â€“ a randomized phase III trial (AGO-OVAR 2.29/ENGOT-ov34). <i>International Journal of Gynecological Cancer</i> , 2020, 30, 1997-2001. | 1.2 | 11 |
| 30 | Assessment of Safety and Efficacy of Combined Trabectedin and Low-Dose Radiotherapy for Patients With Metastatic Soft-Tissue Sarcomas. <i>JAMA Oncology</i> , 2020, 6, 535. | 3.4 | 33 |
| 31 | Niraparib in Patients With Newly Diagnosed Advanced Ovarian Cancer. <i>Obstetrical and Gynecological Survey</i> , 2020, 75, 29-31. | 0.2 | 2 |
| 32 | Pazopanib for treatment of typical solitary fibrous tumours: a multicentre, single-arm, phase 2 trial. <i>Lancet Oncology, The</i> , 2020, 21, 456-466. | 5.1 | 51 |
| 33 | SEOM clinical guidelines for cervical cancer (2019). <i>Clinical and Translational Oncology</i> , 2020, 22, 270-278. | 1.2 | 13 |
| 34 | Patientâ€“reported outcomes from KATHERINE: A phase 3 study of adjuvant trastuzumab emtansine versus trastuzumab in patients with residual invasive disease after neoadjuvant therapy for human epidermal growth factor receptor 2â€“positive breast cancer. <i>Cancer</i> , 2020, 126, 3132-3139. | 2.0 | 14 |
| 35 | Trabectedin and radiotherapy in soft-tissue sarcoma (TRASTS) study: An international, prospective, phase II trial in localized myxoid liposarcomaâ€“A collaborative Spanish (GEIS), Italian (ISG) and French (FSG) group study.. <i>Journal of Clinical Oncology</i> , 2020, 38, 11514-11514. | 0.8 | 7 |
| 36 | IMMUNOSARC: a collaborative Spanish (GEIS) and Italian (ISG) sarcoma groups phase I/II trial of sunitinib and nivolumab in advanced soft tissue and bone sarcoma: Results from the phase II part, bone sarcoma cohort.. <i>Journal of Clinical Oncology</i> , 2020, 38, 11522-11522. | 0.8 | 14 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 37 | Patient-reported outcomes (PRO) in the GARNET trial in patients (pts) with advanced or recurrent dMMR/MSI-H endometrial cancer (EC) treated with dostarlimab.. Journal of Clinical Oncology, 2020, 38, e18032-e18032. | 0.8 | 2 |
| 38 | 3D Culture Modelling: An Emerging Approach for Translational Cancer Research in Sarcomas. Current Medicinal Chemistry, 2020, 27, 4778-4788. | 1.2 | 6 |
| 39 | Patient-reported outcomes (PRO) in patients (pts) with advanced or recurrent dMMR/MSI-H endometrial cancer (EC) treated with dostarlimab in the GARNET trial.. Journal of Clinical Oncology, 2020, 38, 275-275. | 0.8 | 2 |
| 40 | Pazopanib for treatment of advanced extraskeletal myxoid chondrosarcoma: a multicentre, single-arm, phase 2 trial. Lancet Oncology, The, 2019, 20, 1252-1262. | 5.1 | 57 |
| 41 | Primary results from CECILIA, a global single-arm phase II study evaluating bevacizumab (BEV), carboplatin (C) and paclitaxel (P) for advanced cervical cancer (aCC). Annals of Oncology, 2019, 30, v429. | 0.6 | 3 |
| 42 | Niraparib in Patients with Newly Diagnosed Advanced Ovarian Cancer. New England Journal of Medicine, 2019, 381, 2391-2402. | 13.9 | 1,337 |
| 43 | Patient-reported outcomes and final overall survival results from the randomized phase 3 PENELOPE trial evaluating pertuzumab in low tumor human epidermal growth factor receptor 3 (HER3) mRNA-expressing platinum-resistant ovarian cancer. International Journal of Gynecological Cancer, 2019, 29, 1141-1147. | 1.2 | 11 |
| 44 | Trebananib or placebo plus carboplatin and paclitaxel as first-line treatment for advanced ovarian cancer (TRINOVA-3/ENGOT-ov2/GOG-3001): a randomised, double-blind, phase 3 trial. Lancet Oncology, The, 2019, 20, 862-876. | 5.1 | 68 |
| 45 | Myoinvasive Pattern as a Prognostic Marker in Low-Grade, Early-Stage Endometrioid Endometrial Carcinoma. Cancers, 2019, 11, 1845. | 1.7 | 17 |
| 46 | Pazopanib for treatment of advanced malignant and dedifferentiated solitary fibrous tumour: a multicentre, single-arm, phase 2 trial. Lancet Oncology, The, 2019, 20, 134-144. | 5.1 | 97 |
| 47 | Trastuzumab Emtansine for Residual Invasive HER2-Positive Breast Cancer. New England Journal of Medicine, 2019, 380, 617-628. | 13.9 | 1,610 |
| 48 | Patient-reported outcomes (PROs) from KATHERINE: A phase III study of adjuvant trastuzumab emtansine (T-DM1) versus trastuzumab (H) in patients (pts) with residual invasive disease after neoadjuvant therapy for HER2-positive breast cancer.. Journal of Clinical Oncology, 2019, 37, 513-513. | 0.8 | 3 |
| 49 | AGO-OVAR 2.29 (ENGOT-ov34): Atezolizumab in combination with bevacizumab and chemotherapy versus bevacizumab and chemotherapy in recurrent ovarian cancer (ROC).. Journal of Clinical Oncology, 2019, 37, TPS5601-TPS5601. | 0.8 | 3 |
| 50 | Multi-institutional european single-arm phase II trial of pazopanib in advanced typical solitary fibrous tumors (SFT): A collaborative Spanish (GEIS), Italian (ISG), and French (FSG) sarcoma groups study.. Journal of Clinical Oncology, 2019, 37, 11038-11038. | 0.8 | 0 |
| 51 | Phase II Study of Gemcitabine Plus Sirolimus in Previously Treated Patients with Advanced Soft-Tissue Sarcoma: a Spanish Group for Research on Sarcomas (GEIS) Study. Targeted Oncology, 2018, 13, 81-87. | 1.7 | 8 |
| 52 | SEOM clinical guidelines for endometrial cancer (2017). Clinical and Translational Oncology, 2018, 20, 29-37. | 1.2 | 27 |
| 53 | Assessment of socio-economic, physical and mental health status of long-term cancer survivors. Annals of Oncology, 2018, 29, viii624-viii625. | 0.6 | 0 |
| 54 | Long-term response to first-line bevacizumab-based therapy in patients with metastatic breast cancer: results of the observational “LORENA” study. OncoTargets and Therapy, 2018, Volume 11, 5845-5852. | 1.0 | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Predicting Response to Standard First-line Treatment in High-grade Serous Ovarian Carcinoma by Angiogenesis-related Genes. <i>Anticancer Research</i> , 2018, 38, 5393-5400. | 0.5 | 11 |
| 56 | Safety and dose modification for patients receiving niraparib. <i>Annals of Oncology</i> , 2018, 29, 1784-1792. | 0.6 | 125 |
| 57 | The role of glycosyltransferase enzyme GCNT3 in colon and ovarian cancer prognosis and chemoresistance. <i>Scientific Reports</i> , 2018, 8, 8485. | 1.6 | 26 |
| 58 | Nilotinib as Coadjuvant Treatment with Doxorubicin in Patients with Sarcomas: A Phase I Trial of the Spanish Group for Research on Sarcoma. <i>Clinical Cancer Research</i> , 2018, 24, 5239-5249. | 3.2 | 21 |
| 59 | IMMUNOSARC: A collaborative Spanish (GEIS) and Italian (ISG) Sarcoma Groups phase I/II trial of sunitinib plus nivolumab in selected bone and soft tissue sarcoma subtypes—Results of the phase I part.. <i>Journal of Clinical Oncology</i> , 2018, 36, 11515-11515. | 0.8 | 8 |
| 60 | Phase 1b/2 study of olaratumab plus gemcitabine and docetaxel for the treatment of advanced soft tissue sarcoma (STS) (ANNOUNCE 2): Phase 1b results.. <i>Journal of Clinical Oncology</i> , 2018, 36, 11542-11542. | 0.8 | 2 |
| 61 | Preliminary results from CECILIA, an open-label global safety study of bevacizumab (BEV), carboplatin (C) and paclitaxel (P) therapy for metastatic, recurrent or persistent cervical cancer (CC).. <i>Journal of Clinical Oncology</i> , 2018, 36, 5528-5528. | 0.8 | 3 |
| 62 | High-throughput 3-dimensional culture of epithelial ovarian cancer cells as preclinical model of disease. <i>Oncotarget</i> , 2018, 9, 21893-21903. | 0.8 | 21 |
| 63 | Fifth Ovarian Cancer Consensus Conference of the Gynecologic Cancer InterGroup: first-line interventions. <i>Annals of Oncology</i> , 2017, 28, 711-717. | 0.6 | 125 |
| 64 | Malignant bone tumors (other than Ewing's™s): clinical practice guidelines for diagnosis, treatment and follow-up by Spanish Group for Research on Sarcomas (GEIS). <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 80, 1113-1131. | 1.1 | 30 |
| 65 | Gemcitabine plus sirolimus for relapsed and progressing osteosarcoma patients after standard chemotherapy: a multicenter, single-arm phase II trial of Spanish Group for Research on Sarcoma (GEIS). <i>Annals of Oncology</i> , 2017, 28, 2994-2999. | 0.6 | 45 |
| 66 | A definition for aggressive disease in patients with HER-2 negative metastatic breast cancer: an expert consensus of the Spanish Society of Medical Oncology (SEOM). <i>Clinical and Translational Oncology</i> , 2017, 19, 616-624. | 1.2 | 3 |
| 67 | Targeted Sequencing Reveals Low-Frequency Variants in <i>EPHA</i> Genes as Markers of Paclitaxel-Induced Peripheral Neuropathy. <i>Clinical Cancer Research</i> , 2017, 23, 1227-1235. | 3.2 | 16 |
| 68 | Multi-institutional European single-arm phase II trial of pazopanib in advanced malignant/dedifferentiated solitary fibrous tumors (SFT): A collaborative Spanish (GEIS), Italian (ISG), and French (FSG) sarcoma groups study.. <i>Journal of Clinical Oncology</i> , 2017, 35, 11003-11003. | 0.8 | 3 |
| 69 | International single-arm phase II trial of pazopanib in advanced extraskeletal myxoid chondrosarcoma: A Collaborative Spanish (GEIS), Italian (ISG) and French (FSG) Sarcoma Groups study.. <i>Journal of Clinical Oncology</i> , 2017, 35, 11062-11062. | 0.8 | 3 |
| 70 | Open label phase II clinical trial of orteronel (TAK-700) in metastatic or advanced non-resectable granulosa cell ovarian tumors: The Greko II study.. <i>Journal of Clinical Oncology</i> , 2017, 35, 5577-5577. | 0.8 | 1 |
| 71 | Comparison of risk classification between EndoPredict and MammaPrint in ER-positive/HER2-negative primary invasive breast cancer. <i>PLoS ONE</i> , 2017, 12, e0183452. | 1.1 | 11 |
| 72 | CD8+ TILs in early stage epithelial ovarian cancer: A GEICO study.. <i>Journal of Clinical Oncology</i> , 2017, 35, 5543-5543. | 0.8 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 73 | Long-term cancer survivors: Satisfaction with medical care and information received during the diagnosis and treatment of cancer.. Journal of Clinical Oncology, 2017, 35, e21573-e21573. | 0.8 | 0 |
| 74 | Long-term responders to first-line bevacizumab-based therapy in metastatic breast cancer (MBC) patients: Results of the observational study LORENA.. Journal of Clinical Oncology, 2017, 35, e12563-e12563. | 0.8 | 0 |
| 75 | The NER-related gene <i>GTF2H5</i> predicts survival in high-grade serous ovarian cancer patients. Journal of Gynecologic Oncology, 2016, 27, e7. | 1.0 | 30 |
| 76 | SEOM Clinical Guideline in ovarian cancer (2016). Clinical and Translational Oncology, 2016, 18, 1206-1212. | 1.2 | 15 |
| 77 | Randomized Phase II Study of Trabectedin and Doxorubicin Compared With Doxorubicin Alone as First-Line Treatment in Patients With Advanced Soft Tissue Sarcomas: A Spanish Group for Research on Sarcoma Study. Journal of Clinical Oncology, 2016, 34, 2294-2302. | 0.8 | 61 |
| 78 | Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer. New England Journal of Medicine, 2016, 375, 2154-2164. | 13.9 | 1,860 |
| 79 | Final results of a phase 3 study of trebananib plus weekly paclitaxel in recurrent ovarian cancer (TRINOVA-1): Long-term survival, impact of ascites, and progression-free survival-2. Gynecologic Oncology, 2016, 143, 27-34. | 0.6 | 81 |
| 80 | Phosphorylated-insulin growth factor I receptor (p-IGF1R) and metalloproteinase-3 (MMP3) expression in advanced gastrointestinal stromal tumors (GIST). A GEIS 19 study. Clinical Sarcoma Research, 2016, 6, 10. | 2.3 | 1 |
| 81 | <i>CUL4A</i> and <i>ERCC1</i> genes predictive factors for trabectedin efficacy in advanced soft tissue sarcomas (STS): A Spanish Group for Sarcoma Research (GEIS) study.. Journal of Clinical Oncology, 2016, 34, 11048-11048. | 0.8 | 2 |
| 82 | CORAIL trial: Randomized phase III study of lurbinectedin (PM01183) versus pegylated liposomal doxorubicin (PLD) or topotecan (T) in patients with platinum-resistant ovarian cancer.. Journal of Clinical Oncology, 2016, 34, TPS5597-TPS5597. | 0.8 | 6 |
| 83 | Predictive value of angiogenesis-related gene profiling in patients with HER2-negative metastatic breast cancer treated with bevacizumab and weekly paclitaxel. Oncotarget, 2016, 7, 24217-24227. | 0.8 | 8 |
| 84 | Open-label phase II clinical trial of orteronel (TAK-700) in metastatic or advanced nonresectable granulosa cell ovarian tumors: The GREKO II study.. Journal of Clinical Oncology, 2016, 34, TPS2598-TPS2598. | 0.8 | 0 |
| 85 | Prolonged Response to Aflibercept in Ovarian Cancer Relapse: A Case Report. Tumori, 2015, 101, e29-e31. | 0.6 | 3 |
| 86 | SEOM guidelines for cervical cancer. Clinical and Translational Oncology, 2015, 17, 1036-1042. | 1.2 | 27 |
| 87 | Role of Surgery in Patients with Recurrent, Metastatic, or Unresectable Locally Advanced Gastrointestinal Stromal Tumors Sensitive to Imatinib: A Retrospective Analysis of the Spanish Group for Research on Sarcoma (GEIS). Annals of Surgical Oncology, 2015, 22, 2948-2957. | 0.7 | 47 |
| 88 | Phase II trial of gemcitabine plus rapamycin as second line in advanced osteosarcoma: A Spanish Group for Sarcoma Research (GEIS) Study.. Journal of Clinical Oncology, 2015, 33, 10530-10530. | 0.8 | 1 |
| 89 | Long-term functional results and quality of life after open partial laryngectomy and non-surgical larynx preservation.. Journal of Clinical Oncology, 2015, 33, e17056-e17056. | 0.8 | 0 |
| 90 | Open label phase II clinical trial of orteronel (TAK-700) in metastatic or advanced non-resectable granulosa cell ovarian tumors: The Greko II studyâ€™GETHI 2013-01.. Journal of Clinical Oncology, 2015, 33, TPS5612-TPS5612. | 0.8 | 1 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Neutrophil-lymphocyte ratio in stage II-III testicular germ cell tumors before initiating chemotherapy: Correlation with survival.. Journal of Clinical Oncology, 2015, 33, e15564-e15564. | 0.8 | 0 |
| 92 | Early stage ovarian cancer clinical behavior according to FIGO 2014 Staging changes with a focus on IC subtype: data from prospective GEICO registry.. Journal of Clinical Oncology, 2015, 33, 5554-5554. | 0.8 | 1 |
| 93 | Predictive factors of early death after a comprehensive geriatric assessment in elderly cancer patients.. Journal of Clinical Oncology, 2015, 33, e20530-e20530. | 0.8 | 0 |
| 94 | Predictive factors of grade 3-5 toxicity in elderly cancer patients treated with chemotherapy: A prospective multicenter study (TEP study: Toxicity in Elderly Patient).. Journal of Clinical Oncology, 2015, 33, e20535-e20535. | 0.8 | 0 |
| 95 | Continuation of bevacizumab and addition of hormone therapy following weekly paclitaxel therapy in HER2-negative metastatic breast cancer. OncoTargets and Therapy, 2014, 7, 2175. | 1.0 | 11 |
| 96 | NIS Mediates Iodide Uptake in the Female Reproductive Tract and Is a Poor Prognostic Factor in Ovarian Cancer. Journal of Clinical Endocrinology and Metabolism, 2014, 99, E1199-E1208. | 1.8 | 26 |
| 97 | SEOM guideline in ovarian cancer 2014. Clinical and Translational Oncology, 2014, 16, 1067-1071. | 1.2 | 13 |
| 98 | Anti-Angiopoietin Therapy With Trebananib for Recurrent Ovarian Cancer (TRINOVA-1). Obstetrical and Gynecological Survey, 2014, 69, 661-662. | 0.2 | 1 |
| 99 | Anti-angiopoietin therapy with trebananib for recurrent ovarian cancer (TRINOVA-1): a randomised, multicentre, double-blind, placebo-controlled phase 3 trial. Lancet Oncology, The, 2014, 15, 799-808. | 5.1 | 279 |
| 100 | Prognostic/predictive biomarkers in advanced soft tissue sarcomas (STS): Translational research associated to randomized phase II trial comparing trabectedin-doxorubicin versus doxorubicinâ€”A GEIS study.. Journal of Clinical Oncology, 2014, 32, 10500-10500. | 0.8 | 2 |
| 101 | Phase II study of gemcitabine (GEM) plus sirolimus (SIR) in previously treated patients with advanced soft tissue sarcoma (STS): A Spanish Group for Research on Sarcomas (GEIS) study.. Journal of Clinical Oncology, 2014, 32, 10594-10594. | 0.8 | 2 |
| 102 | Predictive value of angiogenesis-related gene profiling in patients with HER2-negative metastatic breast cancer (MBC) treated with bevacizumab and weekly paclitaxel (Bev-Pac).. Journal of Clinical Oncology, 2014, 32, 564-564. | 0.8 | 2 |
| 103 | Nilotinib as co-adjuvant treatment with doxorubicin in sarcomas: Phase I trial resultsâ€”A Spanish Group for Research on Sarcoma (GEIS) study.. Journal of Clinical Oncology, 2014, 32, 10573-10573. | 0.8 | 0 |
| 104 | Head and neck cancer in the elderly: Survival and toxicity with biochemoradiation therapy.. Journal of Clinical Oncology, 2014, 32, e17054-e17054. | 0.8 | 0 |
| 105 | Open-label phase II clinical trial of orteronel (TAK-700) in metastatic or advanced nonresectable granulosa cell ovarian tumors: The GREKO II study.. Journal of Clinical Oncology, 2014, 32, TPS5626-TPS5626. | 0.8 | 0 |
| 106 | Early-stage ovarian cancer: Clinical outcome and analysis of prognostic factorsâ€”Results from a prospective registry of GEICO (Spanish Group for Ovarian Cancer Research).. Journal of Clinical Oncology, 2014, 32, 5582-5582. | 0.8 | 0 |
| 107 | Expression of angiopoietin (Ang) pathway markers and their relationship to progression-free survival (PFS) in TRINOVA-1.. Journal of Clinical Oncology, 2014, 32, 5542-5542. | 0.8 | 0 |
| 108 | Ketoconazole as inhibitor of the enzyme CYP17 in locally advanced or disseminated granulosa cell tumors of the ovary (the GreKo I study) (gethi 11-03).. Journal of Clinical Oncology, 2014, 32, 5558-5558. | 0.8 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | SEOM clinical guidelines for the treatment of osteosarcoma in adults-2013. <i>Clinical and Translational Oncology</i> , 2013, 15, 1037-1043. | 1.2 | 11 |
| 110 | GEICO (Spanish Group for Investigation on Ovarian Cancer) treatment guidelines in ovarian cancer 2012. <i>Clinical and Translational Oncology</i> , 2013, 15, 509-525. | 1.2 | 9 |
| 111 | Efficacy and safety of front-line bevacizumab (BEV), weekly paclitaxel (wPAC), and q3w carboplatin (C) in elderly patients (pts) with ovarian cancer (OC): Subgroup analysis of OCTAVIA.. <i>Journal of Clinical Oncology</i> , 2013, 31, 5544-5544. | 0.8 | 2 |
| 112 | Long-term responders to first-line bevacizumab-based therapy among patients (pts) with HER2-negative metastatic breast cancer (MBC): Retrospective results of an ambispective observational study.. <i>Journal of Clinical Oncology</i> , 2013, 31, e12558-e12558. | 0.8 | 0 |
| 113 | mTOR pathway inhibition in renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2012, 30, 356-361. | 0.8 | 8 |
| 114 | Targeting the endothelin axis in prostate carcinoma. <i>Tumor Biology</i> , 2012, 33, 421-426. | 0.8 | 29 |
| 115 | Phase II study of neoadjuvant high-dose ifosfamide with concurrent radiotherapy followed by surgical resection in high-risk soft tissue sarcoma: A Spanish Group for Research on Sarcomas (GEIS) study.. <i>Journal of Clinical Oncology</i> , 2012, 30, 10052-10052. | 0.8 | 3 |
| 116 | Gene profiling in breast cancer: Time to move forward. <i>Cancer Treatment Reviews</i> , 2011, 37, 416-21. | 3.4 | 8 |
| 117 | Feasibility of a Modified Outpatient Regimen of Intravenous/Intraperitoneal Chemotherapy in Optimally Debulked Stage III Ovarian Cancer Patients. <i>International Journal of Gynecological Cancer</i> , 2011, 21, 1048-1055. | 1.2 | 11 |
| 118 | From targeted therapy in ovarian cancer to personalizing therapy for ovarian cancer. <i>Expert Opinion on Investigational Drugs</i> , 2011, 20, 591-594. | 1.9 | 3 |
| 119 | Angiogenesis as a therapeutic target in urothelial carcinoma. <i>Anti-Cancer Drugs</i> , 2010, 21, 890-896. | 0.7 | 20 |
| 120 | An 8-gene qRT-PCR-based gene expression score that has prognostic value in early breast cancer. <i>BMC Cancer</i> , 2010, 10, 336. | 1.1 | 22 |
| 121 | Molecular characterization of ovarian cancer by gene-expression profiling. <i>Gynecologic Oncology</i> , 2010, 118, 88-92. | 0.6 | 41 |
| 122 | The miR-200 family controls β -tubulin III expression and is associated with paclitaxel-based treatment response and progression-free survival in ovarian cancer patients. <i>Endocrine-Related Cancer</i> , 2010, 18, 85-95. | 1.6 | 188 |
| 123 | Angiogenesis and ovarian cancer. <i>Clinical and Translational Oncology</i> , 2009, 11, 564-571. | 1.2 | 59 |
| 124 | Intra-abdominal desmoplastic small round cell tumour in a 39-year-old man. <i>Clinical and Translational Oncology</i> , 2009, 11, 770-772. | 1.2 | 0 |
| 125 | Aurora kinases as prognostic biomarkers in ovarian carcinoma. <i>Human Pathology</i> , 2009, 40, 631-638. | 1.1 | 35 |
| 126 | Comparison of Prognostic Gene Profiles Using qRT-PCR in Paraffin Samples: A Retrospective Study in Patients with Early Breast Cancer. <i>PLoS ONE</i> , 2009, 4, e5911. | 1.1 | 18 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Cirrhosis-like radiological pattern in patients with breast cancer. <i>Clinical and Translational Oncology</i> , 2008, 10, 111-116. | 1.2 | 3 |
| 128 | Meningeal relapse by follicular lymphoma as a single mass. <i>Clinical and Translational Oncology</i> , 2008, 10, 180-181. | 1.2 | 0 |
| 129 | Angiogenesis-Related Gene Expression Profile with Independent Prognostic Value in Advanced Ovarian Carcinoma. <i>PLoS ONE</i> , 2008, 3, e4051. | 1.1 | 20 |
| 130 | Phospholipid Hydroperoxide Glutathione Peroxidase (PHGPx) expression is downregulated in poorly differentiated breast invasive ductal carcinoma. <i>Free Radical Research</i> , 2007, 41, 681-687. | 1.5 | 25 |
| 131 | High-throughput techniques in breast cancer: A clinical perspective. <i>European Journal of Cancer</i> , 2006, 42, 598-607. | 1.3 | 13 |
| 132 | Clinical Role of Sm-153 EDTMP in the Treatment of Painful Bone Metastatic Disease. <i>Clinical Nuclear Medicine</i> , 2006, 31, 605-610. | 0.7 | 17 |
| 133 | Localisation of COX-2 protein is different in breast ductal carcinoma and adjacent non-tumour ductal epithelium. <i>Clinical and Translational Oncology</i> , 2005, 7, 239-243. | 1.2 | 3 |
| 134 | Breast Cancer Prognosis Determined by Gene Expression Profiling: A Quantitative Reverse Transcriptase Polymerase Chain Reaction Study. <i>Journal of Clinical Oncology</i> , 2005, 23, 7278-7285. | 0.8 | 54 |
| 135 | Breast carcinomas fulfill the Warburg hypothesis and provide metabolic markers of cancer prognosis. <i>Carcinogenesis</i> , 2005, 26, 2095-2104. | 1.3 | 155 |
| 136 | Implications of Oxidative Stress and Cell Membrane Lipid Peroxidation in Human Cancer (Spain). <i>Cancer Causes and Control</i> , 2004, 15, 707-719. | 0.8 | 138 |
| 137 | Phase II study of neoadjuvant treatment of rectal cancer with oxaliplatin, raltitrexed and radiotherapy. <i>Journal of Clinical Oncology</i> , 2004, 22, 3746-3746. | 0.8 | 1 |
| 138 | Phase II study of neoadjuvant treatment of rectal cancer with oxaliplatin, raltitrexed and radiotherapy. <i>Journal of Clinical Oncology</i> , 2004, 22, 3746-3746. | 0.8 | 3 |