

Kazutaka Saito

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5870344/kazutaka-saito-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

193
papers

3,335
citations

32
h-index

50
g-index

207
ext. papers

3,803
ext. citations

3.2
avg. IF

4.82
L-index

#	Paper	IF	Citations
193	Differences in CRP and De Ritis ratio predictive abilities of cancer specific survival between ethnic groups.. <i>Journal of Clinical Oncology</i> , 2022 , 40, 394-394	2.2	
192	Impact of age on functional decline following radical nephrectomy: Analysis of the International Marker Consortium for Renal Cancer (INMARC).. <i>Journal of Clinical Oncology</i> , 2022 , 40, 395-395	2.2	
191	Does presence of diabetes mellitus impact patient oncological outcomes in renal cell carcinoma: A multicenter analysis.. <i>Journal of Clinical Oncology</i> , 2022 , 40, 325-325	2.2	
190	Nonmetastatic castration-resistant prostate cancer treated with salvage focal brachytherapy after external beam radiotherapy. <i>IJU Case Reports</i> , 2021 , 4, 228-230	0.5	0
189	Nonuse of antimicrobial prophylaxis in clean surgeries for adrenal and renal tumors: Results of the risk-based strategy in 1362 consecutive patients. <i>International Journal of Urology</i> , 2021 , 28, 1032-1038	2.3	
188	Significance of Bladder Neck Involvement in Risk Substratification of Intermediate-Risk Non-muscle-invasive Bladder Cancer. <i>European Urology Focus</i> , 2021 , 7, 366-372	5.1	0
187	Outcomes of gasless laparoendoscopic single-port partial nephrectomy in 356 consecutive patients: Feasibility of a clampless and sutureless technique. <i>International Journal of Urology</i> , 2021 , 28, 302-307	2.3	3
186	Early C-reactive protein kinetics predict survival of patients with advanced urothelial cancer treated with pembrolizumab. <i>Cancer Immunology, Immunotherapy</i> , 2021 , 70, 657-665	7.4	7
185	Elevated preoperative C-reactive protein is associated with renal functional decline and non-cancer mortality in surgically treated renal cell carcinoma: analysis from the International Marker Consortium for Renal Cancer (INMARC). <i>BJU International</i> , 2021 , 127, 311-317	5.6	3
184	Potential of Perfusion Magnetic Resonance Imaging to Predict Residual Renal Function after Radical Nephroureterectomy. <i>Urologia Internationalis</i> , 2021 , 105, 507-513	1.9	0
183	Impact of C-reactive protein flare-response on oncological outcomes in patients with metastatic renal cell carcinoma treated with nivolumab 2021 , 9,		8
182	Medium-term oncological and functional outcomes of hemi-gland brachytherapy using iodine-125 seeds for intermediate-risk unilateral prostate cancer. <i>Brachytherapy</i> , 2021 , 20, 842-848	2.4	1
181	Preoperative Elevation of C-Reactive Protein Is a Predictor for Adverse Oncologic Survival Outcomes for Renal Cell Carcinoma: Analysis from the International Marker Consortium Renal Cancer (INMARC). <i>Clinical Genitourinary Cancer</i> , 2021 , 19, e206-e215	3.3	2
180	Longitudinal changes in patient-reported outcomes after artificial urinary sphincter implantation. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2020 , 12, 240-244	1.9	1
179	Contact with renal sinus is associated with poor prognosis in surgically treated pT1 clear cell renal cell carcinoma. <i>International Journal of Urology</i> , 2020 , 27, 657-662	2.3	1
178	Postoperative renal impairment and longitudinal change in renal function after adrenalectomy in patients with Cushing's syndrome. <i>International Journal of Urology</i> , 2020 , 27, 395-400	2.3	3
177	Application of virtual reality in patient explanation of magnetic resonance imaging-ultrasound fusion prostate biopsy. <i>International Journal of Urology</i> , 2020 , 27, 471-472	2.3	6

176	Incidence and Risk Factors of Hypertension Following Partial Nephrectomy in Patients With Renal Tumors: A Cross-sectional Study of Postoperative Home Blood Pressure and Antihypertensive Medications. <i>Clinical Genitourinary Cancer</i> , 2020 , 18, e619-e628	3.3	2
175	Treatment-induced changes in levels of brain natriuretic peptide in prostate cancer patients receiving gonadotropin-releasing hormone antagonists or agonists. <i>International Journal of Urology</i> , 2020 , 27, 354-355	2.3	3
174	Feasibility and outcomes of selective tetramodal bladder-preservation therapy in elderly patients with muscle-invasive bladder cancer. <i>International Journal of Urology</i> , 2020 , 27, 236-243	2.3	2
173	Renal epithelioid angiomyolipoma: Incidence in a Japanese cohort and diagnostic utility of diffusion-weighted magnetic resonance imaging. <i>International Journal of Urology</i> , 2020 , 27, 599-604	2.3	0
172	Value of extra-target prostate biopsy for the detection of magnetic resonance imaging-missed adverse pathology according to the Prostate Imaging Reporting and Data System scores: Spatial analysis using magnetic resonance-ultrasound fusion images. <i>International Journal of Urology</i> , 2020 , 27, 760-766	2.3	1
171	Association of De Ritis Ratio and Neutrophil Lymphocyte Ratio with renal functional decline and all-cause mortality in renal cell carcinoma.. <i>Journal of Clinical Oncology</i> , 2020 , 38, 741-741	2.2	
170	Impact of diabetes mellitus on functional and survival outcomes in renal cell carcinoma: An international multicenter study.. <i>Journal of Clinical Oncology</i> , 2020 , 38, 666-666	2.2	
169	Impact of oncological versus nononcological on survival outcomes in African Americans with renal cell carcinoma.. <i>Journal of Clinical Oncology</i> , 2020 , 38, 749-749	2.2	
168	Impact of Bladder Neck Involvement on Recurrence in Patients With Non-muscle-invasive Bladder Cancer: An Analysis Based on a Time-dependent Model. <i>Clinical Genitourinary Cancer</i> , 2020 , 18, e62-e70	3.3	3
167	Female Gender Predicts Favorable Prognosis in Patients With Non-metastatic Clear Cell Renal Cell Carcinoma Undergoing Curative Surgery: Results From the International Marker Consortium for Renal Cancer (INMARC). <i>Clinical Genitourinary Cancer</i> , 2020 , 18, 111-116.e1	3.3	4
166	METastasis Reporting and Data System for Prostate Cancer as a Prognostic Imaging Marker in Castration-resistant Prostate Cancer. <i>Clinical Genitourinary Cancer</i> , 2020 , 18, e391-e396	3.3	8
165	Clinical Outcomes of Patients With Histologic Variants of Urothelial Carcinoma Treated With Selective Tetramodal Bladder-preservation Therapy Incorporating Consolidative Partial Cystectomy. <i>Clinical Genitourinary Cancer</i> , 2020 , 18, 268-273.e2	3.3	2
164	The Diagnostic Accuracy of Testicular Torsion by Doctors on Duty Using Sonographic Evaluation with Color Doppler. <i>American Journal of Men's Health</i> , 2020 , 14, 1557988320953003	2.2	5
163	SRY-Positive 46, XX Testicular Disorder of Sexual Development With Leydig Cell Tumor. <i>American Journal of Men's Health</i> , 2020 , 14, 1557988320970071	2.2	2
162	Impact of radiotherapy to the primary tumor on the efficacy of pembrolizumab for patients with advanced urothelial cancer: A preliminary study. <i>Cancer Medicine</i> , 2020 , 9, 8355-8363	4.8	5
161	Impact of fluorodeoxyglucose uptake on positron emission tomography/computed tomography on chemosensitivity and survival in patients with metastatic urothelial carcinoma. <i>International Journal of Urology</i> , 2019 , 26, 820-826	2.3	
160	Usefulness of the inchworm sign on DWI for predicting pT1 bladder cancer progression. <i>European Radiology</i> , 2019 , 29, 3881-3888	8	10
159	Selective tetramodal bladder-preservation therapy, incorporating induction chemoradiotherapy and consolidative partial cystectomy with pelvic lymph node dissection for muscle-invasive bladder cancer: oncological and functional outcomes of 107 patients. <i>BJU International</i> , 2019 , 124, 242-250	5.6	23

158	Application of eye tracking as a user interface for urological surgery. <i>International Journal of Urology</i> , 2019 , 26, 680-681	2.3	1
157	Mixed reality computed tomography-based surgical planning for partial nephrectomy using a head-mounted holographic computer. <i>International Journal of Urology</i> , 2019 , 26, 681-682	2.3	16
156	Progressive Site-Directed Therapy for Castration-Resistant Prostate Cancer: Localization of the Progressive Site as a Prognostic Factor. <i>International Journal of Radiation Oncology Biology Physics</i> , 2019 , 105, 376-381	4	28
155	Renal function after bladder-preserving therapy for patients with muscle-invasive bladder cancer: Results of selective bladder-preserving tetramodality therapy consisting of maximal transurethral resection, induction chemoradiotherapy and partial cystectomy. <i>International Journal of Urology</i> , 2019 , 26, 1010-1012	2.3	1
154	Association of elevated C-reactive protein with oncologic outcomes in renal CELL carcinoma: A multicenter analysis.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 596-596	2.2	1
153	miRNA299 involvement in CYP11B2 expression in aldosterone-producing adenoma. <i>European Journal of Endocrinology</i> , 2019 , 181, 69-78	6.5	7
152	Disparities in survival outcomes in African Americans in renal cell carcinoma: Impact of oncological versus nononcological factors.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 595-595	2.2	
151	Clinical outcomes of muscle-invasive bladder cancer patients with hydronephrosis treated with tetra-modality bladder sparing therapy incorporating consolidative partial cystectomy.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 430-430	2.2	
150	Development and validation of a novel scoring index (C-reactive protein, age, race, and tumor size) to predict renal functional decline post partial nephrectomy.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 597-597	2.3	3
149	Incidence and predictive factors of hypoglycemia after pheochromocytoma resection. <i>International Journal of Urology</i> , 2019 , 26, 273-277	2.3	10
148	Predictive ability of renal cortex enhancement in dynamic computed tomography for residual renal function after nephroureterectomy: Comparison with Tc-diethylenetriaminopentacetic acid renography and validation study. <i>International Journal of Urology</i> , 2019 , 26, 266-272	2.3	2
147	Acute kidney injury and intermediate-term renal function after clampless partial nephrectomy. <i>International Journal of Urology</i> , 2019 , 26, 113-118	2.3	6
146	Intensity ratio curve analysis of small renal masses on T2-weighted magnetic resonance imaging: Differentiation of fat-poor angiomyolipoma from renal cell carcinoma. <i>International Journal of Urology</i> , 2018 , 25, 554-560	2.3	3
145	Higher Serum C-reactive Protein Level Represents the Immunosuppressive Tumor Microenvironment in Patients With Clear Cell Renal Cell Carcinoma. <i>Clinical Genitourinary Cancer</i> , 2018 , 16, e1151-e1158	3.3	20
144	Prediction of Intraoperative Urinary Collecting System Entry in Patients with Peripheral Renal Tumors Undergoing Partial Nephrectomy: Usefulness of Tumor-Centered Multiplanar Reconstruction. <i>Urologia Internationalis</i> , 2018 , 100, 85-91	1.9	0
143	Adherent Perinephric Fat in Asian Patients: Predictors and Impact on Perioperative Outcomes of Partial Nephrectomy. <i>Urologia Internationalis</i> , 2018 , 101, 437-442	1.9	19
142	Impact of Immunohistochemistry-Based Subtypes in Muscle-Invasive Bladder Cancer on Response to Chemoradiation Therapy. <i>International Journal of Radiation Oncology Biology Physics</i> , 2018 , 102, 1408-1416	4	12
141	Potential for computer-aided diagnosis using a convolutional neural network algorithm to diagnose fat-poor angiomyolipoma in enhanced computed tomography and T2-weighted magnetic resonance imaging. <i>International Journal of Urology</i> , 2018 , 25, 978-979	2.3	4

140	Computer-aided diagnosis of prostate cancer on magnetic resonance imaging using a convolutional neural network algorithm. <i>BJU International</i> , 2018 , 122, 411-417	5.6	54
139	Multitask Imaging Monitor for Surgical Navigation: Combination of Touchless Interface and Head-Mounted Display. <i>Urologia Internationalis</i> , 2017 , 98, 486-488	1.9	4
138	Utility of computed diffusion-weighted MRI for predicting aggressiveness of prostate cancer. <i>Journal of Magnetic Resonance Imaging</i> , 2017 , 46, 490-496	5.6	15
137	Potential Utility of Diffusion-Weighted Magnetic Resonance Imaging in Diagnosis of Residual Bladder Cancer before Second Transurethral Resection. <i>Urologia Internationalis</i> , 2017 , 98, 298-303	1.9	3
136	Standardization of the apparent diffusion coefficient value of bladder cancer across different centers: Applicability in predicting aggressive pathologic phenotypes. <i>Clinical Imaging</i> , 2017 , 44, 121-126 ^{2.7}	2.7	2
135	Real-time three-dimensional image angle rectification to improve hand-eye coordination in single-port laparoendoscopic surgery. <i>International Journal of Urology</i> , 2017 , 24, 639-640	2.3	2
134	Impact of the Prostate Imaging Reporting and Data System, Version 2, on MRI Diagnosis for Extracapsular Extension of Prostate Cancer. <i>American Journal of Roentgenology</i> , 2017 , 209, W76-W84	5.4	25
133	Stepwise algorithm using computed tomography and magnetic resonance imaging for diagnosis of fat-poor angiomyolipoma in small renal masses: Development and external validation. <i>International Journal of Urology</i> , 2017 , 24, 511-517	2.3	13
132	Early response of C-reactive protein as a predictor of survival in patients with metastatic renal cell carcinoma treated with tyrosine kinase inhibitors. <i>International Journal of Clinical Oncology</i> , 2017 , 22, 1081-1086	4.2	20
131	Absence of renal artery pseudoaneurysm on computed tomography after minimally-invasive partial nephrectomy using clampless and sutureless techniques. <i>International Journal of Urology</i> , 2017 , 24, 472-473 ^{2.3}	2.3	8
130	A novel interactive educational system in the operating room--the IE system. <i>BMC Medical Education</i> , 2016 , 16, 44	3.3	8
129	Words of Wisdom. Re: Evaluation of Functional Outcomes After Laparoscopic Partial Nephrectomy Using Renal Scintigraphy: Clamped vs. Clampless Technique. <i>European Urology</i> , 2016 , 69, 753	10.2	
128	Ureteral Involvement Is Associated with Poor Prognosis in Upper Urinary Tract Urothelial Carcinoma Patients Treated by Nephroureterectomy: A Multicenter Database Study. <i>European Urology Focus</i> , 2016 , 2, 296-302	5.1	10
127	Words of Wisdom. Re: Nephron-sparing Techniques Independently Decrease the Risk of Cardiovascular Events Relative to Radical Nephrectomy in Patients with a T1a-T1b Renal Mass and Normal Preoperative Renal Function. <i>European Urology</i> , 2016 , 69, 538	10.2	1
126	Reply to Umberto Capitano, Paolo Dell'Oglio, Alessandro Larcher, and Francesco Montorsi's Reply to Kazutaka Saito's Words of Wisdom re: Nephron-sparing Techniques Independently Decrease the Risk of Cardiovascular Events Relative to Radical Nephrectomy in Patients with a T1a-T1b Renal Mass and Normal Preoperative Renal Function. <i>Eur Urol</i> . 2016;69:538. <i>European Urology</i> , 2016 , 70, e99	10.2	
125	Candidate selection for quadrant-based focal ablation through a combination of diffusion-weighted magnetic resonance imaging and prostate biopsy. <i>BJU International</i> , 2016 , 117, 94-101 ^{5.6}	5.6	10
124	Recovery of renal function after radical nephrectomy and risk factors for postoperative severe renal impairment: A Japanese multicenter longitudinal study. <i>International Journal of Urology</i> , 2016 , 23, 219-23	2.3	10
123	High Ki-67 Expression Predicts Favorable Survival in Muscle-Invasive Bladder Cancer Patients Treated With Chemoradiation-Based Bladder-Sparing Protocol. <i>Clinical Genitourinary Cancer</i> , 2015 , 13, e243-e251	3.3	17

122	A novel approach to patient self-monitoring of sonographic examinations using a head-mounted display. <i>Journal of Ultrasound in Medicine</i> , 2015 , 34, 29-35	2.9	7
121	Risk stratification for bladder recurrence of upper urinary tract urothelial carcinoma after radical nephroureterectomy. <i>BJU International</i> , 2015 , 115, 705-12	5.6	33
120	Gasless laparoendoscopic single-port clampless sutureless partial nephrectomy for peripheral renal tumors: perioperative outcomes. <i>International Journal of Urology</i> , 2015 , 22, 349-55	2.3	22
119	Head-mounted display for a personal integrated image monitoring system: ureteral stent placement. <i>Urologia Internationalis</i> , 2015 , 94, 117-20	1.9	19
118	Novel image monitoring system using a head-mounted display for assistants in da Vinci surgery. <i>International Journal of Urology</i> , 2015 , 22, 520-1	2.3	6
117	Novel three-dimensional image system for transurethral surgery. <i>International Journal of Urology</i> , 2015 , 22, 714-5	2.3	3
116	Progression of hypertension after partial nephrectomy in patients with renal tumors: A preliminary report. <i>International Journal of Urology</i> , 2015 , 22, 797-8	2.3	7
115	Pharmacokinetic-pharmacodynamic analysis of sunitinib-induced thrombocytopenia in Japanese patients with renal cell carcinoma. <i>Biological and Pharmaceutical Bulletin</i> , 2015 , 38, 402-10	2.3	11
114	Patient's Self-monitoring of Transurethral Surgical Images Using a Head-mounted Display. <i>Urology Case Reports</i> , 2015 , 3, 27-9	0.5	4
113	A Novel Approach to Surgical Instructions for Scrub Nurses by Using See-Through-Type Head-Mounted Display. <i>CIN - Computers Informatics Nursing</i> , 2015 , 33, 335-8	1.4	5
112	Diagnostic performance and safety of a three-dimensional 14-core systematic biopsy method. <i>BJU International</i> , 2015 , 115, 412-8	5.6	3
111	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Partial Nephrectomy 2015 , 43-64		2
110	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Adrenalectomy 2015 , 85-104		1
109	Application of the RoboSurgeon System in Medical Care 2015 , 175-184		
108	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Radical Nephrectomy 2015 , 23-42		
107	Gasless Single-Port RoboSurgeon Partial Cystectomy: A Hybrid Technique Combining an Intravesical and Extravesical Approach 2015 , 143-158		
106	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Radical Prostatectomy 2015 , 105-126		
105	Gasless Two-Port Access RoboSurgeon Retroperitoneoscopic Radical Nephroureterectomy 2015 , 65-84		1

104	Gasless Single-Port RoboSurgeon Retroperitoneoscopic Pelvic Lymph Node Dissection and Inguinal Hernia Prevention 2015 , 159-173		
103	Gasless Single-Port RoboSurgeon Total Cystectomy with Urinary Diversion 2015 , 127-141		
102	Combination of diffusion-weighted magnetic resonance imaging and extended prostate biopsy predicts lobes without significant cancer: application in patient selection for hemiablativ focal therapy. <i>European Urology</i> , 2014 , 65, 186-92	10.2	24
101	Significance of positive urine cytology on progression and cancer-specific mortality of non-muscle-invasive bladder cancer. <i>Clinical Genitourinary Cancer</i> , 2014 , 12, e87-93	3.3	9
100	Apparent diffusion coefficient value as a biomarker reflecting morphological and biological features of prostate cancer. <i>International Urology and Nephrology</i> , 2014 , 46, 555-61	2.3	34
99	Equivalent survival and improved preservation of renal function after distal ureterectomy compared with nephroureterectomy in patients with urothelial carcinoma of the distal ureter: a propensity score-matched multicenter study. <i>International Journal of Urology</i> , 2014 , 21, 1098-104	2.3	18
98	PD13-06 FAVORABLE OUTCOMES OF GASLESS SINGLE-PORT CLAMPLESS PARTIAL NEPHRECTOMY USING NEW THREE DIMENSIONAL HEAD-MOUNTED DISPLAY SYSTEM (ROBOSURGEON SYSTEM). <i>Journal of Urology</i> , 2014 , 191,	2.5	2
97	Impact of bladder neck involvement on progression in patients with primary non-muscle invasive bladder cancer: a prospective validation study. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 38.e29-36	2.8	14
96	Apparent diffusion coefficient value reflects invasive and proliferative potential of bladder cancer. <i>Journal of Magnetic Resonance Imaging</i> , 2014 , 39, 172-8	5.6	71
95	Phase II trial of first-line chemotherapy with gemcitabine, etoposide, and cisplatin for patients with advanced urothelial carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 35.e1-7	2.8	2
94	A three-dimensional head-mounted display system (RoboSurgeon system) for gasless laparoendoscopic single-port partial cystectomy. <i>Wideochirurgia I Inne Techniki Maloinwazyjne</i> , 2014 , 9, 638-43	1.4	6
93	Integrated image monitoring system using head-mounted display for gasless single-port clampless partial nephrectomy. <i>Wideochirurgia I Inne Techniki Maloinwazyjne</i> , 2014 , 9, 634-7	1.4	6
92	New head-mounted display system applied to endoscopic management of upper urinary tract carcinomas. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2014 , 40, 842-5	2	11
91	Pre-operative risk stratification for cancer-specific survival in patients with renal cell carcinoma with venous involvement who underwent nephrectomy. <i>Japanese Journal of Clinical Oncology</i> , 2014 , 44, 756-61	2.8	8
90	Renal function after radical nephrectomy: development and validation of predictive models in Japanese patients. <i>International Journal of Urology</i> , 2014 , 21, 238-42	2.3	15
89	Clinical application of a modern high-definition head-mounted display in sonography. <i>Journal of Ultrasound in Medicine</i> , 2014 , 33, 1499-504	2.9	17
88	Bone abnormal signal incidentally found in pre-biopsy diffusion-weighted MRI for suspected prostate cancer: what does it reflect?. <i>Urologia Internationalis</i> , 2014 , 93, 170-5	1.9	4
87	MP8-09 ANTIMICROBIAL PROPHYLAXIS IS AVOIDABLE IN MINIMALLY INVASIVE NEPHROURETERECTOMY WITH NEGATIVE URINE CULTURE. <i>Journal of Urology</i> , 2014 , 191,	2.5	1

86	Combined resection of a tumor and the inferior vena cava: report of two cases. <i>Surgery Today</i> , 2014 , 44, 166-70	3	0
85	Longitudinal change in health-related quality of life after intensity-modulated radiation monotherapy for clinically localized prostate cancer. <i>Quality of Life Research</i> , 2014 , 23, 1641-50	3.7	9
84	External validation of TNM-C score in three community hospital cohorts for clear cell renal cell carcinoma. <i>Anticancer Research</i> , 2014 , 34, 921-6	2.3	2
83	Apparent diffusion coefficient as a prognostic biomarker of upper urinary tract cancer: a preliminary report. <i>European Radiology</i> , 2013 , 23, 2206-14	8	29
82	Usefulness of pre-biopsy multiparametric magnetic resonance imaging and clinical variables to reduce initial prostate biopsy in men with suspected clinically localized prostate cancer. <i>Journal of Urology</i> , 2013 , 190, 502-8	2.5	58
81	Diagnostic performance of initial transperineal 14-core prostate biopsy to detect significant cancer. <i>International Urology and Nephrology</i> , 2013 , 45, 645-52	2.3	6
80	High diagnostic ability of multiparametric magnetic resonance imaging to detect anterior prostate cancer missed by transrectal 12-core biopsy. <i>Journal of Urology</i> , 2013 , 190, 867-73	2.5	64
79	Prognostic factors of patients with metastatic renal cell carcinoma with removed metastases: a multicenter study of 556 patients. <i>Urology</i> , 2013 , 82, 846-51	1.6	53
78	Successful reduction of hospital-acquired methicillin-resistant <i>Staphylococcus aureus</i> in a urology ward: a 10-year study. <i>BMC Urology</i> , 2013 , 13, 35	2.2	11
77	Role of C-reactive protein in urological cancers: a useful biomarker for predicting outcomes. <i>International Journal of Urology</i> , 2013 , 20, 161-71	2.3	44
76	Sensitivity to chemoradiation predicts development of metastasis in muscle-invasive bladder cancer patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1270-5	2.8	9
75	49 THE ABNORMALITY OF GLUCOSE TOLERANCE TEST IN PHEOCHROMOCYTOMA: ONE OF THE SIGNIFICANT FINDINGS OF POSTOPERATIVE HYPOGLYCEMIA. <i>Journal of Urology</i> , 2013 , 189,	2.5	1
74	840 NEW THREE-DIMENSIONAL HEAD-MOUNTED DISPLAY SYSTEM (ROBOSURGEON SYSTEM): APPLICATION TO THE INITIAL 80 CASES OF GASLESS SINGLE-PORT ACCESS UROLOGIC SURGERIES. <i>Journal of Urology</i> , 2013 , 189,	2.5	11
73	Prognostic impact of pretreatment C-reactive protein for patients with metastatic renal cell carcinoma treated with tyrosine kinase inhibitors. <i>International Journal of Clinical Oncology</i> , 2013 , 18, 884-9	4.2	37
72	Author's reply: CRP kinetics could be prognostic predictors in urothelial cancer. <i>Nature Reviews Urology</i> , 2013 , 10, 182	5.5	
71	Impact of advanced age on biochemical recurrence after radical prostatectomy in Japanese men according to pathological stage. <i>Japanese Journal of Clinical Oncology</i> , 2013 , 43, 410-6	2.8	11
70	Diabetes mellitus with obesity is a predictor of recurrence in patients with non-metastatic renal cell carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2013 , 43, 740-6	2.8	15
69	Impact of lower urinary tract symptoms on prostate cancer risk among Japanese men with prostate-specific antigen. <i>International Journal of Urology</i> , 2013 , 20, 1163-8	2.3	4

68	New three-dimensional head-mounted display system, TMDU-S-3D system, for minimally invasive surgery application: procedures for gasless single-port radical nephrectomy. <i>International Journal of Urology</i> , 2012 , 19, 886-9	2.3	53
67	Performance of prostate-specific antigen mass in estimation of prostate volume in Japanese men with benign prostate hyperplasia. <i>International Journal of Urology</i> , 2012 , 19, 929-35	2.3	2
66	Selective bladder-sparing protocol consisting of induction low-dose chemoradiotherapy plus partial cystectomy with pelvic lymph node dissection against muscle-invasive bladder cancer: oncological outcomes of the initial 46 patients. <i>BJU International</i> , 2012 , 109, 860-6	5.6	42
65	Characteristics and clinical significance of prostate cancers missed by initial transrectal 12-core biopsy. <i>BJU International</i> , 2012 , 109, 665-71	5.6	31
64	Pathology-based risk stratification of muscle-invasive bladder cancer patients undergoing cystectomy for persistent disease after induction chemoradiotherapy in bladder-sparing approaches. <i>BJU International</i> , 2012 , 110, E203-8	5.6	9
63	Prostate-specific antigen response to deferred combined androgen blockade therapy using bicalutamide predicts survival after subsequent oestrogen and docetaxel therapies in patients with castration-resistant prostate cancer. <i>BJU International</i> , 2012 , 110, 1149-55	5.6	7
62	Impact of C-reactive protein kinetics on survival of patients with advanced urothelial carcinoma treated by second-line chemotherapy with gemcitabine, etoposide and cisplatin. <i>BJU International</i> , 2012 , 110, 1478-84	5.6	25
61	Extended biopsy based criteria incorporating cumulative cancer length for predicting clinically insignificant prostate cancer. <i>BJU International</i> , 2012 , 110, E564-9	5.6	8
60	Editorial comment from Dr Saito to impact of hyponatremia on survival of patients with metastatic renal cell carcinoma treated with molecular targeted therapy. <i>International Journal of Urology</i> , 2012 , 19, 1059	2.3	
59	Role of diffusion-weighted magnetic resonance imaging in predicting sensitivity to chemoradiotherapy in muscle-invasive bladder cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012 , 83, e21-7	4	82
58	1105 GASLESS SINGLE-PORT CLAMPLESS PARTIAL NEPHRECTOMY FOR PERIPHERAL RENAL TUMOR SURGICAL, SHORT-TERM ONCOLOGICAL, AND FUNCTIONAL OUTCOMES. <i>Journal of Urology</i> , 2012 , 187,	2.5	1
57	Effect of diabetes mellitus on high-grade prostate cancer detection among Japanese obese patients with prostate-specific antigen less than 10 ng/mL. <i>Urology</i> , 2012 , 79, 1329-34	1.6	12
56	Antimicrobial prophylaxis is not necessary in clean category minimally invasive surgery for renal and adrenal tumors: a prospective study of 373 consecutive patients. <i>Urology</i> , 2012 , 80, 570-5	1.6	19
55	Gasless Single Port Surgery for Renal Cell Carcinoma: Minimum Incision Endoscopic Surgery 2012 ,		1
54	Diagnostic contribution of C-reactive protein kinetics for gastric metastasis from renal cell carcinoma. <i>International Cancer Conference Journal</i> , 2012 , 1, 93-95	0.9	2
53	Development of a nomogram incorporating serum C-reactive protein level to predict overall survival of patients with advanced urothelial carcinoma and its evaluation by decision curve analysis. <i>British Journal of Cancer</i> , 2012 , 107, 1031-6	8.7	27
52	A novel equation and nomogram including body weight for estimating prostate volumes in men with biopsy-proven benign prostatic hyperplasia. <i>Asian Journal of Andrology</i> , 2012 , 14, 703-7	2.8	6
51	C-reactive protein as a biomarker for urological cancers. <i>Nature Reviews Urology</i> , 2011 , 8, 659-66	5.5	78

50	A novel repeat biopsy nomogram based on three-dimensional extended biopsy. <i>Urology</i> , 2011 , 77, 915-206		4
49	Young age as favorable prognostic factor for cancer-specific survival in localized renal cell carcinoma. <i>Urology</i> , 2011 , 77, 842-7	1.6	20
48	Prognostic impact of C-reactive protein for determining overall survival of patients with castration-resistant prostate cancer treated with docetaxel. <i>Urology</i> , 2011 , 78, 1131-5	1.6	41
47	Longitudinal change in renal function after radical nephrectomy in Japanese patients with renal cortical tumors. <i>Journal of Urology</i> , 2011 , 185, 2066-71	2.5	24
46	Diffusion-weighted magnetic resonance imaging in the differentiation of angiomyolipoma with minimal fat from clear cell renal cell carcinoma. <i>International Journal of Urology</i> , 2011 , 18, 727-30	2.3	50
45	Phase-II trial of combination treatment of interferon- γ /cimetidine, cyclooxygenase-2 inhibitor and renin-angiotensin-system inhibitor (I-CCA therapy) for advanced renal cell carcinoma. <i>Cancer Science</i> , 2011 , 102, 137-43	6.9	29
44	Diagnostic performance of diffusion-weighted magnetic resonance imaging in bladder cancer: potential utility of apparent diffusion coefficient values as a biomarker to predict clinical aggressiveness. <i>European Radiology</i> , 2011 , 21, 2178-86	8	123
43	Treatment outcome and prognostic factors in renal cell cancer patients with bone metastasis. <i>Clinical and Experimental Metastasis</i> , 2011 , 28, 405-11	4.7	21
42	1172 ANTIBIOTIC PROPHYLAXIS IS NOT NECESSARY IN MINIMALLY INVASIVE SURGERY FOR RENAL AND ADRENAL TUMORS: A PROSPECTIVE STUDY OF 301 CONSECUTIVE PATIENTS. <i>Journal of Urology</i> , 2011 , 185,	2.5	1
41	Perioperative complications of radical cystectomy after induction chemoradiotherapy in bladder-sparing protocol against muscle-invasive bladder cancer: a single institutional retrospective comparative study with primary radical cystectomy. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 1373-9	2.8	24
40	Interactions between inducible nitric oxide synthase and cyclooxygenase-2 in response to ischaemia-reperfusion of rabbit bladder. <i>BJU International</i> , 2010 , 106, 716-22	5.6	4
39	Editorial comment to Nomogram to predict seminal vesicle invasion using the status of cancer at the base of the prostate on systematic biopsy. <i>International Journal of Urology</i> , 2010 , 17, 540	2.3	
38	Role of C-reactive protein as a biomarker for renal cell carcinoma. <i>Expert Review of Anticancer Therapy</i> , 2010 , 10, 1979-89	3.5	31
37	Initial experience of diffusion-weighted magnetic resonance imaging to assess therapeutic response to induction chemoradiotherapy against muscle-invasive bladder cancer. <i>Urology</i> , 2010 , 75, 387-91	1.6	85
36	Impact of C-reactive protein kinetics on survival of patients with metastatic renal cell carcinoma. <i>European Urology</i> , 2009 , 55, 1145-53	10.2	92
35	Discarding antimicrobial prophylaxis for transurethral resection of bladder tumor: a feasibility study. <i>International Journal of Urology</i> , 2009 , 16, 61-3	2.3	17
34	Safety of transperineal 14-core systematic prostate biopsy in diabetic men. <i>International Journal of Urology</i> , 2009 , 16, 930-5	2.3	21
33	Favourable outcomes of patients with clinical stage T3N0M0 bladder cancer treated with induction low-dose chemo-radiotherapy plus partial or radical cystectomy vs immediate radical cystectomy: a single-institutional retrospective comparative study. <i>BJU International</i> , 2009 , 104, 189-94	5.6	30

32	Development and external validation of a new outcome prediction model for patients with clear cell renal cell carcinoma treated with nephrectomy based on preoperative serum C-reactive protein and TNM classification: the TNM-C score. <i>Journal of Urology</i> , 2009 , 181, 1004-12; discussion 1012	2.5	80
31	Oncological outcome of minimum incision endoscopic radical nephrectomy for pathologically organ confined renal cell carcinoma. <i>International Journal of Urology</i> , 2008 , 15, 44-7	2.3	10
30	Collecting duct carcinoma with acquired cystic disease of the kidney in a long-term hemodialysis patient. <i>International Journal of Urology</i> , 2008 , 15, 93-5	2.3	3
29	Initial experience of functional imaging of upper urinary tract neoplasm by diffusion-weighted magnetic resonance imaging. <i>International Journal of Urology</i> , 2008 , 15, 140-3	2.3	34
28	Female urethral diverticular abscess clearly depicted by diffusion-weighted magnetic resonance imaging. <i>International Journal of Urology</i> , 2008 , 15, 460-1	2.3	6
27	Favorable response to combination treatment of cimetidine, cyclooxygenase-2 inhibitor and renin-angiotensin system inhibitor in metastatic renal cell carcinoma: Report of three cases. <i>International Journal of Urology</i> , 2008 , 15, 848-50	2.3	6
26	Three-dimensional 26-core biopsy-based patient selection criteria for nerve-sparing radical prostatectomy. <i>International Journal of Urology</i> , 2008 , 15, 1061-6	2.3	7
25	History of malignancy is a predictor of prostate cancer detection: incorporation into a pre-biopsy nomogram. <i>International Journal of Urology</i> , 2008 , 15, 1055-60	2.3	8
24	Reactive oxygen species mediate detrusor overactivity via sensitization of afferent pathway in the bladder of anaesthetized rats. <i>BJU International</i> , 2008 , 101, 775-80	5.6	77
23	C-reactive protein level predicts prognosis in patients with muscle-invasive bladder cancer treated with chemoradiotherapy. <i>BJU International</i> , 2008 , 101, 978-81	5.6	58
22	Development, validation, and head-to-head comparison of logistic regression-based nomograms and artificial neural network models predicting prostate cancer on initial extended biopsy. <i>European Urology</i> , 2008 , 54, 601-11	10.2	69
21	Low-dose chemoradiotherapy followed by partial or radical cystectomy against muscle-invasive bladder cancer: an intent-to-treat survival analysis. <i>Urology</i> , 2008 , 72, 384-8	1.6	29
20	Incidence of benign pathologic lesions at partial nephrectomy for presumed RCC renal masses: Japanese dual-center experience with 176 consecutive patients. <i>Urology</i> , 2008 , 72, 598-602	1.6	104
19	Prognostic impact of postoperative C-reactive protein level in patients with metastatic renal cell carcinoma undergoing cytoreductive nephrectomy. <i>Journal of Urology</i> , 2008 , 180, 515-9	2.5	56
18	External validation of the Mayo Clinic cancer specific survival score in a Japanese series of clear cell renal cell carcinoma. <i>Journal of Urology</i> , 2008 , 180, 1290-5; discussion 1295-6	2.5	32
17	Lymphovascular invasion is independently associated with poor prognosis in patients with localized upper urinary tract urothelial carcinoma treated surgically. <i>Journal of Urology</i> , 2007 , 178, 2291-6; discussion 2296	2.5	51
16	Increased preoperative serum C-reactive protein level predicts a poor prognosis in patients with localized renal cell carcinoma. <i>BJU International</i> , 2007 , 99, 77-80	5.6	90
15	The impact of preoperative serum C-reactive protein on the prognosis of patients with upper urinary tract urothelial carcinoma treated surgically. <i>BJU International</i> , 2007 , 100, 269-73	5.6	77

14	Spatial and isoform specific p63 expression in the male human urogenital tract. <i>Journal of Urology</i> , 2006 , 176, 2268-73	2.5	12
13	Loss of uroplakin III expression is associated with a poor prognosis in patients with urothelial carcinoma of the upper urinary tract. <i>BJU International</i> , 2006 , 97, 1322-6	5.6	22
12	Possible involvement of bone morphogenetic protein 2 in heterotopic ossification in metastatic lesion from urothelial carcinoma of bladder. <i>International Journal of Urology</i> , 2006 , 13, 1126-8	2.3	20
11	Prognostic significance of endothelial Per-Arnt-sim domain protein 1/hypoxia-inducible factor-2alpha expression in a subset of tumor associated macrophages in invasive bladder cancer. <i>Journal of Urology</i> , 2004 , 171, 1080-4	2.5	26
10	Localization of aquaporin-7 in human testis and ejaculated sperm: possible involvement in maintenance of sperm quality. <i>Journal of Urology</i> , 2004 , 172, 2073-6	2.5	57
9	Favorable outcome of preoperative low dose chemoradiotherapy against muscle-invasive bladder cancer. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 2003 , 26, 504-7	2.7	10
8	Increased expression of sialyl-Lewis A correlates with poor survival in upper urinary tract urothelial cancer patients. <i>Anticancer Research</i> , 2003 , 23, 3441-6	2.3	11
7	Impaired p63 expression associates with poor prognosis and uroplakin III expression in invasive urothelial carcinoma of the bladder. <i>Clinical Cancer Research</i> , 2003 , 9, 5501-7	12.9	107
6	Endoscope-assisted minilaparotomy (endoscopic minilaparotomy) for retroperitoneal Schwannoma: experience with three cases. <i>Japanese Journal of Clinical Oncology</i> , 2002 , 32, 177-80	2.8	12
5	Endoscopic minilaparotomy partial nephrectomy for solitary renal cell carcinoma smaller than 4 cm. <i>Japanese Journal of Clinical Oncology</i> , 2002 , 32, 417-21	2.8	8
4	Malignant Clear Cell Sugar Tumor of the Kidney: Clear Cell Variant of Epithelioid Angiomyolipoma. <i>Journal of Urology</i> , 2002 , 168, 2533-2534	2.5	35
3	IFN γ enhances the cytotoxicity of 5-fluorouracil against human bladder cancer cells through induction of PD-ECGF / TP. <i>Japanese Journal of Urology</i> , 2002 , 93, 323		
2	Platelet-derived endothelial cell growth factor/thymidine phosphorylase expression in primary bladder cancer. <i>Japanese Journal of Urology</i> , 2002 , 93, 377		
1	Malignant clear cell "sugar" tumor of the kidney: clear cell variant of epithelioid angiomyolipoma. <i>Journal of Urology</i> , 2002 , 168, 2533-4	2.5	9