

IsmaÄ°l DemÄ°r

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/587206/publications.pdf>

Version: 2024-02-01

55

papers

875

citations

394421

19

h-index

580821

25

g-index

56

all docs

56

docs citations

56

times ranked

623

citing authors

| # | ARTICLE | IF | CITATIONS |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1 | Isolation and characterization of entomopathogenic fungi from hazelnut-growing region of Turkey. BioControl, 2010, 55, 279-297. | 2.0 | 71 |
| 2 | Molecular Characterization and Virulence of Beauveria spp. from the Pine Processionary Moth, Thaumetopoea pityocampa (Lepidoptera: Thaumetopoeidae). Mycopathologia, 2010, 170, 269-277. | 3.1 | 45 |
| 3 | Biodiversity and pathogenicity of bacteria associated with the gut microbiota of beet armyworm, Spodoptera exigua Hübner (Lepidoptera: Noctuidae). Microbial Pathogenesis, 2018, 121, 350-358. | 2.9 | 33 |
| 4 | Evaluation of entomopathogenic fungi against the sycamore lace bug, <i>Corythucha ciliata</i> (Say) (Hemiptera: Tingidae). Turk Tarım Ve Ormancılık Dergisi/Turkish Journal of Agriculture and Forestry, 2013, 37, 595-603. | 2.1 | 32 |
| 5 | Culturable bacterial microbiota of <i>Plagiодera versicolora</i> (L.) (Coleoptera: Chrysomelidae) and virulence of the isolated strains. Folia Microbiologica, 2013, 58, 201-210. | 2.3 | 31 |
| 6 | An investigation on the bacterial flora of <i>Agriotes lineatus</i> (Coleoptera: Elateridae) and pathogenicity of the flora members. Crop Protection, 2012, 40, 1-7. | 2.1 | 30 |
| 7 | Molecular Characterization of Chitinase Genes from a Local Isolate of <i>Serratia marcescens</i> and Their Contribution to the Insecticidal Activity of <i>Bacillus thuringiensis</i> Strains. Current Microbiology, 2013, 67, 499-504. | 2.2 | 30 |
| 8 | Isolation and virulence of entomopathogenic fungi against the great spruce bark beetle, <i>Dendroctonus micans</i> (Kugelann) (Coleoptera: Scolytidae). Biocontrol Science and Technology, 2010, 20, 695-701. | 1.3 | 29 |
| 9 | Isolation, characterization and virulence of bacteria from <i>Ostrinia nubilalis</i> (Lepidoptera: Pyralidae). Biologia (Poland), 2012, 67, 767-776. | 1.5 | 29 |
| 10 | Screening of entomopathogenic fungi against the European spruce bark beetle, <i>Dendroctonus micans</i> (Coleoptera: Scolytidae). Biocontrol Science and Technology, 2010, 20, 3-11. | 1.3 | 28 |
| 11 | Molecular characterization, virulence and horizontal transmission of <i>Beauveria pseudobassiana</i> from <i>Dendroctonus micans</i> (<i>Kugelann</i>) (Coleoptera: Scolytidae). Journal of Applied Entomology, 2015, 139, 381-389. | 1.8 | 27 |
| 12 | Virulence and horizontal transmission of <i>Beauveria pseudobassiana</i> S.A. Rehner & Humber in <i>Ips sexdentatus</i> and <i>Ips typographus</i> (Coleoptera: Curculionidae). Turk Tarım Ve Ormancılık Dergisi/Turkish Journal of Agriculture and Forestry, 2016, 40, 241-248. | 2.1 | 26 |
| 13 | Investigations on bacteria as a potential biological control agent of summer chafer, <i>Amphimallon solstitiale</i> L. (Coleoptera: Scarabaeidae). Journal of Microbiology, 2005, 43, 463-8. | 2.8 | 26 |
| 14 | Isolation and identification of bacteria from Thaumetopoea pityocampa Den. and Schiff. (Lep.,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 22 and Biotechnology, 2008, 24, 3005-3015. | 3.6 | 25 |
| 15 | <i>Brevibacterium pityocampae</i> sp. nov., isolated from caterpillars of <i>Thaumetopoea pityocampa</i> (Lepidoptera, Thaumetopoeidae). International Journal of Systematic and Evolutionary Microbiology, 2010, 60, 312-316. | 1.7 | 24 |
| 16 | A new biopesticide from a local <i>Bacillus thuringiensis</i> var. <i>tenebrionis</i> (Xd3) against alder leaf beetle (Coleoptera: Chrysomelidae). World Journal of Microbiology and Biotechnology, 2017, 33, 95. | 3.6 | 23 |
| 17 | Isolation, characterization and pathogenicity of bacteria from <i>Rhynchites bacchus</i> (Coleoptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 and Biotechnology, 2008, 24, 3005-3015. | 3.6 | 22 |
| 18 | Investigating internal bacteria of <i>Spodoptera littoralis</i> (Boisd.) (Lepidoptera: Noctuidae) larvae and some <i>Bacillus</i> strains as biocontrol agents. Turk Tarım Ve Ormancılık Dergisi/Turkish Journal of Agriculture and Forestry, 2014, 38, 99-110. | 2.1 | 22 |

| # | ARTICLE | IF | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | Cloning and expression of chitinase A, B, and C (<i>chiA</i> , <i>chiB</i> , <i>chiC</i>) genes from <i>Serratia marcescens</i> originating from <i>Helicoverpa armigera</i> and determining their activities. <i>Turkish Journal of Biology</i> , 2015, 39, 78-87. | 0.8 | 22 |
| 20 | Microencapsulation of an indigenous isolate of <i>< i>Bacillus thuringiensis</i></i> by spray drying. <i>Journal of Microencapsulation</i> , 2019, 36, 1-9. | 2.8 | 22 |
| 21 | A Novel <i>cry2Ab</i> Gene from the Indigenous Isolate <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> . <i>Journal of Microbiology and Biotechnology</i> , 2012, 22, 133-140. | 2.1 | 19 |
| 22 | Isolation, characterization and virulence of entomopathogenic fungi from <i>Gryllotalpa gryllotalpa</i> (Orthoptera: Gryllotalpidae). <i>Applied Entomology and Zoology</i> , 2016, 51, 213-223. | 1.2 | 17 |
| 23 | Characterisation of three Alphabaculovirus isolates from the gypsy moth, <i>< i>Lymantria dispar</i> <i>dispar</i></i> (Lepidoptera: Erebidae), in Turkey. <i>Biocontrol Science and Technology</i> , 2018, 28, 107-121. | 1.3 | 15 |
| 24 | A new DNA polymerase I from <i>Geobacillus caldoxylosilyticus</i> TK4: cloning, characterization, and mutational analysis of two aromatic residues. <i>Applied Microbiology and Biotechnology</i> , 2009, 84, 105-117. | 3.6 | 14 |
| 25 | Isolation and identification of entomopathogenic nematodes (Nematoda: Rhabditida) from the Eastern Black Sea region and their biocontrol potential against <i>Melolontha melolontha</i> (Coleoptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 TF 2.1 13 2014, 38, 187-197. | 2.1 | 13 |
| 26 | Pathogenicity of selected entomopathogenic fungal isolates against the oak lace bug, <i>Corythucha arcuata</i> Say. (Hemiptera: Tingidae), under controlled conditions. <i>Turk Tarim Ve Ormancılık Dergisi/Turkish Journal of Agriculture and Forestry</i> , 2016, 40, 715-722. | 2.1 | 13 |
| 27 | Pine processionary moth (<i>Thaumetopoea pityocampa</i> , Lepidoptera: Thaumetopoeidae) larvae are highly susceptible to the entomopathogenic fungi <i>Metarhizium brunneum</i> and <i>Beauveria bassiana</i> . <i>Biocontrol Science and Technology</i> , 2017, 27, 1168-1179. | 1.3 | 13 |
| 28 | Development of <i>Beauveria bassiana</i> (Ascomycota: Hypocreales) as a mycoinsecticide to control green peach aphid, <i>Myzus persicae</i> (Homoptera: Aphididae) and investigation of its biocontrol potential. <i>Journal of Asia-Pacific Entomology</i> , 2022, 25, 101878. | 0.9 | 12 |
| 29 | A cytoplasmic polyhedrosis virus isolated from the pine processionary caterpillar, <i>Thaumetopoea pityocampa</i> . <i>Journal of Microbiology and Biotechnology</i> , 2007, 17, 632-7. | 2.1 | 12 |
| 30 | Bacterial isolates from <i>< i>Palomena prasina</i></i> (Hemiptera: Pentatomidae) include potential microbial control agents. <i>Biocontrol Science and Technology</i> , 2014, 24, 1039-1051. | 1.3 | 11 |
| 31 | Genome sequence analysis and organization of the <i>Hyphantria cunea</i> granulovirus (HycuGV-Hc1) from Turkey. <i>Genomics</i> , 2020, 112, 459-466. | 2.9 | 11 |
| 32 | Development of mycoinsecticide formulations with <i>< i>Beauveria bassiana</i></i> and <i>< i>Metarhizium brunneum</i></i> for the control of <i>< i>Orosanga japonica</i></i> (Hemiptera: Ricaniidae). <i>Annals of Applied Biology</i> , 2021, 179, 319-330. | 2.5 | 11 |
| 33 | A new entomopathogenic nematode species from Turkey, <i>Steinernema websteri</i> (Rhabditida: Tj ETQq1 1 0.784314 rgBT /Overlock 10 TF 0.8 10 | 0.8 | 10 |
| 34 | Identification and pathogenicity of bacteria in the Mediterranean corn borer <i>Sesamia nonagrioides</i> Lefebvre (Lepidoptera: Noctuidae). <i>Turkish Journal of Biology</i> , 2015, 39, 31-48. | 0.8 | 9 |
| 35 | Determination of fungal pathogens of <i>Hypera postica</i> (Gyllenhal) (Coleoptera: Curculionidae): isolation, characterization, and susceptibility. <i>Egyptian Journal of Biological Pest Control</i> , 2018, 28, . | 1.8 | 9 |
| 36 | Characterization of a new isolate of <i>Malacosoma neustria</i> nucleopolyhedrovirus (ManeNPV) from Turkey. <i>Turkish Journal of Biology</i> , 0, . | 0.8 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 37 | Highly effective bacterial agents against <i>Cimbex quadrimaculatus</i> (Hymenoptera: Cimbicidae): isolation of bacteria and their insecticidal activities. <i>World Journal of Microbiology and Biotechnology</i> , 2015, 31, 59-67. | 3.6 | 8 |
| 38 | Characterization of a <i>Betabaculovirus</i> from the fall webworm, <i>Hyphantria cunea</i> Drury. (Lepidoptera: Erebidae), in Turkey. <i>Biocontrol Science and Technology</i> , 2018, 28, 1178-1190. | 1.3 | 8 |
| 39 | A novel alphabaculovirus isolated from the cotton bollworm, <i>Helicoverpa armigera</i> (Hubner) (Lepidoptera: Noctuidae): characterization and pathogenicity. <i>Biologia (Poland)</i> , 2018, 73, 545-551. | 1.5 | 8 |
| 40 | Biodiversity and pathogenicity of entomopathogenic fungi associated with the lesser spruce sawfly, <i>Pristiphora abietina</i>. <i>Entomologia Experimentalis Et Applicata</i> , 2021, 169, 414-423. | 1.4 | 8 |
| 41 | First Record of <i>Steinernema kraussei</i> (Rhabditida: Steinernematidae) from Turkey and Its Virulence against <i>Agrotis segetum</i> (Lepidoptera: Noctuidae). <i>Journal of Nematology</i> , 2013, 45, 253-9. | 0.9 | 8 |
| 42 | Characterization of a nucleopolyhedrovirus variant of the gypsy moth, <i>Lymantria dispar</i> (Lepidoptera: Lymantriidae) in Turkey. <i>International Journal of Pest Management</i> , 2018, 64, 119-127. | 1.8 | 7 |
| 43 | Fungal pathogens of <i>Amphimallon solstitiale</i> Linnaeus, 1758 (Coleoptera: Scarabaeidae). <i>Turkiye Entomoloji Dergisi</i> , 0, , 375-384. | 0.6 | 7 |
| 44 | A highly effective nucleopolyhedrovirus against <i>Malacosoma</i> spp. (Lepidoptera: Lasiocampidae) from Turkey: isolation, characterization, phylogeny, and virulence. <i>Turk Tarım Ve Ormancılık Dergisi/Turkish Journal of Agriculture and Forestry</i> , 2014, 38, 462-470. | 2.1 | 6 |
| 45 | Efficacy of native entomopathogenic nematodes from Turkey against the alder leaf beetle, <i>Agelastica alni</i> L. (Coleoptera: Chrysomelidae), under laboratory conditions. <i>Egyptian Journal of Biological Pest Control</i> , 2018, 28, . | 1.8 | 3 |
| 46 | Improvement of delta-endotoxin production from local <i>Bacillus thuringiensis</i> Se13 using Taguchiâ€™s orthogonal array methodology. <i>Turkish Journal of Biochemistry</i> , 2018, 43, 662-670. | 0.5 | 3 |
| 47 | Insecticidal activities of wild type and recombinant invertebrate iridescent viruses on five common pests. <i>Turkiye Entomoloji Dergisi</i> , 0, , 365-373. | 0.6 | 3 |
| 48 | Molecular Characterization and Pathogenicity of <i>Beauveria bassiana</i> Isolated from <i>Rhynchites bacchus</i> L. (Coleoptera: Rhynchitidae). <i>Nevâ€hir Bilim Ve Teknoloji Dergisi</i> , 2015, 3, 33. | 0.1 | 2 |
| 49 | Complete genome sequence analysis of the <i>Malacosoma neustria</i> nucleopolyhedrovirus from Turkey. <i>Virus Genes</i> , 2018, 54, 706-718. | 1.6 | 2 |
| 50 | Stability of <i>Metarhizium anisopliae</i> (Hypocreales: Clavicipitaceae) isolates during repeated <i>in vitro</i> subculture and evaluation of an oil-in-water mycoinsecticide. <i>Canadian Entomologist</i> , 2022, 154, . | 0.8 | 2 |
| 51 | Comparison of DIG-11-dUTP utilization by <i>Geobacillus caldoxylosilyticus</i> TK4, <i>Mycobacterium tuberculosis</i> and <i>Escherichia coli</i> DNA polymerases. <i>World Journal of Microbiology and Biotechnology</i> , 2010, 26, 459-464. | 3.6 | 1 |
| 52 | bassianaâ€™nâ±n MolekÃ¼ller Karakterizasyonu ve Patojenitesi. <i>Nevâ€hir Bilim Ve Teknoloji Dergisi</i> , 2015, 3, 33-33. | 0.1 | 1 |
| 53 | Characterization of a novel baculovirus isolate from <i>Malacosoma neustria</i> (Linnaeus, 1758) (Lepidoptera: Lasiocampidae) in Samsun and its pathogenicity in different hosts. <i>Turkiye Entomoloji Dergisi</i> , 0, , 429-440. | 0.6 | 1 |
| 54 | Bioactivity of a betabaculovirus, <i>Hyphantria cunea</i> granulovirus, in six lepidopteran insects as potential hosts. <i>Turkiye Entomoloji Dergisi</i> , 2021, 45, 417-424. | 0.6 | 1 |

| # | ARTICLE | IF | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 55 | Effect of plant phenolic compounds on the hemocyte concentration and antioxidant enzyme activity in <i>Hyphantria cunea</i> (Drury, 1773) (Lepidoptera: Arctiidae) larvae infected by <i>Hyphantria cunea</i> granulovirus. <i>Turkiye Entomoloji Dergisi</i> , 2022, 46, 37-49. | 0.6 | 1 |