

# Walter S Speidl

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/5873293/walter-s-speidl-publications-by-year.pdf>

**Version:** 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

97  
papers

2,381  
citations

30  
h-index

45  
g-index

117  
ext. papers

2,751  
ext. citations

5.4  
avg, IF

4.12  
L-index

#	Paper	IF	Citations
97	Soluble neprilysin and survival in critically ill patients.. <i>ESC Heart Failure</i> , <b>2022</b> ,	3.7	1
96	Biomarkers of coagulation and fibrinolysis in acute myocardial infarction: a joint position paper of the Association for Acute CardioVascular Care and the European Society of Cardiology Working Group on Thrombosis. <i>European Heart Journal: Acute Cardiovascular Care</i> , <b>2021</b> , 10, 343-355	4.3	4
95	Innate Immune Training with Bacterial Extracts Enhances Lung Macrophage Recruitment to Protect from Betacoronavirus Infection. <i>Journal of Innate Immunity</i> , <b>2021</b> , 1-13	6.9	1
94	Circulating levels of proprotein convertase subtilisin/kexin type 9 (PCSK9) are associated with monocyte subsets in patients with stable coronary artery disease. <i>Journal of Clinical Lipidology</i> , <b>2021</b> , 15, 512-521	4.9	1
93	Alternative activation of human macrophages enhances tissue factor expression and production of extracellular vesicles. <i>Haematologica</i> , <b>2021</b> , 106, 454-463	6.6	8
92	Monocyte subsets predict mortality after cardiac arrest. <i>Journal of Leukocyte Biology</i> , <b>2021</b> , 109, 1139-1146	6.5	2
91	Death is associated to the type of drug-eluting stent in patients with left ventricular dysfunction and elevated natriuretic peptide levels. <i>Scientific Reports</i> , <b>2021</b> , 11, 2443	4.9	
90	Effects of Nicorandil on Inflammation, Apoptosis and Atherosclerotic Plaque Progression. <i>Biomedicines</i> , <b>2021</b> , 9,	4.8	8
89	Pharmacological inhibition of fatty acid oxidation reduces atherosclerosis progression by suppression of macrophage NLRP3 inflammasome activation. <i>Biochemical Pharmacology</i> , <b>2021</b> , 190, 1146-1154	6.34	3
88	The adipokine vaspin is associated with decreased coronary in-stent restenosis in vivo and inhibits migration of human coronary smooth muscle cells in vitro. <i>PLoS ONE</i> , <b>2020</b> , 15, e0232483	3.7	2
87	Epinephrine treatment but not time to ROSC is associated with intestinal injury in patients with cardiac arrest. <i>Resuscitation</i> , <b>2020</b> , 155, 32-38	4	2
86	The ISTH DIC score predicts outcome in non-septic patients admitted to a cardiovascular intensive care unit. <i>European Journal of Internal Medicine</i> , <b>2020</b> , 79, 37-42	3.9	3
85	The pro-inflammatory marker soluble suppression of tumorigenicity-2 (ST2) is reduced especially in diabetic morbidly obese patients undergoing bariatric surgery. <i>Cardiovascular Diabetology</i> , <b>2020</b> , 19, 26	8.7	5
84	Implementation of clinical practices and pathways optimizing ACS patients lipid management: Focus on eight European initiatives. <i>Atherosclerosis Supplements</i> , <b>2020</b> , 42, e59-e64	1.7	2
83	Lipoprotein(a) plasma levels are not associated with survival after acute coronary syndromes: An observational cohort study. <i>PLoS ONE</i> , <b>2020</b> , 15, e0227054	3.7	8
82	Toll-like receptor 2 and 9 expression on circulating neutrophils is associated with increased mortality in critically ill patients. <i>Shock</i> , <b>2020</b> , 54, 35-43	3.4	2
81	Neutrophil Extracellular Trap Degradation by Differently Polarized Macrophage Subsets. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>2020</b> , 40, 2265-2278	9.4	18

80	Interdependence of VA-ECMO output, pulmonary congestion and outcome after cardiac surgery. <i>European Journal of Internal Medicine</i> , <b>2020</b> , 81, 67-70	3.9	1
79	N-terminal pro-brain natriuretic peptide and high-sensitivity troponin T exhibit additive prognostic value for the outcome of critically ill patients. <i>European Heart Journal: Acute Cardiovascular Care</i> , <b>2020</b> , 9, 496-503	4.3	2
78	Release of mitochondrial DNA is associated with mortality in severe acute heart failure. <i>European Heart Journal: Acute Cardiovascular Care</i> , <b>2020</b> , 9, 419-428	4.3	5
77	Intestinal Fatty Acid Binding Protein is Associated With Mortality in Patients With Acute Heart Failure or Cardiogenic Shock. <i>Shock</i> , <b>2019</b> , 51, 410-415	3.4	8
76	Protease-Activated Receptors 1 and 3 are Differentially Expressed on Human Monocyte Subsets and are Upregulated by Lipopolysaccharide Ex Vivo and In Vivo. <i>Thrombosis and Haemostasis</i> , <b>2019</b> , 119, 1394-1402	7	1
75	Differential expression of Plg-R and its effects on migration of proinflammatory monocyte and macrophage subsets. <i>Blood</i> , <b>2019</b> , 134, 561-567	2.2	15
74	Neutrophil extracellular traps and monocyte subsets at the culprit lesion site of myocardial infarction patients. <i>Scientific Reports</i> , <b>2019</b> , 9, 16304	4.9	18
73	Growth differentiation factor-15 predicts poor survival after cardiac arrest. <i>Resuscitation</i> , <b>2019</b> , 143, 22-28	4	3
72	Predictive value of low interleukin-33 in critically ill patients. <i>Cytokine</i> , <b>2018</b> , 103, 109-113	4	13
71	Clopidogrel in Critically Ill Patients. <i>Clinical Pharmacology and Therapeutics</i> , <b>2018</b> , 103, 217-223	6.1	19
70	Pretreatment With Argon Protects Human Cardiac Myocyte-Like Progenitor Cells from Oxygen Glucose Deprivation-Induced Cell Death by Activation of AKT and Differential Regulation of Mapkinases. <i>Shock</i> , <b>2018</b> , 49, 556-563	3.4	9
69	Maternal serum mitochondrial DNA (mtDNA) levels are elevated in preeclampsia - A matched case-control study. <i>Pregnancy Hypertension</i> , <b>2018</b> , 14, 195-199	2.6	13
68	Anti-thrombotic and pro-fibrinolytic effects of levosimendan in human endothelial cells in vitro. <i>Vascular Pharmacology</i> , <b>2017</b> , 90, 44-50	5.9	8
67	Acetylsalicylic acid in critically ill patients: a cross-sectional and a randomized trial. <i>European Journal of Clinical Investigation</i> , <b>2017</b> , 47, 504-512	4.6	4
66	Copeptin Predicts Mortality in Critically Ill Patients. <i>PLoS ONE</i> , <b>2017</b> , 12, e0170436	3.7	9
65	PAI-1 (Plasminogen Activator Inhibitor-1) Expression Renders Alternatively Activated Human Macrophages Proteolytically Quiescent. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>2017</b> , 37, 1913-1922	9.4	18
64	Prasugrel in critically ill patients. <i>Thrombosis and Haemostasis</i> , <b>2017</b> , 117, 1582-1587	7	8
63	Monocyte subset distribution is associated with mortality in critically ill patients. <i>Thrombosis and Haemostasis</i> , <b>2016</b> , 116, 949-957	7	17

62	Expert position paper on prolonged dual antiplatelet therapy in secondary prevention following myocardial infarction. <i>Wiener Klinische Wochenschrift</i> , <b>2016</b> , 128, 450-7	2.3	2
61	Beneficial effects of levosimendan on survival in patients undergoing extracorporeal membrane oxygenation after cardiovascular surgery. <i>British Journal of Anaesthesia</i> , <b>2016</b> , 117, 52-8	5.4	39
60	Variation of lipoprotein(a) plasma levels after premature myocardial infarction. <i>International Journal of Cardiology</i> , <b>2015</b> , 186, 5-6	3.2	4
59	Levosimendan exerts anti-inflammatory effects on cardiac myocytes and endothelial cells in vitro. <i>Thrombosis and Haemostasis</i> , <b>2015</b> , 113, 350-62	7	21
58	Mitochondrial DNA and Toll-Like Receptor-9 Are Associated With Mortality in Critically Ill Patients. <i>Critical Care Medicine</i> , <b>2015</b> , 43, 2633-41	1.4	47
57	Association of small dense LDL serum levels and circulating monocyte subsets in stable coronary artery disease. <i>PLoS ONE</i> , <b>2015</b> , 10, e0123367	3.7	25
56	G-CSF Predicts Cardiovascular Events in Patients with Stable Coronary Artery Disease. <i>PLoS ONE</i> , <b>2015</b> , 10, e0142532	3.7	6
55	Monocyte subset distribution in patients with stable atherosclerosis and elevated levels of lipoprotein(a). <i>Journal of Clinical Lipidology</i> , <b>2015</b> , 9, 533-41	4.9	26
54	Premature myocardial infarction is strongly associated with increased levels of remnant cholesterol. <i>Journal of Clinical Lipidology</i> , <b>2015</b> , 9, 801-806.e1	4.9	24
53	An increase of interleukin-33 serum levels after coronary stent implantation is associated with coronary in-stent restenosis. <i>Cytokine</i> , <b>2014</b> , 67, 65-70	4	21
52	Glycoprotein 130 polymorphism predicts soluble glycoprotein 130 levels. <i>Metabolism: Clinical and Experimental</i> , <b>2014</b> , 63, 647-53	12.7	8
51	Soluble ST2 and interleukin-33 levels in coronary artery disease: relation to disease activity and adverse outcome. <i>PLoS ONE</i> , <b>2014</b> , 9, e95055	3.7	56
50	Small high-density lipoprotein is associated with monocyte subsets in stable coronary artery disease. <i>Atherosclerosis</i> , <b>2014</b> , 237, 589-96	3.1	37
49	An increase of VEGF plasma levels is associated with restenosis of drug-eluting stents. <i>EuroIntervention</i> , <b>2014</b> , 10, 224-30	3.1	13
48	Inflammation and coagulation in atherosclerosis. <i>Hamostaseologie</i> , <b>2013</b> , 33, 269-82	1.9	34
47	Rhabdomyolysis during therapeutic hypothermia in a patient after successful cardio-pulmonary resuscitation. <i>Resuscitation</i> , <b>2013</b> , 84, e79-80	4	5
46	Macrophages transmit potent proangiogenic effects of oxLDL in vitro and in vivo involving HIF-1 $\alpha$ activation: a novel aspect of angiogenesis in atherosclerosis. <i>Journal of Cardiovascular Translational Research</i> , <b>2013</b> , 6, 558-69	3.3	50
45	Novel small leucine-rich repeat protein podocan is a negative regulator of migration and proliferation of smooth muscle cells, modulates neointima formation, and is expressed in human atheroma. <i>Circulation</i> , <b>2013</b> , 128, 2351-63	16.7	18

44	Polymorphism of the complement 5 gene and cardiovascular outcome in patients with atherosclerosis. <i>European Journal of Clinical Investigation</i> , <b>2012</b> , 42, 921-6	4.6	19
43	Premature myocardial infarction is associated with low serum levels of Wnt-1. <i>Atherosclerosis</i> , <b>2012</b> , 222, 251-6	3.1	35
42	High soluble Fas and soluble Fas Ligand serum levels before stent implantation are protective against restenosis. <i>Thrombosis and Haemostasis</i> , <b>2011</b> , 105, 883-91	7	8
41	Atherosclerosis and complement: anaphylatoxin C5a as a new risk marker and therapeutic target. <i>Clinical Lipidology</i> , <b>2011</b> , 6, 123-126		2
40	MUW researcher of the month. <i>Wiener Klinische Wochenschrift</i> , <b>2011</b> , 123, 757-757	2.3	
39	The prognostic impact of soluble apoptosis-stimulating fragment on mortality in patients with carotid atherosclerosis. <i>Stroke</i> , <b>2011</b> , 42, 2465-70	6.7	6
38	The complement component C5a is present in human coronary lesions in vivo and induces the expression of MMP-1 and MMP-9 in human macrophages in vitro. <i>FASEB Journal</i> , <b>2011</b> , 25, 35-44	0.9	70
37	Recombinant apolipoprotein A-I Milano rapidly reverses aortic valve stenosis and decreases leaflet inflammation in an experimental rabbit model. <i>European Heart Journal</i> , <b>2010</b> , 31, 2049-57	9.5	46
36	Coronary late lumen loss of drug eluting stents is associated with increased serum levels of the complement components C3a and C5a. <i>Atherosclerosis</i> , <b>2010</b> , 208, 285-9	3.1	32
35	Increased restenosis rate after implantation of drug-eluting stents in patients with elevated serum activity of matrix metalloproteinase-2 and -9. <i>JACC: Cardiovascular Interventions</i> , <b>2010</b> , 3, 90-7	5	31
34	Up-regulation of reverse cholesterol transport key players and rescue from global inflammation by ApoA-I(Milano). <i>Journal of Cellular and Molecular Medicine</i> , <b>2009</b> , 13, 3226-35	5.6	41
33	Human cardiac fibroblasts express B-type natriuretic peptide: fluvastatin ameliorates its up-regulation by interleukin-1alpha, tumour necrosis factor-alpha and transforming growth factor-beta. <i>Journal of Cellular and Molecular Medicine</i> , <b>2009</b> , 13, 4415-21	5.6	22
32	Quantification of serial changes in plaque burden using multi-detector computed tomography in experimental atherosclerosis. <i>Atherosclerosis</i> , <b>2009</b> , 202, 185-91	3.1	17
31	Safe and sustained overexpression of functional apolipoprotein A-I/high-density lipoprotein in apolipoprotein A-I-null mice by muscular adeno-associated viral serotype 8 vector gene transfer. <i>Journal of Cardiovascular Pharmacology</i> , <b>2009</b> , 54, 405-11	3.1	10
30	Thrombin induces the expression of oncostatin M via AP-1 activation in human macrophages: a link between coagulation and inflammation. <i>Blood</i> , <b>2009</b> , 114, 2812-8	2.2	37
29	Plasminogen activator inhibitor-1 predicts coronary in-stent restenosis of drug-eluting stents. <i>Journal of Thrombosis and Haemostasis</i> , <b>2008</b> , 6, 508-13	15.4	23
28	Rapid change in plaque size, composition, and molecular footprint after recombinant apolipoprotein A-I Milano (ETC-216) administration: magnetic resonance imaging study in an experimental model of atherosclerosis. <i>Journal of the American College of Cardiology</i> , <b>2008</b> , 51, 1104-9	15.1	110
27	Clopidogrel pretreatment abolishes increase of PAI-1 after coronary stent implantation. <i>Thrombosis Research</i> , <b>2008</b> , 123, 79-84	8.2	4

26	In human macrophages the complement component C5a induces the expression of oncostatin M via AP-1 activation. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>2008</b> , 28, 498-503	9.4	36
25	Complement component C5a predicts restenosis after superficial femoral artery balloon angioplasty. <i>Journal of Endovascular Therapy</i> , <b>2007</b> , 14, 62-9	2.5	20
24	Vascular endothelial growth factor is induced by the inflammatory cytokines interleukin-6 and oncostatin m in human adipose tissue in vitro and in murine adipose tissue in vivo. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>2007</b> , 27, 1587-95	9.4	81
23	Early metoprolol administration before coronary reperfusion results in increased myocardial salvage: analysis of ischemic myocardium at risk using cardiac magnetic resonance. <i>Circulation</i> , <b>2007</b> , 115, 2909-16	16.7	118
22	Premature compared with late onset of coronary artery disease: young patients show a severe defect in fibrinolytic response to venous occlusion. <i>Blood Coagulation and Fibrinolysis</i> , <b>2007</b> , 18, 165-71	1	10
21	Mild hyperhomocysteinemia is associated with a decreased fibrinolytic activity in patients after ST-elevation myocardial infarction. <i>Thrombosis Research</i> , <b>2007</b> , 119, 331-6	8.2	10
20	Simvastatin suppresses endotoxin-induced upregulation of toll-like receptors 4 and 2 in vivo. <i>Atherosclerosis</i> , <b>2006</b> , 189, 408-13	3.1	115
19	Monocyte chemoattractant protein (MCP-1) is expressed in human cardiac cells and is differentially regulated by inflammatory mediators and hypoxia. <i>FEBS Letters</i> , <b>2006</b> , 580, 3532-8	3.8	32
18	The complement component C5a induces the expression of plasminogen activator inhibitor-1 in human macrophages via NF-kappaB activation. <i>Journal of Thrombosis and Haemostasis</i> , <b>2006</b> , 4, 1790-7	15.4	56
17	The estrogen metabolite 17beta-dihydroequilenin counteracts interleukin-1-induced expression of inflammatory mediators in human endothelial cells in vitro via NF-B pathway. <i>Thrombosis and Haemostasis</i> , <b>2006</b> , 95, 107-116	7	18
16	The estrogen metabolite 17beta-dihydroequilenin counteracts interleukin-1alpha induced expression of inflammatory mediators in human endothelial cells in vitro via NF-kappaB pathway. <i>Thrombosis and Haemostasis</i> , <b>2006</b> , 95, 107-16	7	7
15	Can a commercial diagnostic ultrasound device accelerate thrombolysis? An in vitro skull model. <i>Stroke</i> , <b>2005</b> , 36, 124-8	6.7	79
14	Complement component C5a predicts future cardiovascular events in patients with advanced atherosclerosis. <i>European Heart Journal</i> , <b>2005</b> , 26, 2294-9	9.5	97
13	The gp130 ligand oncostatin M regulates tissue inhibitor of metalloproteinases-1 through ERK1/2 and p38 in human adult cardiac myocytes and in human adult cardiac fibroblasts: a possible role for the gp130/gp130 ligand system in the modulation of extracellular matrix degradation in the human heart. <i>Journal of Molecular and Cellular Cardiology</i> , <b>2005</b> , 39, 515-24	5.8	30
12	An increase of C-reactive protein is associated with enhanced activation of endogenous fibrinolysis at baseline but an impaired endothelial fibrinolytic response after venous occlusion. <i>Journal of the American College of Cardiology</i> , <b>2005</b> , 45, 30-4	15.1	34
11	Simvastatin blunts endotoxin-induced tissue factor in vivo. <i>Circulation</i> , <b>2005</b> , 111, 1841-6	16.7	123
10	Catecholamines potentiate LPS-induced expression of MMP-1 and MMP-9 in human monocytes and in the human monocytic cell line U937: possible implications for peri-operative plaque instability. <i>FASEB Journal</i> , <b>2004</b> , 18, 603-5	0.9	60
9	Prostaglandin E1 induces vascular endothelial growth factor-1 in human adult cardiac myocytes but not in human adult cardiac fibroblasts via a cAMP-dependent mechanism. <i>Journal of Molecular and Cellular Cardiology</i> , <b>2004</b> , 36, 539-46	5.8	27

8	Pre-existing anticardiolipin antibodies and development of restenosis after coronary balloon angioplasty: predictive value for a complicated course?. <i>Blood Coagulation and Fibrinolysis</i> , <b>2004</b> , 15, 311-6	1.6	3
7	Glycoprotein 130 ligand oncostatin-M induces expression of vascular endothelial growth factor in human adult cardiac myocytes. <i>Cardiovascular Research</i> , <b>2003</b> , 59, 628-38	9.9	32
6	2MHz ultrasound enhances t-PA-mediated thrombolysis: comparison of continuous versus pulsed ultrasound and standing versus travelling acoustic waves. <i>Thrombosis and Haemostasis</i> , <b>2003</b> , 89, 583-589	7	44
5	Circulating t-PA antigen predicts major adverse coronary events in patients with stable coronary artery disease--a 13-year follow-up. <i>Thrombosis and Haemostasis</i> , <b>2003</b> , 90, 344-50	7	11
4	Polymorphic membrane protein (PMP) 20 and PMP 21 of Chlamydia pneumoniae induce proinflammatory mediators in human endothelial cells in vitro by activation of the nuclear factor-kappaB pathway. <i>Journal of Infectious Diseases</i> , <b>2003</b> , 188, 108-13	7	38
3	2MHz ultrasound enhances t-PA-mediated thrombolysis: comparison of continuous versus pulsed ultrasound and standing versus travelling acoustic waves. <i>Thrombosis and Haemostasis</i> , <b>2003</b> , 89, 583-9	7	14
2	Chlamydia pneumoniae in carotid artery atherosclerosis: a comparison of its presence in atherosclerotic plaque, healthy vessels, and circulating leukocytes from the same individuals. <i>Stroke</i> , <b>2002</b> , 33, 2756-61	6.7	41
1	High-sensitivity C-reactive protein in the prediction of coronary events in patients with premature coronary artery disease. <i>American Heart Journal</i> , <b>2002</b> , 144, 449-55	4.9	52