

Domenico Restuccia

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5876384/publications.pdf>

Version: 2024-02-01

77
papers

2,478
citations

186265
28
h-index

214800
47
g-index

78
all docs

78
docs citations

78
times ranked

2071
citing authors

#	ARTICLE	IF	CITATIONS
1	Contribution of different somatosensory afferent input to subcortical somatosensory evoked potentials in humans. <i>Clinical Neurophysiology</i> , 2021, 132, 2357-2364.	1.5	2
2	Somatosensory high frequency oscillations: A useful tool to analyze cortical excitability?. <i>Clinical Neurophysiology</i> , 2020, 131, 468-469.	1.5	1
3	High frequency oscillations after median nerve stimulations in healthy children and adolescents. <i>International Journal of Developmental Neuroscience</i> , 2017, 61, 68-72.	1.6	2
4	Cortical hyperexcitability in healthy children: evidence from habituation and recovery cycle phenomena of somatosensory evoked potentials. <i>Developmental Medicine and Child Neurology</i> , 2016, 58, 855-860.	2.1	7
5	Abnormality of Auditory Mismatch Negativity in Depression and Its Dependence on Stimulus Intensity. <i>Clinical EEG and Neuroscience</i> , 2016, 47, 105-112.	1.7	15
6	Somatosensory high frequency oscillations: A useful tool to analyze dynamic changes in somatosensory pathways?. <i>Clinical Neurophysiology</i> , 2015, 126, 1643-1644.	1.5	4
7	Copper deficiency myelopathy: A report of two cases. <i>Journal of Spinal Cord Medicine</i> , 2015, 38, 559-562.	1.4	29
8	Auditory stimulation enhances thalamic somatosensory high-frequency oscillations in healthy humans: a neurophysiological marker of cross-sensory sensitization?. <i>European Journal of Neuroscience</i> , 2015, 41, 1079-1085.	2.6	5
9	Restless Legs Syndrome and lateralized periodic movements due to a spinal schwannoma. <i>Sleep and Biological Rhythms</i> , 2015, 13, 106-108.	1.0	0
10	Patterns of habituation and clinical fluctuations in migraine. <i>Cephalalgia</i> , 2014, 34, 201-210.	3.9	24
11	A novel TSC2 mutation causing tuberless tuberous sclerosis. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2014, 23, 580-582.	2.0	2
12	Different levels of cortical excitability reflect clinical fluctuations in migraine. <i>Cephalalgia</i> , 2013, 33, 1035-1047.	3.9	20
13	Unmasking of presynaptic cutaneous HFOs burst by DBS lead recordings. <i>Clinical Neurophysiology</i> , 2012, 123, 842-844.	1.5	5
14	Somatosensory High Frequency Oscillations reflect clinical fluctuations in migraine. <i>Clinical Neurophysiology</i> , 2012, 123, 2050-2056.	1.5	31
15	Commentary. <i>Journal of Neurosciences in Rural Practice</i> , 2012, 3, 423-4.	0.8	0
16	High-frequency oscillations after median-nerve stimulation do not undergo habituation: A new insight on their functional meaning?. <i>Clinical Neurophysiology</i> , 2011, 122, 148-152.	1.5	16
17	Tired legs—a gut diagnosis. <i>Lancet, The</i> , 2010, 376, 1798.	13.7	7
18	State Estimation, Response Prediction, and Cerebellar Sensory Processing for Behavioral Control. <i>Cerebellum</i> , 2009, 8, 399-402.	2.5	60

#	ARTICLE	IF	CITATIONS
19	Somatosensory mismatch negativity in healthy children. <i>Developmental Medicine and Child Neurology</i> , 2009, 51, 991-998.	2.1	28
20	Dissociated effects of quiet stance on standard and high-frequency (600Hz) lower limb somatosensory evoked potentials. <i>Clinical Neurophysiology</i> , 2008, 119, 1408-1418.	1.5	9
21	Seeing the pain of others while being in pain: A laser-evoked potentials study. <i>NeuroImage</i> , 2008, 40, 1419-1428.	4.2	104
22	High-frequency ECoG oscillations in the site of onset of epileptic seizures during sleep. <i>Sleep Medicine</i> , 2007, 8, 96-97.	1.6	6
23	Giant subcortical high-frequency SEPs in idiopathic generalized epilepsy: A protective mechanism against seizures?. <i>Clinical Neurophysiology</i> , 2007, 118, 60-68.	1.5	15
24	Parallel spinal pathways generate the middle-latency N1 and the late P2 components of the laser evoked potentials. <i>Clinical Neurophysiology</i> , 2007, 118, 1097-1104.	1.5	28
25	Inhibitory effect of voluntary movement preparation on cutaneous heat pain and laser-evoked potentials. <i>European Journal of Neuroscience</i> , 2007, 25, 1900-1907.	2.6	39
26	Modulation of high-frequency (600Hz) somatosensory-evoked potentials after rTMS of the primary sensory cortex. <i>European Journal of Neuroscience</i> , 2007, 26, 2349-2358.	2.6	39
27	Distraction affects frontal alpha rhythms related to expectancy of pain: An EEG study. <i>NeuroImage</i> , 2006, 31, 1268-1277.	4.2	43
28	Modulation of laser-evoked potentials by experimental cutaneous tonic pain. <i>Neuroscience</i> , 2006, 140, 1301-1310.	2.3	10
29	Cerebellar damage impairs detection of somatosensory input changes. A somatosensory mismatch-negativity study. <i>Brain</i> , 2006, 130, 276-287.	7.6	115
30	Inhibitory effect of capsaicin evoked trigeminal pain on warmth sensation and warmth evoked potentials. <i>Experimental Brain Research</i> , 2005, 160, 29-37.	1.5	28
31	Cerebellar information flow in the thalamus: implications for cortical functions. <i>Thalamus & Related Systems</i> , 2005, 3, 141.	0.5	4
32	Abnormal cortical pain processing in patients with cardiac syndrome X. <i>European Heart Journal</i> , 2005, 26, 975-982.	2.2	74
33	Increase of brain-stem high-frequency SEP subcomponents during light sleep in seizure-free epileptic patients. <i>Clinical Neurophysiology</i> , 2005, 116, 1774-1778.	1.5	9
34	Abnormal cortical pain processing in patients with cardiac syndrome X. <i>International Congress Series</i> , 2005, 1278, 393-396.	0.2	0
35	Segmental inhibition of cutaneous heat sensation and of laser-evoked potentials by experimental muscle pain. <i>Neuroscience</i> , 2005, 136, 301-309.	2.3	21
36	Attentional load of the primary task influences the frontal but not the temporal generators of mismatch negativity. <i>Cognitive Brain Research</i> , 2005, 25, 891-899.	3.0	63

#	ARTICLE	IF	CITATIONS
37	Functional assessment of A? and C fibers in patients with Fabry's disease. <i>Muscle and Nerve</i> , 2004, 30, 708-713.	2.2	36
38	Different neuronal contribution to N20 somatosensory evoked potential and to CO2 laser evoked potentials: an intracerebral recording study. <i>Clinical Neurophysiology</i> , 2004, 115, 211-216.	1.5	38
39	Reduction in amplitude of the subcortical low- and high-frequency somatosensory evoked potentials during voluntary movement: an intracerebral recording study. <i>Clinical Neurophysiology</i> , 2004, 115, 104-111.	1.5	33
40	Parietal generators of low- and high-frequency MN (median nerve) SEPs: data from intracortical human recordings. <i>Clinical Neurophysiology</i> , 2004, 115, 647-657.	1.5	16
41	Influence of modafinil on somatosensory input processing in the human brain-stem. <i>Clinical Neurophysiology</i> , 2004, 115, 919-926.	1.5	9
42	Brain-stem components of high-frequency somatosensory evoked potentials are modulated by arousal changes: nasopharyngeal recordings in healthy humans. <i>Clinical Neurophysiology</i> , 2004, 115, 1392-1398.	1.5	29
43	Abnormal gating of somatosensory inputs in essential tremor. <i>Clinical Neurophysiology</i> , 2003, 114, 120-129.	1.5	27
44	Influence of cholinergic circuitries in generation of high-frequency somatosensory evoked potentials. <i>Clinical Neurophysiology</i> , 2003, 114, 1538-1548.	1.5	45
45	Short-term plastic changes of the human nociceptive system following acute pain induced by capsaicin. <i>Clinical Neurophysiology</i> , 2003, 114, 1879-1890.	1.5	53
46	The human supplementary motor area-proper does not receive direct somatosensory inputs from the periphery: data from stereotactic depth somatosensory evoked potential recordings. <i>Neuroscience Letters</i> , 2003, 344, 161-164.	2.1	21
47	Reduced habituation to experimental pain in migraine patients: a CO2 laser evoked potential study. <i>Pain</i> , 2003, 105, 57-64.	4.2	205
48	Attention-related modifications of ultra-late CO2 laser evoked potentials to human trigeminal nerve stimulation. <i>Neuroscience Letters</i> , 2002, 329, 329-333.	2.1	34
49	Dissociated changes of somatosensory evoked low-frequency scalp responses and 600 Hz bursts after single-dose administration of lorazepam. <i>Brain Research</i> , 2002, 946, 1-11.	2.2	23
50	Modality-related scalp responses after electrical stimulation of cutaneous and muscular upper limb afferents in humans. <i>Muscle and Nerve</i> , 2002, 26, 44-54.	2.2	20
51	Source generators of the early somatosensory evoked potentials to tibial nerve stimulation: an intracerebral and scalp recording study. <i>Clinical Neurophysiology</i> , 2001, 112, 1999-2006.	1.5	27
52	Inhibition of biceps brachii muscle motor area by painful heat stimulation of the skin. <i>Experimental Brain Research</i> , 2001, 139, 168-172.	1.5	50
53	Central scalp projection of the N30 SEP source activity after median nerve stimulation. , 2000, 23, 353-360.		21
54	Scalp distribution of the earliest cortical somatosensory evoked potential to tibial nerve stimulation: proposal of a new recording montage. <i>Clinical Neurophysiology</i> , 2000, 111, 1469-1477.	1.5	13

#	ARTICLE	IF	CITATIONS
55	Sources of cortical responses to painful CO ₂ laser skin stimulation of the hand and foot in the human brain. <i>Clinical Neurophysiology</i> , 2000, 111, 1103-1112.	1.5	125
56	Unmasking of an early laser evoked potential by a point localization task. <i>Clinical Neurophysiology</i> , 2000, 111, 1927-1933.	1.5	25
57	Anatomic Origin of P13 and P14 Scalp Far-Field Potentials. <i>Journal of Clinical Neurophysiology</i> , 2000, 17, 246-257.	1.7	19
58	Different contribution of joint and cutaneous inputs to early scalp somatosensory evoked potentials. , 1999, 22, 910-919.		18
59	Effect of movement on dipolar source activities of somatosensory evoked potentials. <i>Muscle and Nerve</i> , 1999, 22, 1510-1519.	2.2	30
60	Inhibition of the human primary motor area by painful heat stimulation of the skin. <i>Clinical Neurophysiology</i> , 1999, 110, 1475-1480.	1.5	110
61	Functional involvement of cerebral cortex in duchenne muscular dystrophy. , 1998, 21, 662-664.		29
62	Dipolar sources of the early scalp somatosensory evoked potentials to upper limb stimulation. <i>Experimental Brain Research</i> , 1998, 120, 306-315.	1.5	60
63	Dissociation induced by voluntary movement between two different components of the centro-parietal P40 SEP to tibial nerve stimulation. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1998, 108, 190-198.	2.0	21
64	The scalp to earlobe montage as standard in routine SEP recording. Comparison with the non-cephalic reference in patients with lesions of the upper cervical cord. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1998, 108, 414-422.	2.0	14
65	Selective abnormality of the N13 spinal SEP to dermatomal stimulation in patients with cervical monoradiculopathy. <i>Neurophysiologie Clinique</i> , 1998, 28, 221-229.	2.2	6
66	Giant central N20-P22 with normal area 3b N20-P20: an argument in favour of an area 3a generator of early median nerve cortical SEPs?. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1997, 104, 60-67.	2.0	20
67	The pathophysiology of giant SEPs in cortical myoclonus: a scalp topography and dipolar source modelling study. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1997, 104, 122-131.	2.0	47
68	Dipolar generators of the early scalp somatosensory evoked potentials to tibial nerve stimulation in human subjects. <i>Neuroscience Letters</i> , 1997, 238, 49-52.	2.1	25
69	Abnormalities of somatosensory and motor evoked potentials in adrenomyeloneuropathy: Comparison with magnetic resonance imaging and clinical findings. , 1997, 20, 1249-1257.		13
70	Brain-stem somatosensory dysfunction in a case of long-standing left hemispherectomy with removal of the left thalamus: a nasopharyngeal and scalp SEP study. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1996, 100, 184-188.	2.0	19
71	Origin and distribution of P13 and P14 far-field potentials after median nerve stimulation. Scalp, nasopharyngeal and neck recording in healthy subjects and in patients with cervical and cervico-medullary lesions. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1995, 96, 371-384.	2.0	43
72	Recovery after surgery of the spinal N24 SEP in dural arteriovenous malformation of the dorsal cord. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1995, 96, 479-482.	2.0	4

#	ARTICLE	IF	CITATIONS
73	The role of upper limb somatosensory evoked potentials in the management of cervical spondylotic myelopathy: preliminary data. <i>Electroencephalography and Clinical Neurophysiology - Evoked Potentials</i> , 1994, 92, 502-509.	2.0	20
74	Somatosensory and motor evoked potentials in the assessment of cerebrotendinous xanthomatosis before and after treatment with chenodeoxycholic acid: a preliminary study. <i>Journal of the Neurological Sciences</i> , 1992, 112, 139-146.	0.6	11
75	Cisplatin neuropathy: clinical course and neurophysiological findings. <i>Journal of Neurology</i> , 1992, 239, 199-204.	3.6	103
76	Inadequacy of the forehead reference montage for detecting abnormalities of the spinal N13 SEP in cervical cord lesions. <i>Electroencephalography and Clinical Neurophysiology</i> , 1991, 79, 448-456.	0.3	40
77	THE CONTRIBUTION OF MEDIAN NERVE SEPs IN THE FUNCTIONAL ASSESSMENT OF THE CERVICAL SPINAL CORD IN SYRINGOMYELIA. <i>Brain</i> , 1991, 114, 361-379.	7.6	111