

Marco Vugman Wainstein

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5881457/marco-vugman-wainstein-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

48
papers

5,644
citations

15
h-index

52
g-index

52
ext. papers

6,660
ext. citations

6.9
avg, IF

4.76
L-index

#	Paper	IF	Citations
48	Ezetimibe Added to Statin Therapy after Acute Coronary Syndromes. <i>New England Journal of Medicine</i> , 2015 , 372, 2387-97	59.2	2371
47	Alogliptin after acute coronary syndrome in patients with type 2 diabetes. <i>New England Journal of Medicine</i> , 2013 , 369, 1327-35	59.2	1858
46	Effect of darapladib on major coronary events after an acute coronary syndrome: the SOLID-TIMI 52 randomized clinical trial. <i>JAMA - Journal of the American Medical Association</i> , 2014 , 312, 1006-15	27.4	285
45	Effect of vitamins C and E on progression of transplant-associated arteriosclerosis: a randomised trial. <i>Lancet, The</i> , 2002 , 359, 1108-13	4.0	265
44	No-reflow is an independent predictor of death and myocardial infarction after percutaneous coronary intervention. <i>American Heart Journal</i> , 2003 , 145, 42-6	4.9	193
43	Effect of aleglitazar on cardiovascular outcomes after acute coronary syndrome in patients with type 2 diabetes mellitus: the AleCardio randomized clinical trial. <i>JAMA - Journal of the American Medical Association</i> , 2014 , 311, 1515-25	27.4	177
42	Role of endothelin-1 in the active constriction of human atherosclerotic coronary arteries. <i>Circulation</i> , 2001 , 104, 1114-8	16.7	130
41	Endothelial dysfunction as a predictor of cardiovascular disease in type 1 diabetes. <i>World Journal of Diabetes</i> , 2015 , 6, 679-92	4.7	50
40	Elevated serum interleukin-6 is predictive of coronary artery disease in intermediate risk overweight patients referred for coronary angiography. <i>Diabetology and Metabolic Syndrome</i> , 2017 , 9, 67	5.6	41
39	Erectile dysfunction and coronary artery disease: an association of higher risk in younger men. <i>Journal of Sexual Medicine</i> , 2011 , 8, 1445-53	1.1	33
38	Comparison of neutrophil-to-lymphocyte ratio and mean platelet volume in the prediction of adverse events after primary percutaneous coronary intervention in patients with ST-elevation myocardial infarction. <i>Atherosclerosis</i> , 2018 , 274, 212-217	3.1	25
37	Association between myeloperoxidase polymorphisms and its plasma levels with severity of coronary artery disease. <i>Clinical Biochemistry</i> , 2010 , 43, 57-62	3.5	23
36	Rationale, design, and baseline characteristics of the Acetylcystein for Contrast-Induced nephropathy (ACT) Trial: a pragmatic randomized controlled trial to evaluate the efficacy of acetylcysteine for the prevention of contrast-induced nephropathy. <i>Trials</i> , 2009 , 10, 38	2.8	18
35	HOMA-IR is associated with significant angiographic coronary artery disease in non-diabetic, non-obese individuals: a cross-sectional study. <i>Diabetology and Metabolic Syndrome</i> , 2015 , 7, 100	5.6	17
34	Impact of coronary endothelial function on the progression of cardiac transplant-associated arteriosclerosis: effect of anti-oxidant vitamins C and E. <i>Journal of Heart and Lung Transplantation</i> , 2006 , 25, 426-33	5.8	17
33	Plasma endothelin-1 levels after coronary stenting in humans. <i>American Journal of Cardiology</i> , 2003 , 92, 1211-4	3	13
32	Carbonyl groups: Bridging the gap between sleep disordered breathing and coronary artery disease. <i>Free Radical Research</i> , 2010 , 44, 907-12	4	10

31	Cost-effectiveness of sirolimus-eluting stents in percutaneous coronary interventions in Brazil. <i>Arquivos Brasileiros De Cardiologia</i> , 2007 , 88, 464-74	1.2	10
30	Simplifying contrast-induced acute kidney injury prediction after primary percutaneous coronary intervention: the age, creatinine and ejection fraction score. <i>Cardiovascular Intervention and Therapeutics</i> , 2018 , 33, 224-231	2.5	9
29	Atherosclerosis and acute arterial thrombosis in rabbits: a model using balloon desendothelization without dietary intervention. <i>Brazilian Journal of Medical and Biological Research</i> , 1997 , 30, 415-7	2.8	9
28	Inflammatory and oxidative stress markers after intravenous insulin in percutaneous coronary intervention with stent in type 2 diabetes mellitus: a randomized controlled trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, 478-85	5.6	8
27	Coronary stent implantation may seal the inflammatory response in patients with acute coronary syndromes. <i>International Journal of Cardiology</i> , 2008 , 130, 503-4	3.2	8
26	Rescue percutaneous coronary intervention following coronary artery bypass graft--a descriptive analysis of the changing interface between interventional cardiologist and cardiac surgeon. <i>Clinical Cardiology</i> , 2002 , 25, 280-6	3.3	8
25	Severity of obstructive sleep apnea and extension of coronary artery disease. <i>Sleep and Breathing</i> , 2019 , 23, 747-752	3.1	8
24	Obstructive sleep apnea, detected by the Berlin Questionnaire: an associated risk factor for coronary artery disease. <i>Cadernos De Saude Publica</i> , 2012 , 28, 1530-8	3.2	7
23	Lack of association between plasma myeloperoxidase levels and angiographic severity of coronary artery disease in patients with acute coronary syndrome. <i>Inflammation Research</i> , 2011 , 60, 137-42	7.2	7
22	Elevated neutrophil-to-lymphocyte ratio can predict procedural adverse events in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. <i>Coronary Artery Disease</i> , 2019 , 30, 20-25	1.4	6
21	Oxidized-LDL and paraoxonase-1 as biomarkers of coronary artery disease in patients with sleep-disordered breathing. <i>Current Medicinal Chemistry</i> , 2012 , 19, 4359-66	4.3	6
20	Comparison of Two Risk Models in Predicting the Incidence of Contrast-Induced Nephropathy after Percutaneous Coronary Intervention. <i>Journal of Interventional Cardiology</i> , 2016 , 29, 447-453	1.8	6
19	Syntax Score and Major Adverse Cardiac Events in Patients with Suspected Coronary Artery Disease: Results from a Cohort Study in a University-Affiliated Hospital in Southern Brazil. <i>Arquivos Brasileiros De Cardiologia</i> , 2016 , 107, 207-215	1.2	5
18	The Effects of a Flavonoid-Rich Diet on Oxidative Stress, Inflammation, and Lipid Profile after Elective Percutaneous Coronary Intervention: A Randomized Clinical Trial. <i>Preventive Nutrition and Food Science</i> , 2018 , 23, 108-114	2.4	2
17	Comparison of Admission Lung Ultrasound and Left Ventricular End-Diastolic Pressure in Patients Undergoing Primary Percutaneous Coronary Intervention. <i>Circulation: Cardiovascular Imaging</i> , 2021 , 14, e011641	3.9	2
16	Guidewire Self-Extrusion After Entrapment of Distal Protection Device During Saphenous Vein Graft Angioplasty. <i>JACC: Cardiovascular Interventions</i> , 2017 , 10, e3-e5	5	1
15	Ultrasound-guided antecubital vein approach for right heart catheterisation in a Brazilian tertiary centre. <i>Open Heart</i> , 2020 , 7, e001181	3	1
14	Two HEmostasis Methods After Transradial Catheterization: THEMATIC - protocol for a randomized clinical trial. <i>Revista Gaucha De Enfermagem / EENFUFGRS</i> , 2018 , 39, e20170257	0.5	1

13	What seals inflammation in acute coronary syndromes?. <i>International Journal of Cardiology</i> , 2009 , 135, 108	3.2	1
12	Mechanical plaque sealing in patients with acute coronary syndromes. <i>International Journal of Cardiology</i> , 2009 , 135, 105-106	3.2	1
11	Two HEmostasis Methods After Transradial Catheterization: THEMATIC Randomized Clinical Trial. <i>Journal of Cardiovascular Nursing</i> , 2020 , 35, 217-222	2.1	1
10	Diagnostic Accuracy of Perfusional Computed Tomography in Moderate Coronary Stenosis: Comparison With Fractional Flow Reserve. <i>Critical Pathways in Cardiology</i> , 2020 , 19, 9-13	1.3	1
9	Temporal pattern of neutrophil-to-lymphocyte ratio in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. <i>Coronary Artery Disease</i> , 2019 , 30, 631-633	1.4	1
8	Long-term Pattern of Red Cell Distribution Width in Patients With ST-Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. <i>Critical Pathways in Cardiology</i> , 2020 , 19, 43-48	1.3	0
7	Serum ferritin levels may have a pro-atherosclerotic role in coronary artery disease patients with sleep disordered breathing. <i>Journal of Applied Biomedicine</i> , 2015 , 13, 289-298	0.6	
6	Inter-observer variation of Syntax score among cardiac surgeons, clinical and interventional cardiologists. <i>Therapeutic Advances in Cardiovascular Disease</i> , 2020 , 14, 1753944720924254	3.4	
5	Right heart catheterization through ultrasound-guided antecubital vein approach. <i>Revista Brasileira De Cardiologia Invasiva (English Edition)</i> , 2016 , 24, 35-37		
4	Cardiovascular outcomes in patients treated with primary percutaneous coronary intervention in a general tertiary hospital. <i>Revista Brasileira De Cardiologia Invasiva (English Edition)</i> , 2016 , 24, 4-8		
3	Reply to "Mean platelet volume may not have a role in the prediction of adverse events after percutaneous coronary intervention in patients with ST-elevation myocardial infarction". <i>Atherosclerosis</i> , 2018 , 276, 206-207	3.1	
2	Unprotected left main bifurcation restenosis treated with a 2-stent technique. <i>Catheterization and Cardiovascular Interventions</i> , 2013 , 82, E200-5	2.7	
1	Long-term complication after LM bifurcation treatment. <i>Catheterization and Cardiovascular Interventions</i> , 2010 , 75, 1045-9	2.7	