

# Angelo Contarino

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/5882689/angelo-contarino-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

25  
papers

2,374  
citations

14  
h-index

25  
g-index

25  
ext. papers

2,508  
ext. citations

8.7  
avg, IF

4.11  
L-index

#	Paper	IF	Citations
25	Morphine reduces the interest for natural rewards.. <i>Psychopharmacology</i> , <b>2022</b> , 1	4.7	1
24	Dopaminergic neuromodulation of prefrontal cortex activity requires the NMDA receptor coagonist d-serine. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2021</b> , 118,	11.5	3
23	Sex-linked roles of the CRF and the CRF receptor in social behavior. <i>Journal of Neuroscience Research</i> , <b>2020</b> , 98, 1561-1574	4.4	2
22	Long-lasting pseudo-social aggressive behavior in opiate-withdrawn mice. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2020</b> , 97, 109780	5.5	9
21	The CRF receptor mediates social behavior deficits induced by opiate withdrawal. <i>Journal of Neuroscience Research</i> , <b>2020</b> ,	4.4	1
20	Corticotropin-releasing factor receptor 2-deficiency eliminates social behaviour deficits and vulnerability induced by cocaine. <i>British Journal of Pharmacology</i> , <b>2018</b> , 175, 1504-1518	8.6	8
19	CRF receptor-deficiency increases cocaine reward. <i>Neuropharmacology</i> , <b>2017</b> , 117, 41-48	5.5	12
18	The CRF1 and the CRF2 receptor mediate recognition memory deficits and vulnerability induced by opiate withdrawal. <i>Neuropharmacology</i> , <b>2016</b> , 105, 500-507	5.5	8
17	CRF2 Receptor Deficiency Eliminates the Long-Lasting Vulnerability of Motivational States Induced by Opiate Withdrawal. <i>Neuropsychopharmacology</i> , <b>2015</b> , 40, 1990-2000	8.7	10
16	CRF1 receptor-deficiency induces anxiety-like vulnerability to cocaine. <i>Psychopharmacology</i> , <b>2014</b> , 231, 3965-72	4.7	6
15	CRF $\beta$ receptor-deficiency reduces recognition memory deficits and vulnerability to stress induced by cocaine withdrawal. <i>International Journal of Neuropsychopharmacology</i> , <b>2014</b> , 17, 1969-79	5.8	17
14	The corticotropin-releasing factor receptor-2 mediates the motivational effect of opiate withdrawal. <i>Neuropharmacology</i> , <b>2013</b> , 73, 41-7	5.5	10
13	Increased motivation to eat in opiate-withdrawn mice. <i>Psychopharmacology</i> , <b>2012</b> , 221, 675-84	4.7	24
12	Disruption of the CRF(2) receptor pathway decreases the somatic expression of opiate withdrawal. <i>Neuropsychopharmacology</i> , <b>2008</b> , 33, 2878-87	8.7	40
11	Disruption of the CRF/CRF1 receptor stress system exacerbates the somatic signs of opiate withdrawal. <i>Neuron</i> , <b>2007</b> , 53, 577-89	13.9	44
10	Gender- and morphine dose-linked expression of spontaneous somatic opiate withdrawal in mice. <i>Behavioural Brain Research</i> , <b>2006</b> , 170, 110-8	3.4	54
9	The corticotropin-releasing factor receptor-1 pathway mediates the negative affective states of opiate withdrawal. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2005</b> , 102, 18649-54	11.5	87

8	Lack of reward and locomotor stimulation induced by heroin in mu-opioid receptor-deficient mice. <i>European Journal of Pharmacology</i> , <b>2002</b> , 446, 103-9	5.3	62
7	Urocortin-deficient mice show hearing impairment and increased anxiety-like behavior. <i>Nature Genetics</i> , <b>2002</b> , 31, 363-9	36.3	152
6	Automated assessment of conditioning parameters for context and cued fear in mice. <i>Learning and Memory</i> , <b>2002</b> , 9, 89-96	2.8	25
5	Mice deficient for corticotropin-releasing hormone receptor-2 display anxiety-like behaviour and are hypersensitive to stress. <i>Nature Genetics</i> , <b>2000</b> , 24, 410-4	36.3	729
4	Dissociation of locomotor activation and suppression of food intake induced by CRF in CRFR1-deficient mice. <i>Endocrinology</i> , <b>2000</b> , 141, 2698-702	4.8	90
3	Reduced anxiety-like and cognitive performance in mice lacking the corticotropin-releasing factor receptor 1. <i>Brain Research</i> , <b>1999</b> , 835, 1-9	3.7	212
2	Corticotropin releasing factor receptor 1-deficient mice display decreased anxiety, impaired stress response, and aberrant neuroendocrine development. <i>Neuron</i> , <b>1998</b> , 20, 1093-102	13.9	751
1	Conditioned place preference: no tolerance to the rewarding properties of morphine. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , <b>1997</b> , 355, 589-94	3.4	17