

Valeria A Lozano

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5889663/valeria-a-lozano-publications-by-year.pdf>

Version: 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

21
papers

431
citations

13
h-index

20
g-index

21
ext. papers

464
ext. citations

5.2
avg, IF

3.87
L-index

#	Paper	IF	Citations
21	Statistics and Food Quality 2021 , 362-386		
20	Determination of ochratoxin A in coffee and tea samples by coupling second-order multivariate calibration and fluorescence spectroscopy. <i>Talanta</i> , 2020 , 219, 121288	6.2	4
19	Simultaneous determination of quality parameters in yerba mate (<i>Ilex paraguariensis</i>) samples by application of near-infrared (NIR) spectroscopy and partial least squares (PLS). <i>LWT - Food Science and Technology</i> , 2020 , 126, 109290	5.4	14
18	On second-order calibration based on multivariate curve resolution in the presence of highly overlapped profiles. <i>Analytica Chimica Acta</i> , 2020 , 1096, 53-60	6.6	7
17	Comparative chemometric analysis of fluorescence and near infrared spectroscopies for authenticity confirmation and geographical origin of Argentinean extra virgin olive oils. <i>Food Control</i> , 2019 , 96, 22-28	6.2	34
16	Development of a second-order standard addition fluorescence method for the direct determination of riboflavin in human urine samples without previous clean up and separation steps. <i>Microchemical Journal</i> , 2017 , 133, 60-66	4.8	7
15	Simultaneous determination of urea herbicides in water and soil samples based on second-order photoinduced fluorescence data. <i>Analytical Methods</i> , 2016 , 8, 7396-7404	3.2	6
14	Green analytical determination of emerging pollutants in environmental waters using excitation-emission photoinduced fluorescence data and multivariate calibration. <i>Talanta</i> , 2015 , 134, 215-223	6.2	28
13	Novel augmented parallel factor model for four-way calibration of high-performance liquid chromatography fluorescence excitation-emission data. <i>Chemometrics and Intelligent Laboratory Systems</i> , 2015 , 141, 1-11	3.8	21
12	A second-order fluorimetric approach based on a boron dipyrromethene tetraamide derivative for Hg(II) chemosensing in water and fish samples. <i>Analytical Methods</i> , 2014 , 6, 8535-8543	3.2	3
11	Characterization of sildenafil citrate tablets of different sources by near infrared chemical imaging and chemometric tools. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2013 , 85, 207-12	3.5	19
10	Second-order advantage with excitation-emission photoinduced fluorimetry for the determination of the antiepileptic carbamazepine in environmental waters. <i>Analytica Chimica Acta</i> , 2013 , 782, 37-45	6.6	13
9	Four-way multivariate calibration using ultra-fast high-performance liquid chromatography with fluorescence excitation-emission detection. Application to the direct analysis of chlorophylls a and b and pheophytins a and b in olive oils. <i>Chemometrics and Intelligent Laboratory Systems</i> , 2013 , 125, 121-131	3.8	38
8	A novel nylon membrane-rhodamine 6G spirocyclic phenylthiosemicarbazide derivative system as a fluorimetric probe for mercury(II) ion. <i>Analytical Methods</i> , 2012 , 4, 2002	3.2	18
7	Second-order analyte quantitation under identical profiles in one data dimension. A dependency-adapted partial least-squares/residual bilinearization method. <i>Analytical Chemistry</i> , 2010 , 82, 4510-9	7.8	27
6	A novel second-order standard addition analytical method based on data processing with multidimensional partial least-squares and residual bilinearization. <i>Analytica Chimica Acta</i> , 2009 , 651, 165-72	6.6	30
5	Standard addition analysis of fluoroquinolones in human serum in the presence of the interferent salicylate using lanthanide-sensitized excitation-time decay luminescence data and multivariate curve resolution. <i>Talanta</i> , 2009 , 77, 1715-23	6.2	47

4	Chemometric tools for the characterisation of honey produced in La Pampa, Argentina, from their elemental content, using inductively coupled plasma optical emission spectrometry (ICP-OES). <i>Journal of Apicultural Research</i> , 2008 , 47, 102-107	2	3
3	Three-way partial least-squares/residual bilinearization study of second-order lanthanide-sensitized luminescence excitation-time decay data: analysis of benzoic acid in beverage samples. <i>Analytica Chimica Acta</i> , 2008 , 610, 186-95	6.6	35
2	Second-order advantage from kinetic-spectroscopic data matrices in the presence of extreme spectral overlapping A multivariate curve resolution--alternating least-squares approach. <i>Analytica Chimica Acta</i> , 2008 , 614, 46-57	6.6	42
1	Simultaneous determination of sorbic and benzoic acids in commercial juices using the PLS-2 multivariate calibration method and validation by high performance liquid chromatography. <i>Talanta</i> , 2007 , 73, 282-6	6.2	35