Manli Na

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5892399/publications.pdf Version: 2024-02-01



Μάνιι Να

#	Article	IF	CITATIONS
1	Linkage between endosomal escape of LNP-mRNA and loading into EVs for transport to other cells. Nature Communications, 2019, 10, 4333.	12.8	211
2	Human skin commensals augment Staphylococcus aureus pathogenesis. Nature Microbiology, 2018, 3, 881-890.	13.3	80
3	Staphylokinase Control of <i>Staphylococcus aureus</i> Biofilm Formation and Detachment Through Host Plasminogen Activation. Journal of Infectious Diseases, 2016, 213, 139-148.	4.0	61
4	IL-1 Receptor Antagonist Treatment Aggravates Staphylococcal Septic Arthritis and Sepsis in Mice. PLoS ONE, 2015, 10, e0131645.	2.5	40
5	CTLA4 Immunoglobulin but Not Anti–Tumor Necrosis Factor Therapy Promotes Staphylococcal Septic Arthritis in Mice. Journal of Infectious Diseases, 2015, 212, 1308-1316.	4.0	32
6	Glioma Cell Proliferation Controlled by ERK Activity-Dependent Surface Expression of PDGFRA. PLoS ONE, 2014, 9, e87281.	2.5	31
7	Deficiency of the Complement Component 3 but Not Factor B Aggravates Staphylococcus aureus Septic Arthritis in Mice. Infection and Immunity, 2016, 84, 930-939.	2.2	30
8	Gene-Viral Cancer Therapy Using Dual-Regulated Oncolytic Adenovirus with Antiangiogenesis Gene for Increased Efficacy. Molecular Cancer Research, 2008, 6, 568-575.	3.4	29
9	Tissue Plasminogen Activator Coating on Implant Surfaces Reduces Staphylococcus aureus Biofilm Formation. Applied and Environmental Microbiology, 2016, 82, 394-401.	3.1	25
10	The YIN and YANG of lipoproteins in developing and preventing infectious arthritis by Staphylococcus aureus. PLoS Pathogens, 2019, 15, e1007877.	4.7	25
11	Concomitant use of Ad5/35 chimeric oncolytic adenovirus with TRAIL gene and taxol produces synergistic cytotoxicity in gastric cancer cells. Cancer Letters, 2009, 284, 141-148.	7.2	23
12	Galectin-3 Is a Target for Proteases Involved in the Virulence of Staphylococcus aureus. Infection and Immunity, 2017, 85, .	2.2	23
13	Radiological features of experimental staphylococcal septic arthritis by micro computed tomography scan. PLoS ONE, 2017, 12, e0171222.	2.5	20
14	Fiber Mediated Receptor Masking in Non-Infected Bystander Cells Restricts Adenovirus Cell Killing Effect but Promotes Adenovirus Host Co-Existence. PLoS ONE, 2009, 4, e8484.	2.5	18
15	The role of Staphylococcus aureus lipoproteins in hematogenous septic arthritis. Scientific Reports, 2020, 10, 7936.	3.3	17
16	Tofacitinib treatment aggravates Staphylococcus aureus septic arthritis, but attenuates sepsis and enterotoxin induced shock in mice. Scientific Reports, 2020, 10, 10891.	3.3	16
17	The Expression of von Willebrand Factor-Binding Protein Determines Joint-Invading Capacity of Staphylococcus aureus, a Core Mechanism of Septic Arthritis. MBio, 2020, 11, .	4.1	14
18	Staphylococcus aureus lipoproteins promote abscess formation in mice, shielding bacteria from immune killing. Communications Biology, 2021, 4, 432.	4.4	14

Manli Na

#	Article	IF	CITATIONS
19	RAGE Deficiency Impairs Bacterial Clearance in Murine Staphylococcal Sepsis, but Has No Significant Impact on Staphylococcal Septic Arthritis. PLoS ONE, 2016, 11, e0167287.	2.5	9
20	Lack of Receptor for Advanced Glycation End Products Leads to Less Severe Staphylococcal Skin Infection but More Skin Abscesses and Prolonged Wound Healing. Journal of Infectious Diseases, 2018, 218, 791-800.	4.0	8
21	Design of Ad5F35 vectors for coordinated dual gene expression in candidate human hematopoietic stem cells. Experimental Hematology, 2010, 38, 446-452.	0.4	6
22	Both anti-TNF and CTLA4 Ig treatments attenuate the disease severity of staphylococcal dermatitis in mice. PLoS ONE, 2017, 12, e0173492.	2.5	5
23	Adenovirus assembly is impaired by BMI1-related histone deacetylase activity. Virology, 2014, 456-457, 227-237.	2.4	0