Eric Yung

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5902866/publications.pdf Version: 2024-02-01



FRIC YUNC

#	Article	IF	CITATIONS
1	c-MYC interacts with INI1/hSNF5 and requires the SWI/SNF complex for transactivation function. Nature Genetics, 1999, 22, 102-105.	21.4	348
2	In-depth characterization of the microRNA transcriptome in a leukemia progression model. Genome Research, 2008, 18, 1787-1797.	5.5	162
3	MN1 overexpression induces acute myeloid leukemia in mice and predicts ATRA resistance in patients with AML. Blood, 2007, 110, 1639-1647.	1.4	133
4	Inhibition of HIV-1 virion production by a transdominant mutant of integrase interactor 1. Nature Medicine, 2001, 7, 920-926.	30.7	124
5	The methyltransferase G9a regulates HoxA9-dependent transcription in AML. Genes and Development, 2014, 28, 317-327.	5.9	121
6	Unraveling the crucial roles of <i>Meis1</i> in leukemogenesis and normal hematopoiesis. Genes and Development, 2007, 21, 2845-2849.	5.9	87
7	Cell of Origin in AML: Susceptibility to MN1-Induced Transformation Is Regulated by the MEIS1/AbdB-like HOX Protein Complex. Cancer Cell, 2011, 20, 39-52.	16.8	76
8	Specificity of Interaction of INI1/hSNF5 with Retroviral Integrases and Its Functional Significance. Journal of Virology, 2004, 78, 2222-2231.	3.4	66
9	Linkage of Meis1 leukemogenic activity to multiple downstream effectors including Trib2 and Ccl3. Experimental Hematology, 2008, 36, 845-859.	0.4	56
10	HIV-1 replication in cell lines harboring INI1/hSNF5 mutations. Retrovirology, 2006, 3, 56.	2.0	29
11	Linkage of the potent leukemogenic activity of Meis1 to cell-cycle entry and transcriptional regulation of cyclin D3. Blood, 2010, 115, 4071-4082.	1.4	28
12	Constitutive IRF8 expression inhibits AML by activation of repressed immune response signaling. Leukemia, 2015, 29, 157-168.	7.2	25
13	Delineating domains and functions of NUP98 contributing to the leukemogenic activity of NUP98-HOX fusions. Leukemia Research, 2011, 35, 545-550.	0.8	17
14	Cell Fate Decisions in Malignant Hematopoiesis: Leukemia Phenotype Is Determined by Distinct Functional Domains of the MN1 Oncogene. PLoS ONE, 2014, 9, e112671.	2.5	15
15	Elucidating the importance and regulation of key enhancers for human MEIS1 expression. Leukemia, 2022, 36, 1980-1989.	7.2	6
16	Selective B cell depletion upon intravenous infusion of replication-incompetent anti-CD19 CAR lentivirus. Molecular Therapy - Methods and Clinical Development, 2022, 26, 4-14.	4.1	5