

# Anders Christensson

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5902922/publications.pdf>

Version: 2024-02-01

89  
papers

5,078  
citations

147566

31  
h-index

88477

70  
g-index

91  
all docs

91  
docs citations

91  
times ranked

5009  
citing authors

#	ARTICLE	IF	CITATIONS
1	Early life factors in relation to albuminuria and estimated glomerular filtration rate based on cystatin C and creatinine in adults from a Swedish population-based cohort study. <i>Journal of Nephrology</i> , 2022, 35, 889-900.	0.9	9
2	Impaired selective renal filtration captured by eGFR <sub>cysC</sub> /eGFR <sub>crea</sub> ratio is associated with mortality in a population based cohort of older women. <i>Scientific Reports</i> , 2022, 12, 1273.	1.6	6
3	Associations between long-term exposure to low-level air pollution and risk of chronic kidney disease—findings from the Malmö Diet and Cancer cohort. <i>Environment International</i> , 2022, 160, 107085.	4.8	18
4	Updated Pathways in Cardiorenal Continuum after Kidney Transplantation. <i>Transplantation</i> , 2022, 3, 156-168.	0.3	0
5	Adult-onset diabetes in Middle Eastern immigrants to Sweden: Novel subgroups and diabetic complications—The All New Diabetes in Scania cohort diabetic complications and ethnicity. <i>Diabetes/Metabolism Research and Reviews</i> , 2021, 37, e3419.	1.7	21
6	Proteins linked to atherosclerosis and cell proliferation are associated with the shrunken pore syndrome in heart failure patients. <i>Proteomics - Clinical Applications</i> , 2021, 15, e2000089.	0.8	11
7	FC 059EARLY LIFE FACTORS AND ADULT KIDNEY FUNCTION ESTIMATED BY CYSTATIN C AND CREATININE GLOMERULAR FILTRATION RATE EQUATIONS AND ALBUMINURIA: A SWEDISH COHORT STUDY. <i>Nephrology Dialysis Transplantation</i> , 2021, 36, .	0.4	0
8	MO071PROTEINS LINKED TO ATHEROSCLEROSIS AND CELL PROLIFERATION ARE ASSOCIATED WITH SHRUNKEN PORE SYNDROME IN HEART FAILURE PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2021, 36, .	0.4	0
9	MO131THE SHRUNKEN PORE SYNDROME IS ASSOCIATED WITH POOR PROGNOSIS AND LOWER QUALITY OF LIFE IN HEART FAILURE PATIENTS- THE HARVEST-MALMÖ– STUDY. <i>Nephrology Dialysis Transplantation</i> , 2021, 36, .	0.4	0
10	Multiple-Biomarker Panel Estimated GFR Is Not Optimal or Cost-Effective. <i>American Journal of Kidney Diseases</i> , 2021, 77, 823.	2.1	1
11	Secondary hyperparathyroidism, weight loss, and longer term mortality in haemodialysis patients: results from the DOPPS. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2021, 12, 855-865.	2.9	18
12	Potential relationship between eGFR <sub>cystatin C</sub> /eGFR <sub>creatinine</sub> ratio and glomerular basement membrane thickness in diabetic kidney disease. <i>Physiological Reports</i> , 2021, 9, e14939.	0.7	15
13	The Shrunken pore syndrome is associated with poor prognosis and lower quality of life in heart failure patients: the HARVEST—Malmö study. <i>ESC Heart Failure</i> , 2021, 8, 3577-3586.	1.4	13
14	Pro-Enkephalin and its association with renal function in Middle Eastern immigrants and native Swedes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2021, 81, 573-578.	0.6	1
15	The risk of chronic kidney disease in relation to anthropometric measures of obesity: A Swedish cohort study. <i>BMC Nephrology</i> , 2021, 22, 330.	0.8	5
16	Growth differentiation factor-15 and incident chronic kidney disease: a population-based cohort study. <i>BMC Nephrology</i> , 2021, 22, 351.	0.8	9
17	Plasma kidney injury molecule-1 (p-KIM-1) levels and deterioration of kidney function over 16 years. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 265-273.	0.4	43
18	Kidney function and its association to imminent, short- and long-term fracture risk—a longitudinal study in older women. <i>Osteoporosis International</i> , 2020, 31, 97-107.	1.3	6

#	ARTICLE	IF	CITATIONS
19	Patterns of urinary albumin and IgM associate with markers of vascular ageing in young to middle-aged individuals in the MalmÅ¶ offspring study. BMC Cardiovascular Disorders, 2020, 20, 358.	0.7	4
20	<p>Comparison of Hemodialysis Using a Medium Cutoff Dialyzer versus Hemodiafiltration: A Controlled Cross-Over Study</p>. International Journal of Nephrology and Renovascular Disease, 2020, Volume 13, 273-280.	0.8	13
21	Longitudinal Changes in Kidney Function Estimated from Cystatin C and Its Association with Mortality in Elderly Women. Nephron, 2020, 144, 290-298.	0.9	3
22	Shrunken pore syndrome and mortality: a cohort study of patients with measured GFR and known comorbidities. Scandinavian Journal of Clinical and Laboratory Investigation, 2020, 80, 412-422.	0.6	40
23	Impact of longer term phosphorus control on cardiovascular mortality in hemodialysis patients using an area under the curve approach: results from the DOPPS. Nephrology Dialysis Transplantation, 2020, 35, 1794-1801.	0.4	37
24	Low lung function and the risk of incident chronic kidney disease in the MalmÅ¶ Preventive Project cohort. BMC Nephrology, 2020, 21, 124.	0.8	7
25	Genetic Predisposition for Renal Dysfunction and Incidence of CKD in the MalmÅ¶ Diet and Cancer Study. Kidney International Reports, 2019, 4, 1143-1151.	0.4	4
26	Echocardiographic Findings in Patients with Mild to Moderate Chronic Kidney Disease without Symptomatic Heart Failure: A Population-Based Study. CardioRenal Medicine, 2019, 9, 284-296.	0.7	6
27	SP603 Understanding the impact on mortality of long term serum phosphorus control using a 6 month area under the curve approach in the International Dialysis Outcomes and Practice Patterns Study (DOPPS). Nephrology Dialysis Transplantation, 2019, 34, .	0.4	0
28	CKD: A Call for an Age-Adapted Definition. Journal of the American Society of Nephrology: JASN, 2019, 30, 1785-1805.	3.0	198
29	FP424 RENAL FUNCTION AMONG MIDDLE EASTERN IMMIGRANTS IN SWEDEN AND ITS ASSOCIATION TO PRO-ENKEPHALIN. Nephrology Dialysis Transplantation, 2019, 34, .	0.4	0
30	Complement C3 and incident hospitalization due to chronic kidney disease: a population-based cohort study. BMC Nephrology, 2019, 20, 61.	0.8	12
31	Shrunken Pore Syndrome Is Associated With Increased Levels of Atherosclerosis-Promoting Proteins. Kidney International Reports, 2019, 4, 67-79.	0.4	43
32	Cystatin C and creatinine-based eGFR levels and their correlation to long-term morbidity and mortality in older adults. Aging Clinical and Experimental Research, 2019, 31, 1461-1469.	1.4	4
33	Plasma copeptin as a predictor of kidney disease. Nephrology Dialysis Transplantation, 2019, 34, 74-82.	0.4	25
34	Alterations in Serum MicroRNA Profile During Hemodialysis - Potential Biological Implications. Cellular Physiology and Biochemistry, 2018, 46, 793-801.	1.1	2
35	The Impact of the Glomerular Filtration Rate on the Human Plasma Proteome. Proteomics - Clinical Applications, 2018, 12, e1700067.	0.8	37
36	Impact of Kidney Transplantation on Reproductive Hormone Levels in Males: A Longitudinal Study. Nephron, 2018, 138, 192-201.	0.9	18

#	ARTICLE	IF	CITATIONS
37	Blood Lead Levels and Decreased Kidney Function in a Population-Based Cohort. <i>American Journal of Kidney Diseases</i> , 2018, 72, 381-389.	2.1	120
38	Mineral and bone disorder management in hemodialysis patients: comparing PTH control practices in Japan with Europe and North America: the Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>BMC Nephrology</i> , 2018, 19, 253.	0.8	15
39	SP564METABOLIC PATHWAYS ANALYSED FROM SERUM MICRORNA PROFILE DURING HEMODIALYSIS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i538-i538.	0.4	0
40	Soluble Urokinase-type Plasminogen Activator Receptor (suPAR) and Impaired Kidney Function in the Population-based Malmö Diet and Cancer Study. <i>Kidney International Reports</i> , 2017, 2, 239-247.	0.4	33
41	Combining Cystatin C and Creatinine Yields a Reliable Glomerular Filtration Rate Estimation in Older Adults in Contrast to <sup>125</sup> I-Trace Protein and <sup>125</sup> I-Microglobulin. <i>Nephron</i> , 2017, 137, 29-37.	0.9	27
42	Renal function and its association with blood pressure in Middle Eastern immigrants and native Swedes. <i>Journal of Hypertension</i> , 2017, 35, 2493-2500.	0.3	9
43	MicroRNA-155 and Anti-Müllerian Hormone: New Potential Markers of Subfertility in Men with Chronic Kidney Disease. <i>Nephron Extra</i> , 2017, 7, 33-41.	1.1	10
44	High Level of Fasting Plasma Proenkephalin-A Predicts Deterioration of Kidney Function and Incidence of CKD. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 291-303.	3.0	29
45	Prevalence and determinants of differences in cystatin C and creatinine-based estimated glomerular filtration rate in community-dwelling older adults: a cross-sectional study. <i>BMC Nephrology</i> , 2017, 18, 350.	0.8	22
46	Longitudinal Assessment of PTH in Community-Dwelling Older Women—Elevations Are Not Associated With Mortality. <i>Journal of the Endocrine Society</i> , 2017, 1, 615-624.	0.1	7
47	Increased Levels of Copeptin, a Surrogate Marker of Arginine Vasopressin, Are Associated with an Increased Risk of Chronic Kidney Disease in a General Population. <i>American Journal of Nephrology</i> , 2016, 44, 22-28.	1.4	53
48	Iohexol plasma clearance for measuring glomerular filtration rate in clinical practice and research: a review. Part 1: How to measure glomerular filtration rate with iohexol?. <i>CKJ: Clinical Kidney Journal</i> , 2016, 9, 682-699.	1.4	169
49	Iohexol plasma clearance for measuring glomerular filtration rate in clinical practice and research: a review. Part 2: Why to measure glomerular filtration rate with iohexol?. <i>CKJ: Clinical Kidney Journal</i> , 2016, 9, 700-704.	1.4	150
50	The shrunken pore syndrome is associated with declined right ventricular systolic function in a heart failure population—the HARVEST study. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2016, 76, 568-574.	0.6	34
51	Cystatin C and Risk of Diabetes and the Metabolic Syndrome—Biomarker and Genotype Association Analyses. <i>PLoS ONE</i> , 2016, 11, e0155735.	1.1	11
52	Declining Estimated Glomerular Filtration Rate and Its Association with Mortality and Comorbidity Over 10 Years in Elderly Women. <i>Nephron</i> , 2015, 130, 245-255.	0.9	45
53	Reduction in glomerular pore size is not restricted to pregnant women. Evidence for a new syndrome: the "Shrunken pore syndrome". <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2015, 75, 333-340.	0.6	85
54	Cystatin C Is Not Causally Related to Coronary Artery Disease. <i>PLoS ONE</i> , 2015, 10, e0129269.	1.1	26

#	ARTICLE	IF	CITATIONS
55	Male patients with terminal renal failure exhibit low serum levels of antimüllerian hormone. <i>Asian Journal of Andrology</i> , 2015, 17, 149.	0.8	26
56	Male sex and vascular risk factors affect cystatin C-derived renal function in older people without diabetes or overt vascular disease. <i>Age and Ageing</i> , 2014, 43, 411-417.	0.7	11
57	Family History of Myocardial Infarction Increases Risk of Renal Dysfunction in Middle Age. <i>American Journal of Nephrology</i> , 2014, 39, 85-91.	1.4	3
58	Generation of a New Cystatin C-Based Estimating Equation for Glomerular Filtration Rate by Use of 7 Assays Standardized to the International Calibrator. <i>Clinical Chemistry</i> , 2014, 60, 974-986.	1.5	248
59	Replacement of acetate with citrate in dialysis fluid: a randomized clinical trial of short term safety and fluid biocompatibility. <i>BMC Nephrology</i> , 2013, 14, 216.	0.8	58
60	Association of cancer with moderately impaired renal function at baseline in a large, representative, population-based cohort followed for up to 30 years. <i>International Journal of Cancer</i> , 2013, 133, 1452-1458.	2.3	64
61	Evaluation of a new immunoassay for cystatin C, based on a double monoclonal principle, in men with normal and impaired renal function. <i>Nephrology Dialysis Transplantation</i> , 2012, 27, 682-687.	0.4	6
62	Cystatin C as a predictor of all-cause mortality and myocardial infarction in patients with non-ST-elevation acute coronary syndrome. <i>Clinical Biochemistry</i> , 2012, 45, 535-540.	0.8	32
63	Intra-individual short-term variability of prostate-specific antigen and other kallikrein markers in a serial collection of blood from men under evaluation for prostate cancer. <i>BJU International</i> , 2011, 107, 1769-1774.	1.3	10
64	Achievement of recommended treatment targets for bone and mineral metabolism in haemodialysis patients using paricalcitol: An observational study. <i>Scandinavian Journal of Urology and Nephrology</i> , 2011, 45, 196-205.	1.4	4
65	Estimation of the Age-Dependent Decline of Glomerular Filtration Rate from Formulas Based on Creatinine and Cystatin C in the General Elderly Population. <i>Nephron Clinical Practice</i> , 2010, 117, c40-c50.	2.3	32
66	Neutrophil-Derived Proteinase 3 Induces Kallikrein-Independent Release of a Novel Vasoactive Kinin. <i>Journal of Immunology</i> , 2009, 182, 7906-7915.	0.4	50
67	Novel and Conventional Biomarkers for Prediction of Incident Cardiovascular Events in the Community. <i>JAMA - Journal of the American Medical Association</i> , 2009, 302, 49.	3.8	474
68	Increase in percent free prostate-specific antigen in men with chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2008, 24, 1238-1241.	0.4	25
69	Different elimination patterns of $\beta_2$ -microglobulin and cystatin C in haemodialysis, haemodiafiltration and haemofiltration. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2008, 68, 685-691.	0.6	29
70	A Study on the Outcome of Percutaneous Transluminal Renal Angioplasty in Patients with Renal Failure. <i>Nephron Clinical Practice</i> , 2006, 104, c132-c142.	2.3	5
71	Assessment of intra-individual variation in prostate-specific antigen levels in a biennial randomized prostate cancer screening program in Sweden. <i>Prostate</i> , 2005, 65, 216-221.	1.2	25
72	Simple Cystatin C-Based Prediction Equations for Glomerular Filtration Rate Compared with the Modification of Diet in Renal Disease Prediction Equation for Adults and the Schwartz and the Counahan-Barratt Prediction Equations for Children. <i>Clinical Chemistry</i> , 2005, 51, 1420-1431.	1.5	413

#	ARTICLE	IF	CITATIONS
73	RAPID ELIMINATION BY GLOMERULAR FILTRATION OF FREE PROSTATE SPECIFIC ANTIGEN AND HUMAN KALLIKREIN 2 AFTER RENAL TRANSPLANTATION. <i>Journal of Urology</i> , 2004, 171, 1432-1435.	0.2	28
74	Expression of protein C inhibitor (PCI) in benign and malignant prostatic tissues. <i>Prostate</i> , 2003, 57, 196-204.	1.2	34
75	A randomized controlled trial of haemoglobin normalization with epoetin alfa in pre-dialysis and dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2003, 18, 353-361.	0.4	195
76	Serum Cystatin C Is a More Sensitive and More Accurate Marker of Glomerular Filtration Rate than Enzymatic Measurements of Creatinine in Renal Transplantation. <i>Nephron Physiology</i> , 2003, 94, p19-p27.	1.5	67
77	Personal dialysis capacity (PDCTM) test: a multicentre clinical study. <i>Nephrology Dialysis Transplantation</i> , 2003, 18, 788-796.	0.4	34
78	Percent-free prostate specific antigen is elevated in men on haemodialysis or peritoneal dialysis treatment. <i>Nephrology Dialysis Transplantation</i> , 2003, 18, 598-603.	0.4	44
79	Cystine analyses of separate day and night urine as a basis for the management of patients with homozygous cystinuria. <i>Urological Research</i> , 2001, 29, 303-310.	1.5	34
80	Long-term clinical effects of a peritoneal dialysis fluid with less glucose degradation products. <i>Kidney International</i> , 2001, 59, 348-357.	2.6	239
81	Renovascular Disease and Renal Insufficiency - Diagnosis and Treatment. <i>Scandinavian Journal of Urology and Nephrology</i> , 1999, 33, 400-405.	1.4	6
82	Mobilization of a Bacterial Vegetation Visualized During Transesophageal Echocardiography. <i>Echocardiography</i> , 1998, 15, 381-383.	0.3	1
83	The Significance of Serpins in the Regulation of Proteases in the Male Genital Tract. <i>Advances in Experimental Medicine and Biology</i> , 1997, 425, 163-176.	0.8	2
84	Similar treatment success rate after renal transplantation in diabetic and nondiabetic patients due to improved short- and long-term diabetic patient survival. <i>Transplant International</i> , 1996, 9, 557-564.	0.8	20
85	Similar treatment success rate after renal transplantation in diabetic and nondiabetic patients due to improved short- and long-term diabetic patient survival. <i>Transplant International</i> , 1996, 9, 557-564.	0.8	8
86	Complex formation between protein C inhibitor and prostate-specific antigen in vitro and in human semen. <i>FEBS Journal</i> , 1994, 220, 45-53.	0.2	166
87	Serum Prostate Specific Antigen Complexed to $\alpha$ -1-Antichymotrypsin as an Indicator of Prostate Cancer. <i>Journal of Urology</i> , 1993, 150, 100-105.	0.2	629
88	Enzymatic activity of prostate-specific antigen and its reactions with extracellular serine proteinase inhibitors. <i>FEBS Journal</i> , 1990, 194, 755-763.	0.2	578
89	Clinical and Radiological Follow-Up of Chronic Non-Obstructive Pyelonephritis. <i>Scandinavian Journal of Urology and Nephrology</i> , 1988, 22, 299-303.	1.4	2