
Simona Dedoni

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5903596/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Upregulation of p75NTR by Histone Deacetylase Inhibitors Sensitizes Human Neuroblastoma Cells to Targeted Immunotoxin-Induced Apoptosis. International Journal of Molecular Sciences, 2022, 23, 3849.	1.8	1
2	K18- and CAG-hACE2 Transgenic Mouse Models and SARS-CoV-2: Implications for Neurodegeneration Research. Molecules, 2022, 27, 4142.	1.7	7
3	The Neurotrophin Receptor TrkC as a Novel Molecular Target of the Antineuroblastoma Action of Valproic Acid. International Journal of Molecular Sciences, 2021, 22, 7790.	1.8	4
4	Cannabinoid CB1 and CB2 receptors differentially regulate TNF-α-induced apoptosis and LPA1-mediated pro-survival signaling in HT22 hippocampal cells. Life Sciences, 2021, 276, 119407.	2.0	9
5	Valproic acid upregulates the expression of the p75NTR/sortilin receptor complex to induce neuronal apoptosis. Apoptosis: an International Journal on Programmed Cell Death, 2020, 25, 697-714.	2.2	11
6	Antidepressants induce profibrotic responses via the lysophosphatidic acid receptor LPA1. European Journal of Pharmacology, 2020, 873, 172963.	1.7	11
7	Inhibition of TNF-α-induced neuronal apoptosis by antidepressants acting through the lysophosphatidic acid receptor LPA1. Apoptosis: an International Journal on Programmed Cell Death, 2019, 24, 478-498.	2.2	15
8	Downregulation of TrkB Expression and Signaling by Valproic Acid and Other Histone Deacetylase Inhibitors. Journal of Pharmacology and Experimental Therapeutics, 2019, 370, 490-503.	1.3	19
9	Muscarinic Acetylcholine Receptors Potentiate 5′-Adenosine Monophosphate-Activated Protein Kinase Stimulation and Glucose Uptake Triggered by Thapsigargin-Induced Store-Operated Ca2+ Entry in Human Neuroblastoma Cells. Neurochemical Research, 2018, 43, 245-258.	1.6	4
10	The orphan Gâ€protein oupled receptor 75 signaling is activated by the chemokine <scp>CCL</scp> 5. Journal of Neurochemistry, 2018, 146, 526-539.	2.1	32
11	Activation of LPA1 signaling mediates antidepressant-induced neuroprotection from apoptosis triggered by pro-inflammatory cytokines. Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2018, WCP2018, PO2-1-55.	0.0	0
12	Valproic acid up-regulates p75NTR and sortilin expression to induce neuronal cell death. Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2018, WCP2018, PO2-1-77.	0.0	0
13	Interferon-β Inhibits Neurotrophin 3 Signalling and Pro-Survival Activity by Upregulating the Expression of Truncated TrkC-T1 Receptor. Molecular Neurobiology, 2017, 54, 1825-1843.	1.9	9
14	The GABAB positive allosteric modulators CGP7930 and GS39783 stimulate ERK1/2 signalling in cells lacking functional GABAB receptors. European Journal of Pharmacology, 2017, 794, 135-146.	1.7	8
15	<scp>LPA</scp> ₁ is a key mediator of intracellular signalling and neuroprotection triggered by tetracyclic antidepressants in hippocampal neurons. Journal of Neurochemistry, 2017, 143, 183-197.	2.1	14
16	Identification of a binding site of the human immunodeficiency virus envelope protein gp120 to neuronalâ€specific tubulin. Journal of Neurochemistry, 2016, 137, 287-298.	2.1	23
17	LPA ₁ Mediates Antidepressant-Induced ERK1/2 Signaling and Protection from Oxidative Stress in Glial Cells. Journal of Pharmacology and Experimental Therapeutics, 2016, 359, 340-353.	1.3	22
18	Protection from interferonâ€Î²â€induced neuronal apoptosis through stimulation of muscarinic acetylcholine receptors coupled to ERK1/2 activation. British Journal of Pharmacology, 2016, 173, 2910-2928.	2.7	10

SIMONA DEDONI

#	Article	IF	CITATIONS
19	The HIV Protein gp120 Alters Mitochondrial Dynamics in Neurons. Neurotoxicity Research, 2016, 29, 583-593.	1.3	77
20	Antidepressants activate the lysophosphatidic acid receptor LPA 1 to induce insulin-like growth factor-I receptor transactivation, stimulation of ERK1/2 signaling and cell proliferation in CHO-K1 fibroblasts. Biochemical Pharmacology, 2015, 95, 311-323.	2.0	26
21	Interferon-β counter-regulates its own pro-apoptotic action by activating p38 MAPK signalling in human SH-SY5Y neuroblastoma cells. Apoptosis: an International Journal on Programmed Cell Death, 2014, 19, 1509-1526.	2.2	10
22	Type I interferons up-regulate the expression and signalling of p75 NTR/TrkA receptor complex in differentiated human SH-SY5Y neuroblastoma cells. Neuropharmacology, 2014, 79, 321-334.	2.0	14
23	Involvement of store-operated Ca2+ entry in activation of AMP-activated protein kinase and stimulation of glucose uptake by M3 muscarinic acetylcholine receptors in human neuroblastoma cells. Biochimica Et Biophysica Acta - Molecular Cell Research, 2014, 1843, 3004-3017.	1.9	17
24	Preclinical and clinical phase I studies of a new recombinant Filgrastim (BK0023) in comparison with Neupogen®. BMC Pharmacology & Toxicology, 2014, 15, 7.	1.0	21
25	Coincidence Signaling of Dopamine D ₁ -Like and M ₁ Muscarinic Receptors in the Regulation of Cyclic AMP Formation and CREB Phosphorylation in Mouse Prefrontal Cortex. NeuroSignals, 2013, 21, 61-74.	0.5	6
26	δ-Opioid Receptors Stimulate the Metabolic Sensor AMP-Activated Protein Kinase through Coincident Signaling with G _{q/11} -Coupled Receptors. Molecular Pharmacology, 2012, 81, 154-165.	1.0	17
27	Potentiation of dopamine D1-like receptor signaling by concomitant activation of δ- and μ-opioid receptors in mouse medial prefrontal cortex. Neurochemistry International, 2012, 61, 1404-1416.	1.9	19
28	Enzymatic mono-pegylation of glucagon-like peptide 1 towards long lasting treatment of type 2 diabetes. Results in Pharma Sciences, 2012, 2, 58-65.	4.2	15
29	Type I interferons impair BDNFâ€induced cell signaling and neurotrophic activity in differentiated human SH‣Y5Y neuroblastoma cells and mouse primary cortical neurons. Journal of Neurochemistry, 2012, 122, 58-71.	2.1	30
30	The atypical antidepressant mianserin exhibits agonist activity at κâ€opioid receptors. British Journal of Pharmacology, 2012, 167, 1329-1341.	2.7	22
31	Signaling pathways mediating phosphorylation and inactivation of glycogen synthase kinase-3β by the recombinant human δ-opioid receptor stably expressed in Chinese hamster ovary cells. Neuropharmacology, 2011, 60, 1326-1336.	2.0	15
32	δâ€Opioid receptors stimulate GLUT1â€mediated glucose uptake through Src―and IGFâ€1 receptorâ€depender activation of PI3â€kinase signalling in CHO cells. British Journal of Pharmacology, 2011, 163, 624-637.	^{1t} 2.7	26
33	Regulation of PI3K/Akt signaling by N-desmethylclozapine through activation of δ-opioid receptor. European Journal of Pharmacology, 2011, 660, 341-350.	1.7	20
34	Interferonâ€Î² induces apoptosis in human SH‣Y5Y neuroblastoma cells through activation of JAK–STAT signaling and downâ€regulation of PI3K/Akt pathway. Journal of Neurochemistry, 2010, 115, 1421-1433.	2.1	63
35	Direct Agonist Activity of Tricyclic Antidepressants at Distinct Opioid Receptor Subtypes. Journal of Pharmacology and Experimental Therapeutics, 2010, 332, 255-265.	1.3	39
36	Agonist activity of N-desmethylclozapine at δ-opioid receptors of human frontal cortex. European Journal of Pharmacology, 2009, 607, 96-101.	1.7	27

#	Article	IF	CITATIONS
37	Activation of nociceptin/orphanin FQâ€NOP receptor system inhibits tyrosine hydroxylase phosphorylation, dopamine synthesis, and dopamine D ₁ receptor signaling in rat nucleus accumbens and dorsal striatum. Journal of Neurochemistry, 2008, 107, 544-556.	2.1	39
38	Proteinase-activated receptors 1 and 2 in rat olfactory system: Layer-specific regulation of multiple signaling pathways in the main olfactory bulb and induction of neurite retraction in olfactory sensory neurons. Neuroscience, 2007, 146, 1289-1301.	1.1	23