

Robert Feil

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5906050/robert-feil-publications-by-year.pdf>

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

142 papers	11,919 citations	65 h-index	108 g-index
153 ext. papers	13,324 ext. citations	8.7 avg, IF	5.81 L-index

#	Paper	IF	Citations
142	Apolipoprotein E derived from CD11c cells ameliorates atherosclerosis.. <i>IScience</i> , 2022 , 25, 103677	6.1	1
141	Olinciguat, a stimulator of soluble guanylyl cyclase, attenuates inflammation, vaso-occlusion and nephropathy in mouse models of sickle cell disease. <i>British Journal of Pharmacology</i> , 2021 , 178, 3463-3475	8.6	3
140	Visualising and understanding cGMP signals in the cardiovascular system. <i>British Journal of Pharmacology</i> , 2021 ,	8.6	2
139	The C5a/C5a receptor 1 axis controls tissue neovascularization through CXCL4 release from platelets. <i>Nature Communications</i> , 2021 , 12, 3352	17.4	4
138	Analysis of Gene Expression Using lacZ Reporter Mouse Lines. <i>Methods in Molecular Biology</i> , 2021 , 2224, 29-45	1.4	
137	Novel soluble guanylyl cyclase activators increase glomerular cGMP, induce vasodilation and improve blood flow in the murine kidney. <i>British Journal of Pharmacology</i> , 2021 ,	8.6	3
136	Amelanotic B16-F10 Melanoma Compatible with Advanced Three-Dimensional Imaging Modalities. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 2090-2094.e6	4.3	1
135	Guanylyl Cyclase A/cGMP Signaling Slows Hidden, Age- and Acoustic Trauma-Induced Hearing Loss. <i>Frontiers in Aging Neuroscience</i> , 2020 , 12, 83	5.3	5
134	cGMP-dependent protein kinase I in vascular smooth muscle cells improves ischemic stroke outcome in mice. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2019 , 39, 2379-2391	7.3	4
133	Npom-Protected NONOate Enables Light-Triggered NO/cGMP Signalling in Primary Vascular Smooth Muscle Cells. <i>ChemBioChem</i> , 2018 , 19, 1312-1318	3.8	6
132	Real-Time Imaging Reveals Augmentation of Glutamate-Induced Ca Transients by the NO-cGMP Pathway in Cerebellar Granule Neurons. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	3
131	The Absence of Sensory Axon Bifurcation Affects Nociception and Termination Fields of Afferents in the Spinal Cord. <i>Frontiers in Molecular Neuroscience</i> , 2018 , 11, 19	6.1	11
130	cGMP Signaling and Vascular Smooth Muscle Cell Plasticity. <i>Journal of Cardiovascular Development and Disease</i> , 2018 , 5,	4.2	21
129	cGMP Imaging in Brain Slices Reveals Brain Region-Specific Activity of NO-Sensitive Guanylyl Cyclases (NO-GCs) and NO-GC Stimulators. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	5
128	Chronic linacotide treatment reduces colitis-induced neuroplasticity and reverses persistent bladder dysfunction. <i>JCI Insight</i> , 2018 , 3,	9.9	38
127	Regulation of the Natriuretic Peptide Receptor 2 (Npr2) by Phosphorylation of Juxtamembrane Serine and Threonine Residues Is Essential for Bifurcation of Sensory Axons. <i>Journal of Neuroscience</i> , 2018 , 38, 9768-9780	6.6	7
126	GC-B Deficient Mice With Axon Bifurcation Loss Exhibit Compromised Auditory Processing. <i>Frontiers in Neural Circuits</i> , 2018 , 12, 65	3.5	11

125	A shear-dependent NO-cGMP-cGKI cascade in platelets acts as an auto-regulatory brake of thrombosis. <i>Nature Communications</i> , 2018 , 9, 4301	17.4	16
124	cGMP-dependent protein kinase I (cGKI) modulates human hepatic stellate cell activation. <i>Metabolism: Clinical and Experimental</i> , 2018 , 88, 22-30	12.7	14
123	Viagra releases the brakes on melanoma growth. <i>Molecular and Cellular Oncology</i> , 2017 , 4, e1188874	1.2	3
122	Heterotrimeric G Stimulatory Protein β Subunit Is Required for Intestinal Smooth Muscle Contraction in Mice. <i>Gastroenterology</i> , 2017 , 152, 1114-1125.e5	13.3	10
121	Amplified pathogenic actions of angiotensin II in cysteine-rich LIM-only protein 4-negative mouse hearts. <i>FASEB Journal</i> , 2017 , 31, 1620-1638	0.9	5
120	Microglia turnover with aging and in an Alzheimer's model via long-term in vivo single-cell imaging. <i>Nature Neuroscience</i> , 2017 , 20, 1371-1376	25.5	193
119	NO-Sensitive Guanylate Cyclase Isoforms NO-GC1 and NO-GC2 Contribute to Noise-Induced Inner Hair Cell Synaptopathy. <i>Molecular Pharmacology</i> , 2017 , 92, 375-388	4.3	18
118	Oxidant sensor in the cGMP-binding pocket of PKGI regulates nitroxyl-mediated kinase activity. <i>Scientific Reports</i> , 2017 , 7, 9938	4.9	15
117	Cre/lox-assisted non-invasive in vivo tracking of specific cell populations by positron emission tomography. <i>Nature Communications</i> , 2017 , 8, 444	17.4	28
116	Dephosphorylation of the NPR2 guanylyl cyclase contributes to inhibition of bone growth by fibroblast growth factor. <i>ELife</i> , 2017 , 6,	8.9	19
115	cGMP Signaling in Platelets. <i>Cardiac and Vascular Biology</i> , 2017 , 231-252	0.2	2
114	Dorsal root ganglion axon bifurcation tolerates increased cyclic GMP levels: the role of phosphodiesterase 2A and scavenger receptor Npr3. <i>European Journal of Neuroscience</i> , 2016 , 44, 2991-3000	3.5	15
113	Sildenafil Potentiates a cGMP-Dependent Pathway to Promote Melanoma Growth. <i>Cell Reports</i> , 2016 , 14, 2599-610	10.6	49
112	Intercellular signaling via cyclic GMP diffusion through gap junctions restarts meiosis in mouse ovarian follicles. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 5527-32	11.5	99
111	Platelet-derived HMGB1 is a critical mediator of thrombosis. <i>Journal of Clinical Investigation</i> , 2015 , 125, 4638-54	15.9	190
110	Comparative analysis of established and new biosensors for cyclic nucleotides. <i>BMC Pharmacology & Toxicology</i> , 2015 , 16,	2.6	78
109	Real-time imaging of cGMP signals in platelets. <i>BMC Pharmacology & Toxicology</i> , 2015 , 16,	2.6	78
108	The small GTPase Rac1 is required for smooth muscle contraction. <i>Journal of Physiology</i> , 2014 , 592, 915-926	3.6	26

107	Upon the photostability of 8-nitro-cGMP and its caging as a 7-dimethylaminocoumarinyl ester. <i>Chemical Communications</i> , 2014 , 50, 7120-3	5.8	4
106	Transdifferentiation of vascular smooth muscle cells to macrophage-like cells during atherogenesis. <i>Circulation Research</i> , 2014 , 115, 662-7	15.7	283
105	Autonomous functions of murine thyroid hormone receptor TR α and TR β in cochlear hair cells. <i>Molecular and Cellular Endocrinology</i> , 2014 , 382, 26-37	4.4	20
104	Catalytic activity of cGMP-dependent protein kinase type I in intact cells is independent of N-terminal autophosphorylation. <i>PLoS ONE</i> , 2014 , 9, e98946	3.7	4
103	Correlative intravital imaging of cGMP signals and vasodilation in mice. <i>Frontiers in Physiology</i> , 2014 , 5, 394	4.6	18
102	Upregulation of non- β cell-derived vascular endothelial growth factor A increases small clusters of insulin-producing cells in the pancreas. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014 , 122, 308-15	2.3	3
101	Endless: a purine-binding RNA motif that can be expressed in cells. <i>Angewandte Chemie - International Edition</i> , 2014 , 53, 9198-202	16.4	10
100	Endless: A Purine-Binding RNA Motif that Can Be Expressed in Cells. <i>Angewandte Chemie</i> , 2014 , 126, 9352-9356	3.6	4
99	Alterations in the cerebellar (Phospho)proteome of a cyclic guanosine monophosphate (cGMP)-dependent protein kinase knockout mouse. <i>Molecular and Cellular Proteomics</i> , 2014 , 13, 2004-16	7.6	11
98	TGF- β signaling mediates endothelial-to-mesenchymal transition (EndMT) during vein graft remodeling. <i>Science Translational Medicine</i> , 2014 , 6, 227ra34	17.5	241
97	Genetic inducible fate mapping in adult mice using tamoxifen-dependent Cre recombinases. <i>Methods in Molecular Biology</i> , 2014 , 1194, 113-39	1.4	11
96	Defining the molecular targets of cerebellar PKG by quantitative (phospho)proteomics in a knock-out mouse model. <i>BMC Pharmacology & Toxicology</i> , 2013 , 14,	2.6	78
95	Interstitial cells of Cajal integrate excitatory and inhibitory neurotransmission with intestinal slow-wave activity. <i>Nature Communications</i> , 2013 , 4, 1630	17.4	144
94	Visualization of cGMP with cGi biosensors. <i>Methods in Molecular Biology</i> , 2013 , 1020, 89-120	1.4	26
93	Stress-dependent dilated cardiomyopathy in mice with cardiomyocyte-restricted inactivation of cyclic GMP-dependent protein kinase I. <i>European Heart Journal</i> , 2013 , 34, 1233-44	9.5	70
92	Selective involvement of serum response factor in pressure-induced myogenic tone in resistance arteries. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013 , 33, 339-46	9.4	11
91	Inactivation of serum response factor contributes to decrease vascular muscular tone and arterial stiffness in mice. <i>Circulation Research</i> , 2013 , 112, 1035-45	15.7	33
90	α 1-integrin is essential for vasoregulation and smooth muscle survival in vivo. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013 , 33, 2325-35	9.4	15

89	Atrial natriuretic peptide-mediated inhibition of microcirculatory endothelial Ca ²⁺ and permeability response to histamine involves cGMP-dependent protein kinase I and TRPC6 channels. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013 , 33, 2121-9	9.4	34
88	Transgenic mice for cGMP imaging. <i>Circulation Research</i> , 2013 , 113, 365-71	15.7	55
87	Partial loss of contractile marker proteins in human testicular peritubular cells in infertility patients. <i>Andrology</i> , 2013 , 1, 318-24	4.2	32
86	H ₂ O ₂ lowers the cytosolic Ca ²⁺ concentration via activation of cGMP-dependent protein kinase II. <i>Free Radical Biology and Medicine</i> , 2012 , 53, 1574-83	7.8	25
85	cGMP-Prkg1 signaling and Pde5 inhibition shelter cochlear hair cells and hearing function. <i>Nature Medicine</i> , 2012 , 18, 252-9	50.5	67
84	Restoring nitric oxide cytosolic calcium regulation by cyclic guanosine monophosphate protein kinase I alpha transfection in coronary endothelial cells of spontaneously hypertensive rats. <i>Journal of Vascular Research</i> , 2012 , 49, 221-30	1.9	4
83	Presynaptically localized cyclic GMP-dependent protein kinase 1 is a key determinant of spinal synaptic potentiation and pain hypersensitivity. <i>PLoS Biology</i> , 2012 , 10, e1001283	9.7	71
82	cGMP-dependent protein kinase contributes to hydrogen sulfide-stimulated vasorelaxation. <i>PLoS ONE</i> , 2012 , 7, e53319	3.7	97
81	Fumarates improve psoriasis and multiple sclerosis by inducing type II dendritic cells. <i>Journal of Experimental Medicine</i> , 2011 , 208, 2291-303	16.6	284
80	Role of smooth muscle protein SM22 α in glomerular epithelial cell injury. <i>American Journal of Physiology - Renal Physiology</i> , 2011 , 300, F1026-42	4.3	26
79	A cardiac pathway of cyclic GMP-independent signaling of guanylyl cyclase A, the receptor for atrial natriuretic peptide. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 18500-5	11.5	40
78	HCN3 contributes to the ventricular action potential waveform in the murine heart. <i>Circulation Research</i> , 2011 , 109, 1015-23	15.7	52
77	Genetic ablation of cGMP-dependent protein kinase type I causes liver inflammation and fasting hyperglycemia. <i>Diabetes</i> , 2011 , 60, 1566-76	0.9	33
76	IRAG determines nitric oxide- and atrial natriuretic peptide-mediated smooth muscle relaxation. <i>Cardiovascular Research</i> , 2010 , 86, 496-505	9.9	43
75	Novel insights into the mechanisms mediating the local antihypertrophic effects of cardiac atrial natriuretic peptide: role of cGMP-dependent protein kinase and RGS2. <i>Basic Research in Cardiology</i> , 2010 , 105, 583-95	11.8	60
74	cGMP-dependent protein kinase type I is implicated in the regulation of the timing and quality of sleep and wakefulness. <i>PLoS ONE</i> , 2009 , 4, e4238	3.7	45
73	Regulate axon branching by the cyclic GMP pathway via inhibition of glycogen synthase kinase 3 in dorsal root ganglion sensory neurons. <i>Journal of Neuroscience</i> , 2009 , 29, 1350-60	6.6	61
72	The tumor suppressor p53 transcriptionally regulates cGKI expression during neuronal maturation and is required for cGMP-dependent growth cone collapse. <i>Journal of Neuroscience</i> , 2009 , 29, 15155-60	6.6	29

71	The commonly used cGMP-dependent protein kinase type I (cGKI) inhibitor Rp-8-Br-PET-cGMPS can activate cGKI in vitro and in intact cells. <i>Journal of Biological Chemistry</i> , 2009 , 284, 556-562	5.4	35
70	cGMP-dependent protein kinase I, the circadian clock, sleep and learning. <i>Communicative and Integrative Biology</i> , 2009 , 2, 298-301	1.7	18
69	C-type natriuretic peptide is a bifurcation factor for sensory neurons. <i>BMC Pharmacology</i> , 2009 , 9,		78
68	cGMP signalling in the mammalian brain: role in synaptic plasticity and behaviour. <i>Handbook of Experimental Pharmacology</i> , 2009 , 549-79	3.2	122
67	Inducible Cre mice. <i>Methods in Molecular Biology</i> , 2009 , 530, 343-63	1.4	162
66	Altered nitric oxide calcium responsiveness of aortic smooth muscle cells in spontaneously hypertensive rats depends on low expression of cyclic guanosine monophosphate-dependent protein kinase type I. <i>Journal of Hypertension</i> , 2009 , 27, 1258-67	1.9	7
65	cGMP-dependent protein kinase I is crucial for angiogenesis and postnatal vasculogenesis. <i>PLoS ONE</i> , 2009 , 4, e4879	3.7	19
64	LKB1 signaling in mesenchymal cells required for suppression of gastrointestinal polyposis. <i>Nature Genetics</i> , 2008 , 40, 455-9	36.3	88
63	Myosin light chain kinase is central to smooth muscle contraction and required for gastrointestinal motility in mice. <i>Gastroenterology</i> , 2008 , 135, 610-20	13.3	132
62	Tamoxifen-inducible gene deletion in the cardiac conduction system. <i>Journal of Molecular and Cellular Cardiology</i> , 2008 , 45, 62-9	5.8	84
61	NO/cGMP-dependent modulation of synaptic transmission. <i>Handbook of Experimental Pharmacology</i> , 2008 , 529-60	3.2	131
60	Cyclic guanosine monophosphate-dependent protein kinase I promotes adhesion of primary vascular smooth muscle cells. <i>Molecular Biology of the Cell</i> , 2008 , 19, 4434-41	3.5	17
59	Anemia and splenomegaly in cGKI-deficient mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 6771-6	11.5	125
58	Role of smooth muscle cGMP/cGKI signaling in murine vascular restenosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2008 , 28, 1244-50	9.4	29
57	Severe intestinal obstruction on induced smooth muscle-specific ablation of the transcription factor SRF in adult mice. <i>Gastroenterology</i> , 2007 , 133, 1948-59	13.3	33
56	Inducible mouse model of chronic intestinal pseudo-obstruction by smooth muscle-specific inactivation of the SRF gene. <i>Gastroenterology</i> , 2007 , 133, 1960-70	13.3	37
55	Rescue of cGMP kinase I knockout mice by smooth muscle specific expression of either isozyme. <i>Circulation Research</i> , 2007 , 101, 1096-103	15.7	93
54	The receptor guanylyl cyclase Npr2 is essential for sensory axon bifurcation within the spinal cord. <i>Journal of Cell Biology</i> , 2007 , 179, 331-40	7.3	73

53	Ablation of connexin43 in smooth muscle cells of the mouse intestine: functional insights into physiology and morphology. <i>Cell and Tissue Research</i> , 2007 , 327, 333-42	4.2	15
52	Conditional somatic mutagenesis in the mouse using site-specific recombinases. <i>Handbook of Experimental Pharmacology</i> , 2007 , 3-28	3.2	55
51	Function of cGMP-dependent protein kinases as revealed by gene deletion. <i>Physiological Reviews</i> , 2006 , 86, 1-23	47.9	327
50	Mdm2, but not Mdm4, protects terminally differentiated smooth muscle cells from p53-mediated caspase-3-independent cell death. <i>Cell Death and Differentiation</i> , 2006 , 13, 2089-98	12.7	51
49	Control of intestinal motility by the Ca(v)1.2 L-type calcium channel in mice. <i>FASEB Journal</i> , 2006 , 20, 1260-2	0.9	41
48	cGMP-dependent protein kinase type I inhibits TAB1-p38 mitogen-activated protein kinase apoptosis signaling in cardiac myocytes. <i>Journal of Biological Chemistry</i> , 2006 , 281, 32831-40	5.4	69
47	Ablation of connexin43 in uterine smooth muscle cells of the mouse causes delayed parturition. <i>Journal of Cell Science</i> , 2006 , 119, 1715-22	5.3	87
46	The cGMP/protein kinase G pathway contributes to dihydropyridine-sensitive calcium response and cytokine production in TH2 lymphocytes. <i>Journal of Biological Chemistry</i> , 2006 , 281, 12421-7	5.4	22
45	A sensitive method for determining the phosphorylation status of natriuretic peptide receptors: cGK-Ialpha does not regulate NPR-A. <i>Biochemistry</i> , 2006 , 45, 1295-303	3.2	24
44	Sequential activation of p38 and ERK pathways by cGMP-dependent protein kinase leading to activation of the platelet integrin alphaIIb beta3. <i>Blood</i> , 2006 , 107, 965-72	2.2	128
43	A heretical view on the role of NO and cGMP in vascular proliferative diseases. <i>Trends in Molecular Medicine</i> , 2005 , 11, 71-5	11.5	41
42	Distribution of cGMP-dependent protein kinase type I and its isoforms in the mouse brain and retina. <i>Neuroscience</i> , 2005 , 135, 863-8	3.9	58
41	Insights into cGMP signalling derived from cGMP kinase knockout mice. <i>Frontiers in Bioscience - Landmark</i> , 2005 , 10, 1279-89	2.8	54
40	Elevated blood pressure linked to primary hyperaldosteronism and impaired vasodilation in BK channel-deficient mice. <i>Circulation</i> , 2005 , 112, 60-8	16.7	195
39	Function of cGMP-dependent protein kinases in the nervous system. <i>Reviews in the Neurosciences</i> , 2005 , 16, 23-41	4.7	68
38	cGMP-dependent protein kinase mediates NO- but not acetylcholine-induced dilations in resistance vessels in vivo. <i>Hypertension</i> , 2004 , 44, 952-5	8.5	66
37	Cerebellar ataxia and Purkinje cell dysfunction caused by Ca2+-activated K+ channel deficiency. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 9474-8	11.5	331
36	An essential role of Cav1.2 L-type calcium channel for urinary bladder function. <i>FASEB Journal</i> , 2004 , 18, 1159-61	0.9	93

35	Reduced inflammatory hyperalgesia with preservation of acute thermal nociception in mice lacking cGMP-dependent protein kinase I. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 3253-7	11.5	88
34	IRAG is essential for relaxation of receptor-triggered smooth muscle contraction by cGMP kinase. <i>EMBO Journal</i> , 2004 , 23, 4222-31	13	99
33	SM22alpha modulates vascular smooth muscle cell phenotype during atherogenesis. <i>Circulation Research</i> , 2004 , 94, 863-5	15.7	113
32	GPIIb-dependent platelet activation is dependent on Src kinases but not MAP kinase or cGMP-dependent kinase. <i>Blood</i> , 2004 , 103, 2601-9	2.2	79
31	Significance and therapeutic potential of the natriuretic peptides/cGMP/cGMP-dependent protein kinase pathway in vascular regeneration. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 3404-9	11.5	139
30	The hyperpolarization-activated channel HCN4 is required for the generation of pacemaker action potentials in the embryonic heart. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 15235-40	11.5	357
29	Hippocampal cGMP-dependent protein kinase I supports an age- and protein synthesis-dependent component of long-term potentiation but is not essential for spatial reference and contextual memory. <i>Journal of Neuroscience</i> , 2003 , 23, 6005-12	6.6	80
28	Individual cerebellar Purkinje cells express different cGMP phosphodiesterases (PDEs): in vivo phosphorylation of cGMP-specific PDE (PDE5) as an indicator of cGMP-dependent protein kinase (PKG) activation. <i>Journal of Neuroscience</i> , 2003 , 23, 6452-9	6.6	82
27	Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. <i>EMBO Journal</i> , 2003 , 22, 216-24	13	389
26	Impaired insulin secretion and glucose tolerance in beta cell-selective Ca(v)1.2 Ca ²⁺ channel null mice. <i>EMBO Journal</i> , 2003 , 22, 3844-54	13	174
25	Dominant role of smooth muscle L-type calcium channel Cav1.2 for blood pressure regulation. <i>EMBO Journal</i> , 2003 , 22, 6027-34	13	226
24	cGMP-dependent protein kinase II modulates mPer1 and mPer2 gene induction and influences phase shifts of the circadian clock. <i>Current Biology</i> , 2003 , 13, 725-33	6.3	67
23	A stimulatory role for cGMP-dependent protein kinase in platelet activation. <i>Cell</i> , 2003 , 112, 77-86	56.2	221
22	Physiology and pathophysiology of vascular signaling controlled by guanosine 3',5'-cyclic monophosphate-dependent protein kinase [corrected]. <i>Circulation</i> , 2003 , 108, 2172-83	16.7	267
21	Signaling through NO and cGMP-dependent protein kinases. <i>Annals of Medicine</i> , 2003 , 35, 21-7	1.5	144
20	A proatherogenic role for cGMP-dependent protein kinase in vascular smooth muscle cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 13519-24	11.5	67
19	Impairment of LTD and cerebellar learning by Purkinje cell-specific ablation of cGMP-dependent protein kinase I. <i>Journal of Cell Biology</i> , 2003 , 163, 295-302	7.3	122
18	Alternative splicing of cGMP-dependent protein kinase I and nitrate tolerance. <i>Circulation Research</i> , 2003 , 93, e143	15.7	0

17	Cyclic GMP-dependent protein kinases and the cardiovascular system: insights from genetically modified mice. <i>Circulation Research</i> , 2003 , 93, 907-16	15.7	239
16	Direct activation of PDE5 by cGMP: long-term effects within NO/cGMP signaling. <i>Journal of Cell Biology</i> , 2003 , 160, 719-27	7.3	139
15	Effects of cGMP-Dependent Protein Kinase Knockouts 2003 , 511-514		
14	p8-deficient fibroblasts grow more rapidly and are more resistant to adriamycin-induced apoptosis. <i>Oncogene</i> , 2002 , 21, 1685-94	9.2	75
13	Functional reconstitution of vascular smooth muscle cells with cGMP-dependent protein kinase I isoforms. <i>Circulation Research</i> , 2002 , 90, 1080-6	15.7	96
12	cGMP-mediated signaling via cGKIalpha is required for the guidance and connectivity of sensory axons. <i>Journal of Cell Biology</i> , 2002 , 159, 489-98	7.3	93
11	Regulation of cGMP-specific phosphodiesterase (PDE5) phosphorylation in smooth muscle cells. <i>Journal of Biological Chemistry</i> , 2002 , 277, 3310-7	5.4	167
10	cGMP-dependent protein kinase I mediates the negative inotropic effect of cGMP in the murine myocardium. <i>Circulation Research</i> , 2002 , 90, 18-20	15.7	148
9	Temporally controlled somatic mutagenesis in smooth muscle. <i>Genesis</i> , 2000 , 28, 15-22	1.9	132
8	Functional embryonic cardiomyocytes after disruption of the L-type alpha1C (Cav1.2) calcium channel gene in the mouse. <i>Journal of Biological Chemistry</i> , 2000 , 275, 39193-9	5.4	207
7	Engineering the mouse genome by site-specific recombination. <i>Current Opinion in Biotechnology</i> , 1999 , 10, 470-6	11.4	103
6	A chimeric Cre recombinase inducible by synthetic, but not by natural ligands of the glucocorticoid receptor. <i>Nucleic Acids Research</i> , 1998 , 26, 4086-90	20.1	78
5	Regulation of Cre recombinase activity by mutated estrogen receptor ligand-binding domains. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 237, 752-7	3.4	757
4	Ligand-activated site-specific recombination in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 10887-90	11.5	707
3	Functional cGMP-dependent protein kinase is phosphorylated in its catalytic domain at threonine-516. <i>Biochemistry</i> , 1995 , 34, 13152-8	3.2	29
2	High-level expression of functional cGMP-dependent protein kinase using the baculovirus system. <i>FEBS Letters</i> , 1993 , 336, 163-7	3.8	17
1	Purification and characterization of a novel thermostable 4-alpha-glucanotransferase of <i>Thermotoga maritima</i> cloned in <i>Escherichia coli</i> . <i>FEBS Journal</i> , 1992 , 207, 81-8		85