

# William G Wierda

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/5916319/william-g-wierda-publications-by-citations.pdf>

**Version:** 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

172  
papers

19,339  
citations

69  
h-index

138  
g-index

180  
ext. papers

22,360  
ext. citations

7  
avg, IF

6.25  
L-index

#	Paper	IF	Citations
172	Targeting BTK with ibrutinib in relapsed chronic lymphocytic leukemia. <i>New England Journal of Medicine</i> , <b>2013</b> , 369, 32-42	59.2	1656
171	Targeting BCL2 with Venetoclax in Relapsed Chronic Lymphocytic Leukemia. <i>New England Journal of Medicine</i> , <b>2016</b> , 374, 311-22	59.2	1164
170	Early results of a chemoimmunotherapy regimen of fludarabine, cyclophosphamide, and rituximab as initial therapy for chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , <b>2005</b> , 23, 4079-88	2.2	811
169	Use of CAR-Transduced Natural Killer Cells in CD19-Positive Lymphoid Tumors. <i>New England Journal of Medicine</i> , <b>2020</b> , 382, 545-553	59.2	652
168	Substantial susceptibility of chronic lymphocytic leukemia to BCL2 inhibition: results of a phase I study of navitoclax in patients with relapsed or refractory disease. <i>Journal of Clinical Oncology</i> , <b>2012</b> , 30, 488-96	2.2	622
167	Acalabrutinib (ACP-196) in Relapsed Chronic Lymphocytic Leukemia. <i>New England Journal of Medicine</i> , <b>2016</b> , 374, 323-32	59.2	621
166	Long-term results of the fludarabine, cyclophosphamide, and rituximab regimen as initial therapy of chronic lymphocytic leukemia. <i>Blood</i> , <b>2008</b> , 112, 975-80	2.2	568
165	Venetoclax in relapsed or refractory chronic lymphocytic leukaemia with 17p deletion: a multicentre, open-label, phase 2 study. <i>Lancet Oncology, The</i> , <b>2016</b> , 17, 768-778	21.7	536
164	Three-year follow-up of treatment-naïve and previously treated patients with CLL and SLL receiving single-agent ibrutinib. <i>Blood</i> , <b>2015</b> , 125, 2497-506	2.2	529
163	Ofatumumab as single-agent CD20 immunotherapy in fludarabine-refractory chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , <b>2010</b> , 28, 1749-55	2.2	483
162	Chemoimmunotherapy with fludarabine, cyclophosphamide, and rituximab for relapsed and refractory chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , <b>2005</b> , 23, 4070-8	2.2	447
161	Phase I First-in-Human Study of Venetoclax in Patients With Relapsed or Refractory Non-Hodgkin Lymphoma. <i>Journal of Clinical Oncology</i> , <b>2017</b> , 35, 826-833	2.2	442
160	Ibrutinib as initial therapy for elderly patients with chronic lymphocytic leukaemia or small lymphocytic lymphoma: an open-label, multicentre, phase 1b/2 trial. <i>Lancet Oncology, The</i> , <b>2014</b> , 15, 48-58	21.7	372
159	Lenalidomide induces complete and partial remissions in patients with relapsed and refractory chronic lymphocytic leukemia. <i>Blood</i> , <b>2008</b> , 111, 5291-7	2.2	355
158	Fludarabine, cyclophosphamide, and rituximab treatment achieves long-term disease-free survival in IGHV-mutated chronic lymphocytic leukemia. <i>Blood</i> , <b>2016</b> , 127, 303-9	2.2	347
157	CD40-ligand (CD154) gene therapy for chronic lymphocytic leukemia. <i>Blood</i> , <b>2000</b> , 96, 2917-2924	2.2	294
156	Safety and activity of ibrutinib plus rituximab for patients with high-risk chronic lymphocytic leukaemia: a single-arm, phase 2 study. <i>Lancet Oncology, The</i> , <b>2014</b> , 15, 1090-9	21.7	283

155	Single-agent ibrutinib in treatment-naïve and relapsed/refractory chronic lymphocytic leukemia: a 5-year experience. <i>Blood</i> , <b>2018</b> , 131, 1910-1919	2.2	267
154	Prognostic nomogram and index for overall survival in previously untreated patients with chronic lymphocytic leukemia. <i>Blood</i> , <b>2007</b> , 109, 4679-85	2.2	264
153	Diverse marrow stromal cells protect CLL cells from spontaneous and drug-induced apoptosis: development of a reliable and reproducible system to assess stromal cell adhesion-mediated drug resistance. <i>Blood</i> , <b>2009</b> , 114, 4441-50	2.2	260
152	Ibrutinib and Venetoclax for First-Line Treatment of CLL. <i>New England Journal of Medicine</i> , <b>2019</b> , 380, 2095-2103	59.2	256
151	Outcomes of patients with chronic lymphocytic leukemia after discontinuing ibrutinib. <i>Blood</i> , <b>2015</b> , 125, 2062-7	2.2	255
150	Relative value of ZAP-70, CD38, and immunoglobulin mutation status in predicting aggressive disease in chronic lymphocytic leukemia. <i>Blood</i> , <b>2008</b> , 112, 1923-30	2.2	254
149	High-level expression of the T-cell chemokines CCL3 and CCL4 by chronic lymphocytic leukemia B cells in nurse-like cell cocultures and after BCR stimulation. <i>Blood</i> , <b>2009</b> , 113, 3050-8	2.2	238
148	Venetoclax for chronic lymphocytic leukaemia progressing after ibrutinib: an interim analysis of a multicentre, open-label, phase 2 trial. <i>Lancet Oncology</i> , <b>2018</b> , 19, 65-75	21.7	228
147	Clonal evolution in patients with chronic lymphocytic leukaemia developing resistance to BTK inhibition. <i>Nature Communications</i> , <b>2016</b> , 7, 11589	17.4	220
146	Experience with alemtuzumab plus rituximab in patients with relapsed and refractory lymphoid malignancies. <i>Blood</i> , <b>2003</b> , 101, 3413-5	2.2	220
145	Chronic lymphocytic leukaemia. <i>Nature Reviews Disease Primers</i> , <b>2017</b> , 3, 16096	51.1	219
144	Acalabrutinib with or without obinutuzumab versus chlorambucil and obinutuzumab for treatment-naïve chronic lymphocytic leukaemia (ELEVATE TN): a randomised, controlled, phase 3 trial. <i>Lancet</i> , <b>2020</b> , 395, 1278-1291	40	201
143	Alemtuzumab as treatment for residual disease after chemotherapy in patients with chronic lymphocytic leukemia. <i>Cancer</i> , <b>2003</b> , 98, 2657-63	6.4	186
142	Complex karyotype is a stronger predictor than del(17p) for an inferior outcome in relapsed or refractory chronic lymphocytic leukemia patients treated with ibrutinib-based regimens. <i>Cancer</i> , <b>2015</b> , 121, 3612-21	6.4	185
141	Venetoclax for Patients With Chronic Lymphocytic Leukemia With 17p Deletion: Results From the Full Population of a Phase II Pivotal Trial. <i>Journal of Clinical Oncology</i> , <b>2018</b> , 36, 1973-1980	2.2	174
140	Overexpression of the CXCR5 chemokine receptor, and its ligand, CXCL13 in B-cell chronic lymphocytic leukemia. <i>Blood</i> , <b>2007</b> , 110, 3316-25	2.2	170
139	Prognostic value of miR-155 in individuals with monoclonal B-cell lymphocytosis and patients with B chronic lymphocytic leukemia. <i>Blood</i> , <b>2013</b> , 122, 1891-9	2.2	157
138	Fludarabine, cyclophosphamide, and rituximab chemoimmunotherapy is highly effective treatment for relapsed patients with CLL. <i>Blood</i> , <b>2011</b> , 117, 3016-24	2.2	152

137	Pharmacological and Protein Profiling Suggests Venetoclax (ABT-199) as Optimal Partner with Ibrutinib in Chronic Lymphocytic Leukemia. <i>Clinical Cancer Research</i> , <b>2015</b> , 21, 3705-15	12.9	147
136	Lenalidomide as initial therapy of elderly patients with chronic lymphocytic leukemia. <i>Blood</i> , <b>2011</b> , 118, 3489-98	2.2	145
135	Ofatumumab is active in patients with fludarabine-refractory CLL irrespective of prior rituximab: results from the phase 2 international study. <i>Blood</i> , <b>2011</b> , 118, 5126-9	2.2	139
134	De novo deletion 17p13.1 chronic lymphocytic leukemia shows significant clinical heterogeneity: the M. D. Anderson and Mayo Clinic experience. <i>Blood</i> , <b>2009</b> , 114, 957-64	2.2	134
133	Mechanism of action of SNS-032, a novel cyclin-dependent kinase inhibitor, in chronic lymphocytic leukemia. <i>Blood</i> , <b>2009</b> , 113, 4637-45	2.2	132
132	Phase II study of lenalidomide and rituximab as salvage therapy for patients with relapsed or refractory chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , <b>2013</b> , 31, 584-91	2.2	127
131	Isoform-selective phosphoinositide 3 kinase inhibitors inhibit CXCR4 signaling and overcome stromal cell-mediated drug resistance in chronic lymphocytic leukemia: a novel therapeutic approach. <i>Blood</i> , <b>2009</b> , 113, 5549-57	2.2	125
130	Phase I-II study of oxaliplatin, fludarabine, cytarabine, and rituximab combination therapy in patients with Richter's syndrome or fludarabine-refractory chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , <b>2008</b> , 26, 196-203	2.2	124
129	High expression of activation-induced cytidine deaminase (AID) and splice variants is a distinctive feature of poor-prognosis chronic lymphocytic leukemia. <i>Blood</i> , <b>2003</b> , 101, 4903-8	2.2	121
128	Randomized trial of ibrutinib vs ibrutinib plus rituximab in patients with chronic lymphocytic leukemia. <i>Blood</i> , <b>2019</b> , 133, 1011-1019	2.2	120
127	Eradication of bone marrow minimal residual disease may prompt early treatment discontinuation in CLL. <i>Blood</i> , <b>2014</b> , 123, 3727-32	2.2	109
126	Mitochondrial Reprogramming Underlies Resistance to BCL-2 Inhibition in Lymphoid Malignancies. <i>Cancer Cell</i> , <b>2019</b> , 36, 369-384.e13	24.3	107
125	Chemoimmunotherapy with O-FC in previously untreated patients with chronic lymphocytic leukemia. <i>Blood</i> , <b>2011</b> , 117, 6450-8	2.2	107
124	Multivariable model for time to first treatment in patients with chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , <b>2011</b> , 29, 4088-95	2.2	105
123	Correlation between FDG/PET, histology, characteristics, and survival in 332 patients with chronic lymphoid leukemia. <i>Blood</i> , <b>2014</b> , 123, 2783-90	2.2	97
122	Economic Burden of Chronic Lymphocytic Leukemia in the Era of Oral Targeted Therapies in the United States. <i>Journal of Clinical Oncology</i> , <b>2017</b> , 35, 166-174	2.2	94
121	Efficacy of venetoclax in relapsed chronic lymphocytic leukemia is influenced by disease and response variables. <i>Blood</i> , <b>2019</b> , 134, 111-122	2.2	94
120	Second cancers in patients with chronic lymphocytic leukemia who received frontline fludarabine, cyclophosphamide and rituximab therapy: distribution and clinical outcomes. <i>Leukemia and Lymphoma</i> , <b>2015</b> , 56, 1643-50	1.9	93

119	Venetoclax for patients with chronic lymphocytic leukemia who progressed during or after idelalisib therapy. <i>Blood</i> , <b>2018</b> , 131, 1704-1711	2.2	93
118	Comprehensive Safety Analysis of Venetoclax Monotherapy for Patients with Relapsed/Refractory Chronic Lymphocytic Leukemia. <i>Clinical Cancer Research</i> , <b>2018</b> , 24, 4371-4379	12.9	90
117	Eliminating minimal residual disease as a therapeutic end point: working toward cure for patients with CLL. <i>Blood</i> , <b>2016</b> , 127, 279-86	2.2	89
116	Homoharringtonine reduced Mcl-1 expression and induced apoptosis in chronic lymphocytic leukemia. <i>Blood</i> , <b>2011</b> , 117, 156-64	2.2	85
115	The antileukemia activity of a human anti-CD40 antagonist antibody, HCD122, on human chronic lymphocytic leukemia cells. <i>Blood</i> , <b>2008</b> , 112, 711-20	2.2	85
114	The natural history of fludarabine-refractory chronic lymphocytic leukemia patients who fail alemtuzumab or have bulky lymphadenopathy. <i>Leukemia and Lymphoma</i> , <b>2007</b> , 48, 1931-9	1.9	85
113	Long-term outcomes for patients with chronic lymphocytic leukemia who discontinue ibrutinib. <i>Cancer</i> , <b>2017</b> , 123, 2268-2273	6.4	83
112	Comparison of Acalabrutinib, A Selective Bruton Tyrosine Kinase Inhibitor, with Ibrutinib in Chronic Lymphocytic Leukemia Cells. <i>Clinical Cancer Research</i> , <b>2017</b> , 23, 3734-3743	12.9	82
111	Acalabrutinib monotherapy in patients with relapsed/refractory chronic lymphocytic leukemia: updated phase 2 results. <i>Blood</i> , <b>2020</b> , 135, 1204-1213	2.2	81
110	Pirtobrutinib in relapsed or refractory B-cell malignancies (BRUIN): a phase 1/2 study. <i>Lancet, The</i> , <b>2021</b> , 397, 892-901	4.0	81
109	NCCN Guidelines Insights: Non-Hodgkin's Lymphomas, Version 3.2016. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , <b>2016</b> , 14, 1067-79	7.3	80
108	Phase 1/2 study of lumiliximab combined with fludarabine, cyclophosphamide, and rituximab in patients with relapsed or refractory chronic lymphocytic leukemia. <i>Blood</i> , <b>2010</b> , 115, 489-95	2.2	75
107	Therapeutic advances in leukemia and myelodysplastic syndrome over the past 40 years. <i>Cancer</i> , <b>2008</b> , 113, 1933-52	6.4	74
106	Relevance of the immunoglobulin VH somatic mutation status in patients with chronic lymphocytic leukemia treated with fludarabine, cyclophosphamide, and rituximab (FCR) or related chemoimmunotherapy regimens. <i>Blood</i> , <b>2009</b> , 113, 3168-71	2.2	73
105	Ibrutinib Therapy Increases T Cell Repertoire Diversity in Patients with Chronic Lymphocytic Leukemia. <i>Journal of Immunology</i> , <b>2017</b> , 198, 1740-1747	5.3	71
104	Frontline chemoimmunotherapy with fludarabine, cyclophosphamide, alemtuzumab, and rituximab for high-risk chronic lymphocytic leukemia. <i>Blood</i> , <b>2011</b> , 118, 2062-8	2.2	70
103	Characteristics associated with important clinical end points in patients with chronic lymphocytic leukemia at initial treatment. <i>Journal of Clinical Oncology</i> , <b>2009</b> , 27, 1637-43	2.2	67
102	A pilot study of lower doses of ibrutinib in patients with chronic lymphocytic leukemia. <i>Blood</i> , <b>2018</b> , 132, 2249-2259	2.2	63

101	A retrospective comparison of three sequential groups of patients with Recurrent/Refractory chronic lymphocytic leukemia treated with fludarabine-based regimens. <i>Cancer</i> , <b>2006</b> , 106, 337-45	6.4	56
100	Outcomes of first-line treatment for chronic lymphocytic leukemia with 17p deletion. <i>Haematologica</i> , <b>2014</b> , 99, 1350-5	6.6	55
99	Targeted multigene deep sequencing of Bruton tyrosine kinase inhibitor-resistant chronic lymphocytic leukemia with disease progression and Richter transformation. <i>Cancer</i> , <b>2019</b> , 125, 559-574	6.4	51
98	NCCN Guidelines Insights: Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma, Version 1.2017. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , <b>2017</b> , 15, 293-311	7.3	49
97	Cyclophosphamide, fludarabine, alemtuzumab, and rituximab as salvage therapy for heavily pretreated patients with chronic lymphocytic leukemia. <i>Blood</i> , <b>2011</b> , 118, 2085-93	2.2	49
96	Myelosuppression after frontline fludarabine, cyclophosphamide, and rituximab in patients with chronic lymphocytic leukemia: analysis of persistent and new-onset cytopenia. <i>Cancer</i> , <b>2013</b> , 119, 3805-11	6.4	48
95	Non-Hodgkin's lymphomas. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , <b>2011</b> , 9, 484-560	5.0	47
94	Novel immune-based treatment strategies for chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , <b>2005</b> , 23, 6325-32	2.2	46
93	International prognostic score for asymptomatic early-stage chronic lymphocytic leukemia. <i>Blood</i> , <b>2020</b> , 135, 1859-1869	2.2	45
92	Long-term Follow-up of Treatment with Ibrutinib and Rituximab in Patients with High-Risk Chronic Lymphocytic Leukemia. <i>Clinical Cancer Research</i> , <b>2017</b> , 23, 2154-2158	12.9	43
91	Venetoclax (VEN) Monotherapy for Patients with Chronic Lymphocytic Leukemia (CLL) Who Relapsed after or Were Refractory to Ibrutinib or Idelalisib. <i>Blood</i> , <b>2016</b> , 128, 637-637	2.2	43
90	Phase 1b study of venetoclax-obinutuzumab in previously untreated and relapsed/refractory chronic lymphocytic leukemia. <i>Blood</i> , <b>2019</b> , 133, 2765-2775	2.2	42
89	Fludarabine, cyclophosphamide, mitoxantrone plus rituximab (FCM-R) in frontline CLL. <i>Leukemia Research</i> , <b>2010</b> , 34, 284-8	2.7	41
88	Fludarabine and mitoxantrone for patients with chronic lymphocytic leukemia. <i>Cancer</i> , <b>2004</b> , 100, 2583-91	3.4	38
87	Chronic lymphocytic leukemia/small lymphocytic lymphoma, version 1.2015. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , <b>2015</b> , 13, 326-62	7.3	37
86	Multiple-dose granulocyte-macrophage-colony-stimulating factor plus 23-valent polysaccharide pneumococcal vaccine in patients with chronic lymphocytic leukemia: a prospective, randomized trial of safety and immunogenicity. <i>Cancer</i> , <b>2008</b> , 113, 383-7	6.4	36
85	Evaluation of 230 patients with relapsed/refractory deletion 17p chronic lymphocytic leukaemia treated with ibrutinib from 3 clinical trials. <i>British Journal of Haematology</i> , <b>2018</b> , 182, 504-512	4.5	32
84	Ibrutinib (Ibr) Plus Venetoclax (Ven) for First-Line Treatment of Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Results from the MRD Cohort of the Phase 2 CAPTIVATE Study. <i>Blood</i> , <b>2019</b> , 134, 35-35	2.2	30



83	LDOC1 mRNA is differentially expressed in chronic lymphocytic leukemia and predicts overall survival in untreated patients. <i>Blood</i> , <b>2011</b> , 117, 4076-84	2.2	28
82	AMG-176, an Mcl-1 Antagonist, Shows Preclinical Efficacy in Chronic Lymphocytic Leukemia. <i>Clinical Cancer Research</i> , <b>2020</b> , 26, 3856-3867	12.9	26
81	Ibrutinib (Ibr) Plus Venetoclax (Ven) for First-Line Treatment of Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): 1-Year Disease-Free Survival (DFS) Results From the MRD Cohort of the Phase 2 CAPTIVATE Study. <i>Blood</i> , <b>2020</b> , 136, 16-17	2.2	26
80	Clinical implications of cancer gene mutations in patients with chronic lymphocytic leukemia treated with lenalidomide. <i>Blood</i> , <b>2018</b> , 131, 1820-1832	2.2	25
79	B-cell Receptor Signaling Regulates Metabolism in Chronic Lymphocytic Leukemia. <i>Molecular Cancer Research</i> , <b>2017</b> , 15, 1692-1703	6.6	23
78	Population pharmacokinetics of ofatumumab in patients with chronic lymphocytic leukemia, follicular lymphoma, and rheumatoid arthritis. <i>Journal of Clinical Pharmacology</i> , <b>2014</b> , 54, 818-27	2.9	23
77	Three newly approved drugs for chronic lymphocytic leukemia: incorporating ibrutinib, idelalisib, and obinutuzumab into clinical practice. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , <b>2015</b> , 15, 385-91	2	22
76	Ofatumumab and Lenalidomide for Patients with Relapsed or Refractory Chronic Lymphocytic Leukemia: Correlation between Responses and Immune Characteristics. <i>Clinical Cancer Research</i> , <b>2016</b> , 22, 2359-67	12.9	22
75	What is the best frontline therapy for patients with CLL and 17p deletion?. <i>Current Hematologic Malignancy Reports</i> , <b>2011</b> , 6, 36-46	4.4	22
74	Investigational immunotherapeutics for B-cell malignancies. <i>Journal of Clinical Oncology</i> , <b>2010</b> , 28, 884-92.2		21
73	KTE-X19 anti-CD19 CAR T-cell therapy in adult relapsed/refractory acute lymphoblastic leukemia: ZUMA-3 phase 1 results. <i>Blood</i> , <b>2021</b> , 138, 11-22	2.2	21
72	Phase 2 CAPTIVATE results of ibrutinib (ibr) plus venetoclax (ven) in first-line chronic lymphocytic leukemia (CLL).. <i>Journal of Clinical Oncology</i> , <b>2018</b> , 36, 7502-7502	2.2	20
71	Association of gene mutations with time-to-first treatment in 384 treatment-naive chronic lymphocytic leukaemia patients. <i>British Journal of Haematology</i> , <b>2019</b> , 187, 307-318	4.5	19
70	Immunohistochemical detection of ZAP70 in chronic lymphocytic leukemia predicts immunoglobulin heavy chain gene mutation status and time to progression. <i>Modern Pathology</i> , <b>2010</b> , 23, 1518-23	9.8	19
69	Minimal residual disease undetectable by next-generation sequencing predicts improved outcome in CLL after chemoimmunotherapy. <i>Blood</i> , <b>2019</b> , 134, 1951-1959	2.2	19
68	Acalabrutinib in treatment-naive chronic lymphocytic leukemia. <i>Blood</i> , <b>2021</b> , 137, 3327-3338	2.2	18
67	Self-administered, subcutaneous alemtuzumab to treat residual disease in patients with chronic lymphocytic leukemia. <i>Cancer</i> , <b>2011</b> , 117, 116-24	6.4	17
66	Ibrutinib Plus Venetoclax for First-Line Treatment of Chronic Lymphocytic Leukemia: Primary Analysis Results From the Minimal Residual Disease Cohort of the Randomized Phase II CAPTIVATE Study. <i>Journal of Clinical Oncology</i> , <b>2021</b> , 39, 3853-3865	2.2	17

65	Serial minimal residual disease (MRD) monitoring during first-line FCR treatment for CLL may direct individualized therapeutic strategies. <i>Leukemia</i> , <b>2018</b> , 32, 2388-2398	10.7	17
64	Ibrutinib Plus Venetoclax for First-line Treatment of Chronic Lymphocytic Leukemia: A Nonrandomized Phase 2 Trial. <i>JAMA Oncology</i> , <b>2021</b> , 7, 1213-1219	13.4	17
63	Evaluation of bendamustine in combination with fludarabine in primary chronic lymphocytic leukemia cells. <i>Blood</i> , <b>2014</b> , 123, 3780-9	2.2	16
62	Initial therapy for patients with chronic lymphocytic leukemia. <i>Seminars in Oncology</i> , <b>2006</b> , 33, 202-9	5.5	16
61	Statin and aspirin use is associated with improved outcome of FCR therapy in relapsed/refractory chronic lymphocytic leukemia. <i>Blood</i> , <b>2014</b> , 123, 1424-6	2.2	15
60	Efficacy and safety in a 4-year follow-up of the ELEVATE-TN study comparing acalabrutinib with or without obinutuzumab versus obinutuzumab plus chlorambucil in treatment-naïve chronic lymphocytic leukemia.. <i>Leukemia</i> , <b>2022</b> ,	10.7	15
59	Routine sequencing in CLL has prognostic implications and provides new insight into pathogenesis and targeted treatments. <i>British Journal of Haematology</i> , <b>2019</b> , 185, 852-864	4.5	14
58	$\kappa$ -microglobulin normalization within 6 months of ibrutinib-based treatment is associated with superior progression-free survival in patients with chronic lymphocytic leukemia. <i>Cancer</i> , <b>2016</b> , 122, 565-73	6.4	14
57	Ofatumumab Combined with Fludarabine and Cyclophosphamide (O-FC) Shows High Activity in Patients with Previously Untreated Chronic Lymphocytic Leukemia (CLL): Results From a Randomized, Multicenter, International, Two-Dose, Parallel Group, Phase II Trial.. <i>Blood</i> , <b>2009</b> , 114, 207-207	2.2	14
56	Autologous CD33-CAR-T cells for treatment of relapsed/refractory acute myelogenous leukemia. <i>Leukemia</i> , <b>2021</b> , 35, 3282-3286	10.7	14
55	Targeting BCL2 in Chronic Lymphocytic Leukemia and Other Hematologic Malignancies. <i>Drugs</i> , <b>2019</b> , 79, 1287-1304	12.1	13
54	Current and investigational therapies for patients with CLL. <i>Hematology American Society of Hematology Education Program</i> , <b>2006</b> , 2006, 285-94	3.1	13
53	Phase 1 TRANSCEND CLL 004 study of lisocabtagene maraleucel in patients with relapsed/refractory CLL or SLL. <i>Blood</i> , <b>2021</b> ,	2.2	13
52	Ofatumumab monotherapy in fludarabine-refractory chronic lymphocytic leukemia: final results from a pivotal study. <i>Haematologica</i> , <b>2015</b> , 100, e311-4	6.6	12
51	Azacitidine in fludarabine-refractory chronic lymphocytic leukemia: a phase II study. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , <b>2013</b> , 13, 292-5	2	12
50	Ofatumumab retreatment and maintenance in fludarabine-refractory chronic lymphocytic leukaemia patients. <i>British Journal of Haematology</i> , <b>2015</b> , 170, 40-9	4.5	12
49	Ibrutinib: a paradigm shift in management of CLL. <i>Expert Review of Hematology</i> , <b>2014</b> , 7, 705-17	2.8	12
48	Circulating CD52 and CD20 levels at end of treatment predict for progression and survival in patients with chronic lymphocytic leukaemia treated with fludarabine, cyclophosphamide and rituximab (FCR). <i>British Journal of Haematology</i> , <b>2010</b> , 148, 386-93	4.5	12



47	Genetics and molecular biology of chronic lymphocytic leukemia. <i>Current Treatment Options in Oncology</i> , <b>2005</b> , 6, 215-25	5.4	12
46	Salvage Therapy with Combined Cyclophosphamide (C), Fludarabine (F), Alemtuzumab (A), and Rituximab (R) (CFAR) for Heavily Pre-Treated Patients with CLL.. <i>Blood</i> , <b>2005</b> , 106, 719-719	2.2	12
45	An Ongoing Phase 1/2a Study of ABT-263; Pharmacokinetics (PK), Safety and Anti-Tumor Activity in Patients (pts) with Relapsed or Refractory Chronic Lymphocytic Leukemia (CLL).. <i>Blood</i> , <b>2009</b> , 114, 883-883	2.2	12
44	Gene therapy and active immune therapy of hematologic malignancies. <i>Best Practice and Research in Clinical Haematology</i> , <b>2007</b> , 20, 557-68	4.2	11
43	Combined Cyclophosphamide, Fludarabine, Alemtuzumab, and Rituximab (CFAR) Is Active for Relapsed and Refractory Patients with CLL.. <i>Blood</i> , <b>2004</b> , 104, 340-340	2.2	11
42	Fludarabine, cyclophosphamide, and multiple-dose rituximab as frontline therapy for chronic lymphocytic leukemia. <i>Cancer</i> , <b>2015</b> , 121, 3869-76	6.4	10
41	Chronic lymphocytic leukemia: treatment options for patients with refractory disease. <i>Cancer</i> , <b>2009</b> , 115, 3830-41	6.4	10
40	Fixed-duration ibrutinib plus venetoclax for first-line treatment of CLL: primary analysis of the CAPTIVATE FD cohort.. <i>Blood</i> , <b>2022</b> ,	2.2	10
39	Creating novel translation inhibitors to target pro-survival proteins in chronic lymphocytic leukemia. <i>Leukemia</i> , <b>2019</b> , 33, 1663-1674	10.7	9
38	Fludarabine, cyclophosphamide and rituximab plus granulocyte macrophage colony-stimulating factor as frontline treatment for patients with chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , <b>2014</b> , 55, 828-33	1.9	9
37	Killing of chronic lymphocytic leukemia by the combination of fludarabine and oxaliplatin is dependent on the activity of XPF endonuclease. <i>Clinical Cancer Research</i> , <b>2011</b> , 17, 4731-41	12.9	9
36	Chronic lymphocytic leukemia: new concepts for future therapy. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , <b>2010</b> , 10, 369-78	2	9
35	Dynamic changes of the normal B lymphocyte repertoire in CLL in response to ibrutinib or FCR chemo-immunotherapy. <i>OncImmunology</i> , <b>2018</b> , 7, e1417720	7.2	7
34	Lenalidomide-induced graft-vs.-Leukemia effect in a patient with chronic lymphocytic leukemia who relapsed after allogeneic stem cell transplant. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , <b>2014</b> , 14, e105-9	2	7
33	Combined Ibrutinib and Venetoclax for First-Line Treatment for Patients with Chronic Lymphocytic Leukemia (CLL): Focus on MRD Results. <i>Blood</i> , <b>2020</b> , 136, 42-43	2.2	7
32	Clinical Efficacy of Lenalidomide in Fludarabine-Refractory Chronic Lymphocytic Leukemia Patients.. <i>Blood</i> , <b>2007</b> , 110, 3108-3108	2.2	7
31	Combination Therapy with Lenalidomide and Rituximab in Patients with Relapsed Chronic Lymphocytic Leukemia (CLL).. <i>Blood</i> , <b>2009</b> , 114, 206-206	2.2	7
30	Incidental Richter transformation in chronic lymphocytic leukemia patients during temporary interruption of ibrutinib. <i>Blood Advances</i> , <b>2020</b> , 4, 4508-4511	7.8	7

29	PET-positive lymphadenopathy in CLL-Not always Richter transformation. <i>American Journal of Hematology</i> , <b>2017</b> , 92, 405-406	7.1	6
28	Lenalidomide as Initial Treatment of Elderly Patients with Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , <b>2008</b> , 112, 45-45	2.2	6
27	Final Analysis From the International Trial of Single-Agent Ofatumumab In Patients with Fludarabine-Refractory Chronic Lymphocytic Leukemia. <i>Blood</i> , <b>2010</b> , 116, 921-921	2.2	6
26	Measurable residual disease in chronic lymphocytic leukemia: expert review and consensus recommendations. <i>Leukemia</i> , <b>2021</b> , 35, 3059-3072	10.7	6
25	Venetoclax for chronic lymphocytic leukaemia patients who progress after more than one B-cell receptor pathway inhibitor. <i>British Journal of Haematology</i> , <b>2019</b> , 185, 961-966	4.5	6
24	Long-term follow-up of patients receiving allogeneic stem cell transplant for chronic lymphocytic leukaemia: mixed T-cell chimerism is associated with high relapse risk and inferior survival. <i>British Journal of Haematology</i> , <b>2017</b> , 177, 567-577	4.5	5
23	A phase I-II trial of fludarabine, bendamustine and rituximab (FBR) in previously treated patients with CLL. <i>Oncotarget</i> , <b>2017</b> , 8, 22104-22112	3.3	5
22	Ibrutinib, fludarabine, cyclophosphamide, and obinutuzumab (iFCG) regimen for chronic lymphocytic leukemia (CLL) with mutated IGHV and without TP53 aberrations. <i>Leukemia</i> , <b>2021</b> , 35, 3421-3429	10.7	5
21	Tumour debulking and reduction in predicted risk of tumour lysis syndrome with single-agent ibrutinib in patients with chronic lymphocytic leukaemia. <i>British Journal of Haematology</i> , <b>2019</b> , 186, 184-188	4.5	5
20	A phase II trial of eltrombopag for patients with chronic lymphocytic leukaemia (CLL) and thrombocytopenia. <i>British Journal of Haematology</i> , <b>2019</b> , 185, 606-608	4.5	5
19	Expression of BCL2 alternative proteins and association with outcome in CLL patients treated with venetoclax. <i>Leukemia and Lymphoma</i> , <b>2021</b> , 62, 1129-1135	1.9	5
18	Tumour lysis syndrome in patients with chronic lymphocytic leukaemia treated with BCL-2 inhibitors: risk factors, prophylaxis, and treatment recommendations. <i>Lancet Haematology</i> , <b>2020</b> , 7, e168-e176	14.6	4
17	Proteomics profiling identifies induction of caveolin-1 in chronic lymphocytic leukemia cells by bone marrow stromal cells. <i>Leukemia and Lymphoma</i> , <b>2018</b> , 59, 1427-1438	1.9	4
16	Updates to the management of chronic lymphocytic leukemia. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , <b>2015</b> , 13, 662-5	7.3	4
15	The Role of BTK Inhibition in the Treatment of Chronic Lymphocytic Leukemia: A Clinical View. <i>Journal of Experimental Pharmacology</i> , <b>2021</b> , 13, 923-935	3	4
14	CXCL13 plasma levels function as a biomarker for disease activity in patients with chronic lymphocytic leukemia. <i>Leukemia</i> , <b>2021</b> , 35, 1610-1620	10.7	4
13	Sustained long-lasting responses after lenalidomide discontinuation in patients with chronic lymphocytic leukemia. <i>Leukemia</i> , <b>2018</b> , 32, 2278-2281	10.7	3
12	Associations of ofatumumab exposure and treatment outcomes in patients with untreated CLL receiving chemoimmunotherapy. <i>Leukemia and Lymphoma</i> , <b>2017</b> , 58, 348-356	1.9	2

11	The landscape of genetic mutations in patients with chronic lymphocytic leukaemia and complex karyotype. <i>British Journal of Haematology</i> , <b>2019</b> , 187, e1-e4	4.5	2
10	Immunologic monitoring in chronic lymphocytic leukemia. <i>Current Oncology Reports</i> , <b>2003</b> , 5, 419-25	6.3	2
9	Clinical outcome of allogeneic stem cell transplantation in patients with B-cell lymphoid malignancies following treatment with targeted small molecule inhibitors.. <i>Leukemia and Lymphoma</i> , <b>2022</b> , 1-9	1.9	2
8	Cyclin-dependent kinase inhibitor fadraciclib (CYC065) depletes anti-apoptotic protein and synergizes with venetoclax in primary chronic lymphocytic leukemia cells.. <i>Leukemia</i> , <b>2022</b> ,	10.7	2
7	Critical appraisal of the role of rituximab in the treatment of patients with previously untreated or treated chronic lymphocytic leukemia (CLL). <i>Journal of Blood Medicine</i> , <b>2010</b> , 1, 115-22	2.3	1
6	Chronic Lymphocytic Leukemia: How to Assess Prognosis in 2007. <i>Clinical Leukemia</i> , <b>2007</b> , 1, 162-167		1
5	RPPA-based proteomics recognizes distinct epigenetic signatures in chronic lymphocytic leukemia with clinical consequences. <i>Leukemia</i> , <b>2021</b> ,	10.7	1
4	Clinical and molecular characteristics and treatment patterns of adolescent and young adult patients with chronic lymphocytic leukaemia. <i>British Journal of Haematology</i> , <b>2021</b> , 194, 61-68	4.5	1
3	Proteomic profiling based classification of CLL provides prognostication for modern therapy and identifies novel therapeutic targets.. <i>Blood Cancer Journal</i> , <b>2022</b> , 12, 43	7	1
2	New aspects of the treatment of chronic lymphocytic leukemia. <i>Current Hematologic Malignancy Reports</i> , <b>2006</b> , 1, 251-7	4.4	
1	HLA Homozygosity and Haplotype Bias Among Patients with Chronic Lymphocytic Leukemia: Implications for Disease Control by Physiologic Immune Surveillance. <i>Blood</i> , <b>2010</b> , 116, 1370-1370	2.2	