

# Hannu Juhani Koponen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5921551/publications.pdf>

Version: 2024-02-01

99  
papers

2,797  
citations

218592

26  
h-index

197736

49  
g-index

101  
all docs

101  
docs citations

101  
times ranked

3932  
citing authors

#	ARTICLE	IF	CITATIONS
1	Work incapacity among family caregivers: a record linkage study. <i>Journal of Epidemiology and Community Health</i> , 2022, 76, 580-585.	2.0	1
2	A longitudinal study of restless legs symptoms among patients with depression. <i>Nordic Journal of Psychiatry</i> , 2022, 76, 584-590.	0.7	3
3	Incidence and mortality of alcohol-related dementia and Wernicke-Korsakoff syndrome: A nationwide register study. <i>International Journal of Geriatric Psychiatry</i> , 2022, 37, .	1.3	10
4	Characteristics and predictors of off-label use of antipsychotics in general population sample. <i>Acta Psychiatrica Scandinavica</i> , 2022, 146, 227-239.	2.2	4
5	Criminal Behavior in the Four Years Preceding Diagnosis of Neurocognitive Disorder: A Nationwide Register Study in Finland. <i>American Journal of Geriatric Psychiatry</i> , 2021, 29, 657-665.	0.6	10
6	Physical activity and ability to meet different work requirements among adult working men with or without current depressive symptoms. <i>International Archives of Occupational and Environmental Health</i> , 2021, 94, 451-458.	1.1	1
7	Use of antidepressants among Finnish family caregivers: a nationwide register-based study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2021, 56, 2209-2216.	1.6	4
8	Predictors of response to pharmacological treatments in treatment-resistant schizophrenia – A systematic review and meta-analysis. <i>Schizophrenia Research</i> , 2021, 236, 123-134.	1.1	10
9	Does baseline leisure-time physical activity level predict future depressive symptoms or physical activity among depressive patients? Findings from a Finnish five-year cohort study. <i>Nordic Journal of Psychiatry</i> , 2021, 75, 356-361.	0.7	1
10	The influence of restless legs symptoms on musculoskeletal pain in depression. <i>Scandinavian Journal of Pain</i> , 2020, 20, 603-610.	0.5	0
11	The profiles of health care utilization among a non-depressed population and patients with depressive symptoms with and without clinical depression. <i>Scandinavian Journal of Primary Health Care</i> , 2019, 37, 312-318.	0.6	18
12	Leisure time physical activity and its relation to psychiatric comorbidities in depression. Findings from Finnish Depression and Metabolic Syndrome in Adults (FDMSA) study. <i>Journal of Affective Disorders</i> , 2019, 259, 150-153.	2.0	8
13	Prevalence of restless legs symptoms according to depressive symptoms and depression type: a cross-sectional study. <i>Nordic Journal of Psychiatry</i> , 2018, 72, 51-56.	0.7	15
14	Elevation of tumor necrosis factor alpha levels is associated with restless legs symptoms in clinically depressed patients. <i>Journal of Psychosomatic Research</i> , 2018, 115, 1-5.	1.2	5
15	T78. LONG-TERM PROGNOSIS OF SCHIZOPHRENIA - RESULTS FROM THE NORTHERN FINLAND BIRTH COHORT 1966. <i>Schizophrenia Bulletin</i> , 2018, 44, S144-S145.	2.3	0
16	Does level of leisure time physical activity, in a sample of patients with depression, predict health care utilization over a subsequent 5-year period? Findings from a Finnish cohort study. <i>Mental Health and Physical Activity</i> , 2018, 15, 40-44.	0.9	5
17	Antipsychotic and benzodiazepine use and brain morphology in schizophrenia and affective psychoses – Systematic reviews and birth cohort study. <i>Psychiatry Research - Neuroimaging</i> , 2018, 281, 43-52.	0.9	3
18	Musculoskeletal Pain in Melancholic and Atypical Depression. <i>Pain Medicine</i> , 2017, 18, pnw202.	0.9	8

#	ARTICLE	IF	CITATIONS
19	Lifetime antipsychotic medication and cognitive performance in schizophrenia at age 43 years in a general population birth cohort. <i>Psychiatry Research</i> , 2017, 247, 130-138.	1.7	68
20	Physical restraints and associations with neuropsychiatric symptoms and personal characteristics in residential care: a cross-sectional study. <i>International Journal of Geriatric Psychiatry</i> , 2017, 32, 1418-1424.	1.3	5
21	Long-term antipsychotic and benzodiazepine use and brain volume changes in schizophrenia: The Northern Finland Birth Cohort 1966 study. <i>Psychiatry Research - Neuroimaging</i> , 2017, 266, 73-82.	0.9	21
22	Cost-utility analysis of vortioxetine versus agomelatine, bupropion SR, sertraline and venlafaxine XR after treatment switch in major depressive disorder in Finland. <i>Expert Review of Pharmacoeconomics and Outcomes Research</i> , 2017, 17, 293-302.	0.7	17
23	Antipsychotic drug use and associations with neuropsychiatric symptoms in persons with impaired cognition: a cross-sectional study. <i>Nordic Journal of Psychiatry</i> , 2016, 70, 621-625.	0.7	7
24	Depressive symptoms are associated with poor outcome for lumbar spine surgery. <i>Scandinavian Journal of Pain</i> , 2016, 12, 13-17.	0.5	10
25	Use of psychiatric medications in schizophrenia and other psychoses in a general population sample. <i>Psychiatry Research</i> , 2016, 235, 160-168.	1.7	11
26	Talking about paranoid experiences: Interactional dynamics of accomplishing disclosure of psychotic symptoms in triadic conversation. <i>Communication and Medicine</i> , 2016, 12, 25-40.	0.1	3
27	Cardiometabolic risk factors in patients referred to depression nurse case managers. <i>Nordic Journal of Psychiatry</i> , 2015, 69, 262-267.	0.7	11
28	Use of anti-dementia drugs in home care and residential care and associations with neuropsychiatric symptoms: a cross-sectional study. <i>BMC Geriatrics</i> , 2015, 15, 100.	1.1	8
29	Association between suicidal behaviour and impaired glucose metabolism in depressive disorders. <i>BMC Psychiatry</i> , 2015, 15, 163.	1.1	68
30	Altered resting-state activity in seasonal affective disorder. <i>Human Brain Mapping</i> , 2014, 35, 161-172.	1.9	30
31	Use of antidepressant medication and suicidal ideation in the Northern Finland Birth Cohort 1966. <i>Human Psychopharmacology</i> , 2014, 29, 559-567.	0.7	3
32	Lifetime use of antipsychotic medication and its relation to change of verbal learning and memory in midlife schizophrenia - An observational 9-year follow-up study. <i>Schizophrenia Research</i> , 2014, 158, 134-141.	1.1	66
33	Longitudinal Changes in Total Brain Volume in Schizophrenia: Relation to Symptom Severity, Cognition and Antipsychotic Medication. <i>PLoS ONE</i> , 2014, 9, e101689.	1.1	92
34	Association between vitamin b12 levels and melancholic depressive symptoms: a Finnish population-based study. <i>BMC Psychiatry</i> , 2013, 13, 145.	1.1	32
35	Association between the duration of untreated psychosis and short- and long-term outcome in schizophrenia within the Northern Finland 1966 Birth Cohort. <i>Schizophrenia Research</i> , 2013, 143, 3-10.	1.1	25
36	Financial satisfaction and its relationship to depressive symptoms in middle-aged and older adults: Results from the FIN-D2D survey. <i>International Journal of Social Psychiatry</i> , 2013, 59, 239-246.	1.6	17

#	ARTICLE	IF	CITATIONS
37	Disordered semantic activation in disorganized discourse in schizophrenia: a new pragmatic linguistic tool for structure and meaning reconstruction. <i>International Journal of Language and Communication Disorders</i> , 2013, 48, 320-328.	0.7	4
38	Use of antipsychotic medication and suicidality in the Northern Finland Birth Cohort 1966. <i>Human Psychopharmacology</i> , 2012, 27, 476-485.	0.7	9
39	Outcome of patients after market withdrawal of thioridazine: A retrospective analysis in a nationwide cohort. <i>Pharmacoepidemiology and Drug Safety</i> , 2012, 21, 1227-1231.	0.9	28
40	Prevalence of metabolic syndrome in subjects with melancholic and non-melancholic depressive symptoms. A Finnish population-based study. <i>Journal of Affective Disorders</i> , 2012, 136, 543-549.	2.0	40
41	Association between folate intake and melancholic depressive symptoms. A Finnish population-based study. <i>Journal of Affective Disorders</i> , 2012, 138, 473-478.	2.0	12
42	Lifetime leisure-time physical activity and the risk of depressive symptoms at the ages of 65-74 years: The FIN-D2D survey. <i>Preventive Medicine</i> , 2012, 54, 313-315.	1.6	16
43	Psychological Distress Predicts the Development of the Metabolic Syndrome: A Prospective Population-Based Study. <i>Psychosomatic Medicine</i> , 2011, 73, 158-165.	1.3	58
44	Separable developmental trajectories in schizophrenia from womb to grave. Lifespan results from the Northern Finland 1966 Birth Cohort (NFBC 1966). <i>International Clinical Psychopharmacology</i> , 2011, 26, e95-e96.	0.9	0
45	Psychological distress and C-reactive protein: do health behaviours and pathophysiological factors modify the association?. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2011, 261, 277-284.	1.8	20
46	Association of Depressive Symptoms With Impaired Glucose Regulation, Screen-Detected, and Previously Known Type 2 Diabetes. <i>Diabetes Care</i> , 2011, 34, 71-76.	4.3	32
47	Meta-analysis of Paternal Age and Schizophrenia Risk in Male Versus Female Offspring. <i>Schizophrenia Bulletin</i> , 2011, 37, 1039-1047.	2.3	167
48	Advanced Paternal Age, Mortality, and Suicide in the General Population. <i>Journal of Nervous and Mental Disease</i> , 2010, 198, 404-411.	0.5	15
49	Agreement between self-reported and pharmacy data on medication use in the Northern Finland 1966 Birth Cohort. <i>International Journal of Methods in Psychiatric Research</i> , 2010, 19, 88-96.	1.1	42
50	Rate of Cannabis Use Disorders in Clinical Samples of Patients With Schizophrenia: A Meta-analysis. <i>Schizophrenia Bulletin</i> , 2010, 36, 1115-1130.	2.3	275
51	The prevalence and predictive value of individual criteria for metabolic syndrome in schizophrenia: A Northern Finland 1966 Birth Cohort Study. <i>World Journal of Biological Psychiatry</i> , 2010, 11, 262-267.	1.3	18
52	Depressive symptoms and 10-year risk for cardiovascular morbidity and mortality. <i>World Journal of Biological Psychiatry</i> , 2010, 11, 834-839.	1.3	42
53	Association between duration of untreated psychosis and brain morphology in schizophrenia within the Northern Finland 1966 Birth Cohort. <i>Schizophrenia Research</i> , 2010, 123, 145-152.	1.1	35
54	Effect of menopause and use of contraceptives/hormone therapy on association of C-reactive protein and depression: A population-based study. <i>Journal of Psychosomatic Research</i> , 2010, 68, 573-579.	1.2	5

#	ARTICLE	IF	CITATIONS
55	Negative symptoms and their predictors in schizophrenia within the Northern Finland 1966 Birth Cohort. <i>Psychiatry Research</i> , 2010, 178, 121-125.	1.7	21
56	Leisure-time physical activity and metabolic syndrome plus depressive symptoms in the FIN-D2D survey. <i>Preventive Medicine</i> , 2010, 51, 466-470.	1.6	16
57	Gender-specific association of psychological distress with cardiovascular risk scores. <i>Scandinavian Journal of Primary Health Care</i> , 2010, 28, 36-40.	0.6	12
58	Management of depression in the presence of pain symptoms. <i>Psychiatria Danubina</i> , 2010, 22, 4-13.	0.2	14
59	Ziprasidone versus olanzapine, risperidone or quetiapine in patients with chronic schizophrenia: A 12-week open-label, multicentre clinical trial. <i>World Journal of Biological Psychiatry</i> , 2009, 10, 710-718.	1.3	14
60	Depressive symptomatology is associated with decreased interleukin-1 beta and increased interleukin-1 receptor antagonist levels in males. <i>Psychiatry Research</i> , 2009, 167, 73-79.	1.7	47
61	Suicides Among Older Persons in Finland and Time Since Hospitalization Discharge. <i>Psychiatric Services</i> , 2009, 60, 390-393.	1.1	12
62	Schizophrenia and sudden cardiac deathâ€”A review. <i>Nordic Journal of Psychiatry</i> , 2008, 62, 342-345.	0.7	99
63	Metabolic Syndrome Predisposes to Depressive Symptoms. <i>Journal of Clinical Psychiatry</i> , 2008, 69, 178-182.	1.1	145
64	Socio-demographic and clinical predictors of occupational status in schizophrenic psychosesâ€”follow-up within the Northern Finland 1966 Birth Cohort. <i>Psychiatry Research</i> , 2007, 150, 217-225.	1.7	24
65	A double-blind study of the efficacy of venlafaxine extended-release, paroxetine, and placebo in the treatment of panic disorder. <i>Depression and Anxiety</i> , 2007, 24, 1-14.	2.0	88
66	Rates and previous disease history in old age suicide. <i>International Journal of Geriatric Psychiatry</i> , 2007, 22, 38-46.	1.3	36
67	Efficacy of Duloxetine for the Treatment of Generalized Anxiety Disorder. <i>Primary Care Companion To the Journal of Clinical Psychiatry</i> , 2007, 09, 100-107.	0.6	87
68	Patterns of psychiatric hospitalizations in schizophrenic psychoses within the Northern Finland 1966 Birth Cohort. <i>Nordic Journal of Psychiatry</i> , 2006, 60, 286-293.	0.7	13
69	Are females at special risk of obesity if they become psychotic? The longitudinal Northern Finland 1966 Birth Cohort Study. <i>Schizophrenia Research</i> , 2006, 84, 15-19.	1.1	25
70	Statistical methodologies in psychopharmacology: a review. <i>Human Psychopharmacology</i> , 2006, 21, 195-203.	0.7	8
71	Risk factors for schizophrenia. Follow-up data from the Northern Finland 1966 Birth Cohort Study. <i>World Psychiatry</i> , 2006, 5, 168-71.	4.8	16
72	Deramciclane in the treatment of generalized anxiety disorder: A placebo-controlled, double-blind, dose-finding study. <i>European Neuropsychopharmacology</i> , 2005, 15, 617-623.	0.3	16

#	ARTICLE	IF	CITATIONS
73	Sedative Drug Use in the Home-Dwelling Elderly. <i>Annals of Pharmacotherapy</i> , 2004, 38, 2017-2022.	0.9	41
74	Childhood central nervous system infections and risk for schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2004, 254, 9-13.	1.8	95
75	Developmental precursors of psychosis. <i>Current Psychiatry Reports</i> , 2004, 6, 168-175.	2.1	17
76	Reversal of antipsychotic-induced weight gain during quetiapine treatment. <i>International Clinical Psychopharmacology</i> , 2003, 18, 355-356.	0.9	2
77	Weight gain and glucose and lipid metabolism disturbances during antipsychotic medication. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2002, 252, 294-298.	1.8	23
78	Neurochemistry and Delirium. <i>Dementia and Geriatric Cognitive Disorders</i> , 1999, 10, 339-341.	0.7	18
79	A Controlled, Prospective, 1-Year Trial of Citalopram in the Treatment of Panic Disorder. <i>Journal of Clinical Psychiatry</i> , 1998, 59, 528-534.	1.1	95
80	Risperidone in the Treatment of Psychosis and Concomitant Buccolinguomasticatory Dyskinesia in the Elderly. , 1997, 12, 412-413.		3
81	Adolescent conduct disorder, depression, and central nervous system demyelination in magnetic resonance imaging:A report with two cases. <i>Nordic Journal of Psychiatry</i> , 1996, 50, 419-421.	0.7	1
82	Paroxetine increases serum trimipramine concentration. A report of two cases. <i>Human Psychopharmacology</i> , 1995, 10, 345-347.	0.7	17
83	A long-term follow-up study of cerebrospinal fluid 5-hydroxyindoleacetic acid in delirium. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 1994, 244, 131-134.	1.8	14
84	A long-term follow-up study of cerebrospinal fluid acetylcholinesterase in delirium. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 1994, 243, 347-351.	1.8	10
85	Serum mianserin and ageing. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1994, 18, 833-845.	2.5	166
86	Normal neuropathological findings after neuroleptic malignant syndrome. <i>Human Psychopharmacology</i> , 1993, 8, 217-219.	0.7	0
87	Neuroleptic malignant syndrome during remoxipride treatment. A case report. <i>European Neuropsychopharmacology</i> , 1993, 3, 517-519.	0.3	4
88	A prospective study of delirium in elderly patients admitted to a psychiatric hospital. <i>Psychological Medicine</i> , 1993, 23, 103-109.	2.7	51
89	Delirium During Fluoxetine Treatment: A Case Report. <i>Annals of Clinical Psychiatry</i> , 1993, 5, 255-257.	0.6	10
90	Clinical picture and long-term course of epileptic seizures that occur during clozapine treatment. <i>Psychiatry Research</i> , 1992, 44, 107-112.	1.7	30

#	ARTICLE	IF	CITATIONS
91	A longitudinal study of cerebrospinal fluid acetylcholinesterase in delirium: Changes at the acute stage and at one-year followup. <i>Psychiatry Research</i> , 1991, 38, 135-142.	1.7	5
92	Delirium in the Elderly: A Brief Overview. <i>International Psychogeriatrics</i> , 1991, 3, 177-179.	0.6	1
93	Electroencephalographic Indices for Diagnosis of Delirium. <i>International Psychogeriatrics</i> , 1991, 3, 249-251.	0.6	4
94	Cerebrospinal fluid somatostatin in delirium. II Changes at the acute stage and at one year follow-up. <i>Psychological Medicine</i> , 1990, 20, 501-505.	2.7	12
95	Epileptic Attack, Delirium, and a Creutzfeldt-Jakob-Like Syndrome during Mianserin Treatment. <i>Neuropsychobiology</i> , 1990, 23, 164-168.	0.9	11
96	Decreased somatostatin-like immunoreactivity in the cerebrospinal fluid of chronic schizophrenic patients with cognitive impairment. <i>Psychiatry Research</i> , 1990, 33, 307-312.	1.7	25
97	CSF beta-endorphin-like immunoreactivity in delirium. <i>Biological Psychiatry</i> , 1989, 25, 938-944.	0.7	15
98	Cerebrospinal fluid somatostatin in delirium. <i>Psychological Medicine</i> , 1989, 19, 605-609.	2.7	16
99	Benefits and risks of off label use of antipsychotics in insomnia and anxiety – APSY Oulu project. <i>Nordic Journal of Psychiatry</i> , 0, , 1-1.	0.7	0