David C Henshall

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/592401/publications.pdf

Version: 2024-02-01

201 papers

10,850 citations

28190 55 h-index 40881

g-index

210 all docs

210 docs citations

times ranked

210

10494 citing authors

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Increased hippocampal neurogenesis in Alzheimer's disease. Proceedings of the National Academy of Sciences of the United States of America, 2004, 101, 343-347. | 3.3 | 932 |
| 2 | Silencing microRNA-134 produces neuroprotective and prolonged seizure-suppressive effects. Nature Medicine, 2012, 18, 1087-1094. | 15.2 | 423 |
| 3 | To die or not to die for neurons in ischemia, traumatic brain injury and epilepsy: a review on the stress-activated signaling pathways and apoptotic pathways. Progress in Neurobiology, 2003, 69, 103-142. | 2.8 | 272 |
| 4 | Endotoxin Preconditioning Prevents Cellular Inflammatory Response During Ischemic Neuroprotection in Mice. Stroke, 2004, 35, 2576-2581. | 1.0 | 225 |
| 5 | Neuroprotective Actions of FK506 in Experimental Stroke: <i>In Vivo</i> Evidence against an Antiexcitotoxic Mechanism. Journal of Neuroscience, 1997, 17, 6939-6946. | 1.7 | 203 |
| 6 | MicroRNAs in epilepsy: pathophysiology and clinical utility. Lancet Neurology, The, 2016, 15, 1368-1376. | 4.9 | 200 |
| 7 | CREB-Mediated Bcl-2 Protein Expression after Ischemic Preconditioning. Journal of Cerebral Blood Flow and Metabolism, 2005, 25, 234-246. | 2.4 | 198 |
| 8 | Epilepsy and Apoptosis Pathways. Journal of Cerebral Blood Flow and Metabolism, 2005, 25, 1557-1572. | 2.4 | 196 |
| 9 | miRNA Expression Profile after Status Epilepticus and Hippocampal Neuroprotection by Targeting miR-132. American Journal of Pathology, 2011, 179, 2519-2532. | 1.9 | 194 |
| 10 | Activation of Bcl-2-Associated Death Protein and Counter-Response of Akt within Cell Populations during Seizure-Induced Neuronal Death. Journal of Neuroscience, 2002, 22, 8458-8465. | 1.7 | 176 |
| 11 | Seizure suppression and neuroprotection by targeting the purinergic P2X7 receptor during status epilepticus in mice. FASEB Journal, 2012, 26, 1616-1628. | 0.2 | 173 |
| 12 | Endotoxin Preconditioning Protects against the Cytotoxic Effects of TNFα after Stroke: A Novel Role for TNFα in LPS-Ischemic Tolerance. Journal of Cerebral Blood Flow and Metabolism, 2007, 27, 1663-1674. | 2.4 | 142 |
| 13 | Differential DNA methylation profiles of coding and non-coding genes define hippocampal sclerosis in human temporal lobe epilepsy. Brain, 2015, 138, 616-631. | 3.7 | 140 |
| 14 | Unilateral hippocampal CA3-predominant damage and short latency epileptogenesis after intra-amygdala microinjection of kainic acid in mice. Brain Research, 2008, 1213, 140-151. | 1.1 | 137 |
| 15 | Neuroinflammatory targets and treatments for epilepsy validated in experimental models. Epilepsia, 2017, 58, 27-38. | 2.6 | 131 |
| 16 | Increased neocortical expression of the <scp>P</scp> 2X7 receptor after status epilepticus and anticonvulsant effect of <scp>P</scp> 2X7 receptor antagonist <scp>A</scp> â€438079. Epilepsia, 2013, 54, 1551-1561. | 2.6 | 130 |
| 17 | Involvement of Caspaseâ€3â€Like Protease in the Mechanism of Cell Death Following Focally Evoked Limbic Seizures. Journal of Neurochemistry, 2000, 74, 1215-1223. | 2.1 | 127 |
| 18 | Transient P2X7 Receptor Antagonism Produces Lasting Reductions in Spontaneous Seizures and Gliosis in Experimental Temporal Lobe Epilepsy. Journal of Neuroscience, 2016, 36, 5920-5932. | 1.7 | 127 |

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 19 | Reduced Mature MicroRNA Levels in Association with Dicer Loss in Human Temporal Lobe Epilepsy with Hippocampal Sclerosis. PLoS ONE, 2012, 7, e35921. | 1.1 | 121 |
| 20 | Induction of Oxidative DNA Damage in the Peri-Infarct Region After Permanent Focal Cerebral Ischemia. Journal of Neurochemistry, 2002, 75, 1716-1728. | 2.1 | 120 |
| 21 | MicroRNA and epilepsy. Current Opinion in Neurology, 2014, 27, 199-205. | 1.8 | 109 |
| 22 | LifeTime and improving European healthcare through cell-based interceptive medicine. Nature, 2020, 587, 377-386. | 13.7 | 108 |
| 23 | Cleavage of Bid May Amplify Caspase-8-Induced Neuronal Death Following Focally Evoked Limbic Seizures. Neurobiology of Disease, 2001, 8, 568-580. | 2.1 | 105 |
| 24 | Activation of Poly(ADP-Ribose) Polymerase in the Rat Hippocampus May Contribute to Cellular Recovery Following Sublethal Transient Global Ischemia. Journal of Neurochemistry, 2002, 74, 1636-1645. | 2.1 | 103 |
| 25 | Epilepsy and microRNA. Neuroscience, 2013, 238, 218-229. | 1.1 | 103 |
| 26 | Differential DNA Methylation Patterns Define Status Epilepticus and Epileptic Tolerance. Journal of Neuroscience, 2012, 32, 1577-1588. | 1.7 | 102 |
| 27 | Antagomirs targeting microRNA-134 increase hippocampal pyramidal neuron spine volume in vivo and protect against pilocarpine-induced status epilepticus. Brain Structure and Function, 2015, 220, 2387-2399. | 1.2 | 101 |
| 28 | microRNA targeting of the P2X7 purinoceptor opposes a contralateral epileptogenic focus in the hippocampus. Scientific Reports, 2015, 5, 17486. | 1.6 | 98 |
| 29 | CHOP regulates the p53–MDM2 axis and is required for neuronal survival after seizures. Brain, 2013, 136, 577-592. | 3.7 | 95 |
| 30 | Cerebrospinal fluid microRNAs are potential biomarkers of temporal lobe epilepsy and status epilepticus. Scientific Reports, 2017, 7, 3328. | 1.6 | 93 |
| 31 | MicroRNAs as regulators of brain function and targets for treatment of epilepsy. Nature Reviews Neurology, 2020, 16, 506-519. | 4.9 | 92 |
| 32 | The Epigenetics of Epilepsy and Its Progression. Neuroscientist, 2018, 24, 186-200. | 2.6 | 91 |
| 33 | Formation of a tumour necrosis factor receptor 1 molecular scaffolding complex and activation of apoptosis signal-regulating kinase 1 during seizure-induced neuronal death. European Journal of Neuroscience, 2003, 17 , 2065-2076. | 1.2 | 88 |
| 34 | Dual-center, dual-platform microRNA profiling identifies potential plasma biomarkers of adult temporal lobe epilepsy. EBioMedicine, 2018, 38, 127-141. | 2.7 | 88 |
| 35 | ATPergic signalling during seizures and epilepsy. Neuropharmacology, 2016, 104, 140-153. | 2.0 | 86 |
| 36 | A microRNAâ€129â€5p/Rbfox crosstalk coordinates homeostatic downscaling of excitatory synapses. EMBO Journal, 2017, 36, 1770-1787. | 3.5 | 85 |

| # | Article | IF | Citations |
|----|--|-----|-----------|
| 37 | Expression profiling the microRNA response to epileptic preconditioning identifies miR-184 as a modulator of seizure-induced neuronal death. Experimental Neurology, 2012, 237, 346-354. | 2.0 | 81 |
| 38 | Bim regulation may determine hippocampal vulnerability after injurious seizures and in temporal lobe epilepsy. Journal of Clinical Investigation, 2004, 113, 1059-1068. | 3.9 | 78 |
| 39 | Characterization of neuronal death induced by focally evoked limbic seizures in the C57BL/6 mouse. Journal of Neuroscience Research, 2002, 69, 614-621. | 1.3 | 77 |
| 40 | Endoplasmic Reticulum Stress and Apoptosis Signaling in Human Temporal Lobe Epilepsy. Journal of Neuropathology and Experimental Neurology, 2006, 65, 217-225. | 0.9 | 72 |
| 41 | MicroRNA-Mediated Downregulation of the Potassium Channel Kv4.2 Contributes to Seizure Onset. Cell Reports, 2016, 17, 37-45. | 2.9 | 71 |
| 42 | Elevation of plasma tRNA fragments precedes seizures in human epilepsy. Journal of Clinical Investigation, 2019, 129, 2946-2951. | 3.9 | 71 |
| 43 | Expression of deathâ€associated protein kinase and recruitment to the tumor necrosis factor signaling pathway following brief seizures. Journal of Neurochemistry, 2003, 86, 1260-1270. | 2.1 | 68 |
| 44 | Hippocampal transcriptome after status epilepticus in mice rendered seizure damage-tolerant by epileptic preconditioning features suppressed calcium and neuronal excitability pathways. Neurobiology of Disease, 2008, 32, 442-453. | 2.1 | 68 |
| 45 | Cell Signaling Underlying Epileptic Behavior. Frontiers in Behavioral Neuroscience, 2011, 5, 45. | 1.0 | 68 |
| 46 | Epigenetics and Epilepsy. Cold Spring Harbor Perspectives in Medicine, 2015, 5, a022731. | 2.9 | 68 |
| 47 | Reduced hippocampal damage and epileptic seizures after <i>status epilepticus</i> in mice lacking proapoptotic Puma. FASEB Journal, 2010, 24, 853-861. | 0.2 | 65 |
| 48 | Formation of the Base Modification 8-Hydroxyl-2′ - Deoxyguanosine and DNA Fragmentation Following Seizures Induced by Systemic Kainic Acid in the Rat. Journal of Neurochemistry, 2001, 74, 302-309. | 2.1 | 63 |
| 49 | Convulsant Doses of a Dopamine D1 Receptor Agonist Result in Erk-Dependent Increases in Zif268 and Arc/Arg3.1 Expression in Mouse Dentate Gyrus. PLoS ONE, 2011, 6, e19415. | 1.1 | 63 |
| 50 | Proteins and microRNAs are differentially expressed in tear fluid from patients with Alzheimer's disease. Scientific Reports, 2019, 9, 15437. | 1.6 | 63 |
| 51 | Potent Anti-seizure Effects of Locked Nucleic Acid Antagomirs Targeting miR-134 in Multiple Mouse and Rat Models of Epilepsy. Molecular Therapy - Nucleic Acids, 2017, 6, 45-56. | 2.3 | 62 |
| 52 | Bim regulation may determine hippocampal vulnerability after injurious seizures and in temporal lobe epilepsy. Journal of Clinical Investigation, 2004, 113, 1059-1068. | 3.9 | 62 |
| 53 | <i>In vivo</i> Contributions of BH3-Only Proteins to Neuronal Death Following Seizures, Ischemia, and Traumatic Brain Injury. Journal of Cerebral Blood Flow and Metabolism, 2011, 31, 1196-1210. | 2.4 | 61 |
| 54 | lgG Leakage May Contribute to Neuronal Dysfunction in Drug-Refractory Epilepsies With Blood-Brain Barrier Disruption. Journal of Neuropathology and Experimental Neurology, 2012, 71, 826-838. | 0.9 | 60 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | P2X7 Receptor Inhibition Interrupts the Progression of Seizures in Immature Rats and Reduces Hippocampal Damage. CNS Neuroscience and Therapeutics, 2014, 20, 556-564. | 1.9 | 58 |
| 56 | Critical Evaluation of P2X7 Receptor Antagonists in Selected Seizure Models. PLoS ONE, 2016, 11, e0156468. | 1.1 | 57 |
| 57 | Increased Expression of MicroRNA-29a in ALS Mice: Functional Analysis of Its Inhibition. Journal of Molecular Neuroscience, 2014, 53, 231-241. | 1.1 | 56 |
| 58 | MicroRNAs in the pathophysiology and treatment of status epilepticus. Frontiers in Molecular Neuroscience, 2013, 6, 37. | 1.4 | 55 |
| 59 | Contribution of apoptosis-associated signaling pathways to epileptogenesis: lessons from Bcl-2 family knockouts. Frontiers in Cellular Neuroscience, 2013, 7, 110. | 1.8 | 54 |
| 60 | Neurodevelopmental alterations and seizures developed by mouse model of infantile hypophosphatasia are associated with purinergic signalling deregulation. Human Molecular Genetics, 2016, 25, 4143-4156. | 1.4 | 54 |
| 61 | Identification of clinically relevant biomarkers of epileptogenesis — a strategic roadmap. Nature Reviews Neurology, 2021, 17, 231-242. | 4.9 | 54 |
| 62 | Apoptosis, Bcl-2 family proteins and caspases: the ABCs of seizure-damage and epileptogenesis?. International Journal of Physiology, Pathophysiology and Pharmacology, 2009, 1, 97-115. | 0.8 | 54 |
| 63 | "TORNADO―– Theranostic One-Step RNA Detector; microfluidic disc for the direct detection of microRNA-134 in plasma and cerebrospinal fluid. Scientific Reports, 2017, 7, 1750. | 1.6 | 53 |
| 64 | Bcl-w Protects Hippocampus during Experimental Status Epilepticus. American Journal of Pathology, 2007, 171, 1258-1268. | 1.9 | 52 |
| 65 | microRNA and Epilepsy. Advances in Experimental Medicine and Biology, 2015, 888, 41-70. | 0.8 | 52 |
| 66 | Bax Regulates Neuronal Ca ²⁺ Homeostasis. Journal of Neuroscience, 2015, 35, 1706-1722. | 1.7 | 52 |
| 67 | Effects of hypoxia-induced neonatal seizures on acute hippocampal injury and later-life seizure susceptibility and anxiety-related behavior in mice. Neurobiology of Disease, 2015, 83, 100-114. | 2.1 | 52 |
| 68 | Expression and function of the metabotropic purinergic P2Y receptor family in experimental seizure models and patients with drugâ€refractory epilepsy. Epilepsia, 2017, 58, 1603-1614. | 2.6 | 51 |
| 69 | Modulators of neuronal cell death in epilepsy. Current Opinion in Pharmacology, 2008, 8, 75-81. | 1.7 | 50 |
| 70 | Protective neuronal induction of ATF5 in endoplasmic reticulum stress induced by status epilepticus. Brain, 2013, 136, 1161-1176. | 3.7 | 49 |
| 71 | EpimiRBase: a comprehensive database of microRNA-epilepsy associations. Bioinformatics, 2016, 32, 1436-1438. | 1.8 | 48 |
| 72 | Death-associated protein kinase expression in human temporal lobe epilepsy. Annals of Neurology, 2004, 55, 485-494. | 2.8 | 47 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | NMDA receptorâ€mediated excitotoxic neuronal apoptosis <i>in vitro</i> and <i>in vivo</i> occurs in an ER stress and PUMA independent manner. Journal of Neurochemistry, 2008, 105, 891-903. | 2.1 | 47 |
| 74 | Identification of a Novel Bcl-2-interacting Mediator of Cell Death (Bim) E3 Ligase, Tripartite Motif-containing Protein 2 (TRIM2), and Its Role in Rapid Ischemic Tolerance-induced Neuroprotection. Journal of Biological Chemistry, 2011, 286, 19331-19339. | 1.6 | 47 |
| 75 | Bok Is Not Pro-Apoptotic But Suppresses Poly ADP-Ribose Polymerase-Dependent Cell Death Pathways and Protects against Excitotoxic and Seizure-Induced Neuronal Injury. Journal of Neuroscience, 2016, 36, 4564-4578. | 1.7 | 47 |
| 76 | Involvement of micro <scp>RNA</scp> s in epileptogenesis. Epilepsia, 2016, 57, 1015-1026. | 2.6 | 47 |
| 77 | Microvascular stabilization via blood-brainÂbarrier regulation prevents seizure activity. Nature Communications, 2022, 13, 2003. | 5.8 | 47 |
| 78 | Development of a model of seizure-induced hippocampal injury with features of programmed cell death in the BALB/c mouse. Journal of Neuroscience Research, 2004, 76, 121-128. | 1.3 | 46 |
| 79 | microRNAs in the pathophysiology of epilepsy. Neuroscience Letters, 2018, 667, 47-52. | 1.0 | 46 |
| 80 | Upregulation of Mitochondrial Base-Excision Repair Capability within Rat Brain after Brief Ischemia. Journal of Cerebral Blood Flow and Metabolism, 2003, 23, 88-98. | 2.4 | 45 |
| 81 | Microarray profile of seizure damage-refractory hippocampal CA3 in a mouse model of epileptic preconditioning. Neuroscience, 2007, 150, 467-477. | 1.1 | 45 |
| 82 | Dopamine D1 vs D5 receptor-dependent induction of seizures in relation to DARPP-32, ERK1/2 and GluR1-AMPA signalling. Neuropharmacology, 2008, 54, $1051-1061$. | 2.0 | 45 |
| 83 | P2X receptors as targets for the treatment of status epilepticus. Frontiers in Cellular Neuroscience, 2013, 7, 237. | 1.8 | 45 |
| 84 | Effects of P2X7 receptor antagonists on hypoxia-induced neonatal seizures in mice. Neuropharmacology, 2017, 116, 351-363. | 2.0 | 44 |
| 85 | Transgenic Overexpression of 14-3-3 Zeta Protects Hippocampus against Endoplasmic Reticulum Stress and Status Epilepticus In Vivo. PLoS ONE, 2013, 8, e54491. | 1.1 | 44 |
| 86 | Caspase-3 Cleavage and Nuclear Localization of Caspase-Activated DNase in Human Temporal Lobe Epilepsy. Journal of Cerebral Blood Flow and Metabolism, 2006, 26, 583-589. | 2.4 | 43 |
| 87 | Isoform- and subcellular fraction-specific differences in hippocampal 14-3-3 levels following experimentally evoked seizures and in human temporal lobe epilepsy. Journal of Neurochemistry, 2006, 99, 561-569. | 2.1 | 42 |
| 88 | P2X purinoceptors as a link between hyperexcitability and neuroinflammation in status epilepticus. Epilepsy and Behavior, 2015, 49, 8-12. | 0.9 | 42 |
| 89 | Elevated Plasma microRNA-206 Levels Predict Cognitive Decline and Progression to Dementia from Mild Cognitive Impairment. Biomolecules, 2019, 9, 734. | 1.8 | 41 |
| 90 | A systems approach delivers a functional microRNA catalog and expanded targets for seizure suppression in temporal lobe epilepsy. Proceedings of the National Academy of Sciences of the United States of America, 2020, 117, 15977-15988. | 3.3 | 41 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 91 | Caspase-2 activation is redundant during seizure-induced neuronal death. Journal of Neurochemistry, 2001, 77, 886-895. | 2.1 | 40 |
| 92 | Experimental Neonatal Status Epilepticus and the Development of Temporal Lobe Epilepsy with Unilateral Hippocampal Sclerosis. American Journal of Pathology, 2010, 176, 330-342. | 1.9 | 40 |
| 93 | Mutation of Semaphorin-6A Disrupts Limbic and Cortical Connectivity and Models Neurodevelopmental Psychopathology. PLoS ONE, 2011, 6, e26488. | 1.1 | 40 |
| 94 | Bcl-w expression is increased in brain regions affected by focal cerebral ischemia in the rat. Neuroscience Letters, 2000, 279, 193-195. | 1.0 | 39 |
| 95 | Evidence of tumor necrosis factor receptor 1 signaling in human temporal lobe epilepsy. Experimental Neurology, 2006, 202, 410-420. | 2.0 | 39 |
| 96 | Precise Targeting of miRNA Sites Restores CFTR Activity in CF Bronchial Epithelial Cells. Molecular Therapy, 2020, 28, 1190-1199. | 3.7 | 39 |
| 97 | Opportunities and challenges for microRNA-targeting therapeutics for epilepsy. Trends in Pharmacological Sciences, 2021, 42, 605-616. | 4.0 | 39 |
| 98 | Depletion of 14â€3â€3 zeta elicits endoplasmic reticulum stress and cell death, and increases vulnerability to kainateâ€induced injury in mouse hippocampal cultures. Journal of Neurochemistry, 2008, 106, 978-988. | 2.1 | 38 |
| 99 | Context-Specific Switch from Anti- to Pro-epileptogenic Function of the P2Y ₁ Receptor in Experimental Epilepsy. Journal of Neuroscience, 2019, 39, 5377-5392. | 1.7 | 37 |
| 100 | P2X7 receptor in epilepsy; role in pathophysiology and potential targeting for seizure control. International Journal of Physiology, Pathophysiology and Pharmacology, 2012, 4, 174-87. | 0.8 | 36 |
| 101 | Spatio-temporally restricted blood–brain barrier disruption after intra-amygdala kainic acid-induced status epilepticus in mice. Epilepsy Research, 2013, 103, 167-179. | 0.8 | 35 |
| 102 | Kainic Acid-Induced Seizures Modulate Akt (SER473) Phosphorylation in the Hippocampus of Dopamine D2 Receptor Knockout Mice. Journal of Molecular Neuroscience, 2013, 49, 202-210. | 1.1 | 35 |
| 103 | Genome-wide microRNA profiling of plasma from three different animal models identifies biomarkers of temporal lobe epilepsy. Neurobiology of Disease, 2020, 144, 105048. | 2.1 | 35 |
| 104 | Subcellular distribution of Bcl-2 family proteins and 14-3-3 within the hippocampus during seizure-induced neuronal death in the rat. Neuroscience Letters, 2004, 356, 163-166. | 1.0 | 34 |
| 105 | Elevated p53 and lower MDM2 expression in hippocampus from patients with intractable temporal lobe epilepsy. Epilepsy Research, 2007, 77, 151-156. | 0.8 | 34 |
| 106 | Targeting microRNA-134 for seizure control and disease modification in epilepsy. EBioMedicine, 2019, 45, 646-654. | 2.7 | 34 |
| 107 | Spatio-temporal profile of DNA fragmentation and its relationship to patterns of epileptiform activity following focally evoked limbic seizures. Brain Research, 2000, 858, 290-302. | 1.1 | 33 |
| 108 | miRNA-Mediated Regulation of Adult Hippocampal Neurogenesis; Implications for Epilepsy. Brain Plasticity, 2017, 3, 43-59. | 1.9 | 33 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Bi-directional genetic modulation of GSK-3 \hat{l}^2 exacerbates hippocampal neuropathology in experimental status epilepticus. Cell Death and Disease, 2018, 9, 969. | 2.7 | 32 |
| 110 | Discovery and validation of blood micro <scp>RNA</scp> s as molecular biomarkers of epilepsy: Ways to close current knowledge gaps. Epilepsia Open, 2018, 3, 427-436. | 1.3 | 32 |
| 111 | A calcium-sensitive feed-forward loop regulating the expression of the ATP-gated purinergic P2X7 receptor via specificity protein 1 and microRNA-22. Biochimica Et Biophysica Acta - Molecular Cell Research, 2017, 1864, 255-266. | 1.9 | 31 |
| 112 | Expression and Differential Processing of Caspases 6 and 7 in Relation to Specific Epileptiform EEG Patterns Following Limbic Seizures. Neurobiology of Disease, 2002, 10, 71-87. | 2.1 | 29 |
| 113 | Relationship Between Seizure-Induced Transcription of the DNA Damage-Inducible Gene GADD45, DNA Fragmentation, and Neuronal Death in Focally Evoked Limbic Epilepsy. Journal of Neurochemistry, 2002, 73, 1573-1583. | 2.1 | 29 |
| 114 | Expression, interaction, and proteolysis of death-associated protein kinase and p53 within vulnerable and resistant hippocampal subfields following seizures. Hippocampus, 2004, 14, 326-336. | 0.9 | 29 |
| 115 | Regulatory Mechanisms of the RNA Modification m6A and Significance in Brain Function in Health and Disease. Frontiers in Cellular Neuroscience, 2021, 15, 671932. | 1.8 | 29 |
| 116 | High Throughput qPCR Expression Profiling of Circulating MicroRNAs Reveals Minimal Sex- and Sample Timing-Related Variation in Plasma of Healthy Volunteers. PLoS ONE, 2015, 10, e0145316. | 1.1 | 29 |
| 117 | Complex spectrum of phenobarbital effects in a mouse model of neonatal hypoxia-induced seizures. Scientific Reports, 2018, 8, 9986. | 1.6 | 28 |
| 118 | Antagonizing Increased <i>miR-135a</i> Levels at the Chronic Stage of Experimental TLE Reduces Spontaneous Recurrent Seizures. Journal of Neuroscience, 2019, 39, 5064-5079. | 1.7 | 28 |
| 119 | Altered Biogenesis and MicroRNA Content of Hippocampal Exosomes Following Experimental Status Epilepticus. Frontiers in Neuroscience, 2019, 13, 1404. | 1.4 | 27 |
| 120 | MicroRNA-22 Controls Aberrant Neurogenesis and Changes in Neuronal Morphology After Status Epilepticus. Frontiers in Molecular Neuroscience, 2018, 11, 442. | 1.4 | 26 |
| 121 | Hippocampal damage after intra-amygdala kainic acid-induced status epilepticus and seizure preconditioning-mediated neuroprotection in SJL mice. Epilepsy Research, 2010, 88, 151-161. | 0.8 | 24 |
| 122 | BH3â€only protein Bid is dispensable for seizureâ€induced neuronal death and the associated nuclear accumulation of apoptosisâ€inducing factor. Journal of Neurochemistry, 2010, 115, 92-101. | 2.1 | 24 |
| 123 | Bi-lateral changes to hippocampal cholesterol levels during epileptogenesis and in chronic epilepsy following focal-onset status epilepticus in mice. Brain Research, 2012, 1480, 81-90. | 1.1 | 23 |
| 124 | Antagomirs and micro <scp>RNA</scp> in status epilepticus. Epilepsia, 2013, 54, 17-19. | 2.6 | 23 |
| 125 | Spatiotemporal progression of ubiquitin-proteasome system inhibition after status epilepticus suggests protective adaptation against hippocampal injury. Molecular Neurodegeneration, 2017, 12, 21. | 4.4 | 23 |
| 126 | Circulating P2X7 Receptor Signaling Components as Diagnostic Biomarkers for Temporal Lobe Epilepsy. Cells, 2021, 10, 2444. | 1.8 | 23 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Seizure preconditioning and epileptic tolerance: models and mechanisms. International Journal of Physiology, Pathophysiology and Pharmacology, 2009, 1, 180-191. | 0.8 | 23 |
| 128 | Interaction of 14-3-3 with Bid during seizure-induced neuronal death. Journal of Neurochemistry, 2004, 86, 460-469. | 2.1 | 22 |
| 129 | Mitochondrial localization of the Forkhead box class O transcription factor <scp>FOXO</scp> 3a in brain. Journal of Neurochemistry, 2013, 124, 749-756. | 2.1 | 21 |
| 130 | Elevated blood purine levels as a biomarker of seizures and epilepsy. Epilepsia, 2021, 62, 817-828. | 2.6 | 21 |
| 131 | Epigenetic principles underlying epileptogenesis and epilepsy syndromes. Neurobiology of Disease, 2021, 148, 105179. | 2.1 | 20 |
| 132 | Systemic delivery of antagomirs during blood-brain barrier disruption is disease-modifying in experimental epilepsy. Molecular Therapy, 2021, 29, 2041-2052. | 3.7 | 20 |
| 133 | CHD2-Related CNS Pathologies. International Journal of Molecular Sciences, 2021, 22, 588. | 1.8 | 20 |
| 134 | Increased expression of the ATPâ€gated P2X7 receptor reduces responsiveness to antiâ€convulsants during status epilepticus in mice. British Journal of Pharmacology, 2022, 179, 2986-3006. | 2.7 | 20 |
| 135 | Expression, proteolysis and activation of caspases 6 and 7 during rat C6 glioma cell apoptosis. Neuroscience Letters, 2002, 324, 33-36. | 1.0 | 19 |
| 136 | Bclâ€⊋ homology domain 3â€only proteins Puma and Bim mediate the vulnerability of CA1 hippocampal neurons to proteasome inhibition <i>inâ€fvivo</i> . European Journal of Neuroscience, 2011, 33, 401-408. | 1.2 | 19 |
| 137 | Genetic deletion of microRNA-22 blunts the inflammatory transcriptional response to status epilepticus and exacerbates epilepsy in mice. Molecular Brain, 2020, 13, 114. | 1.3 | 18 |
| 138 | Increased Bcl-w expression following focally evoked limbic seizures in the rat. Neuroscience Letters, 2001, 305, 153-156. | 1.0 | 17 |
| 139 | Dysregulation of Specialized Delay/Interference-Dependent Working Memory Following Loss of Dysbindin-1A in Schizophrenia-Related Phenotypes. Neuropsychopharmacology, 2017, 42, 1349-1360. | 2.8 | 17 |
| 140 | Manipulating MicroRNAs in Murine Models: Targeting the Multi-Targeting in Epilepsy. Epilepsy Currents, 2017, 17, 43-47. | 0.4 | 17 |
| 141 | Spared <scp>CA</scp> 1 pyramidal neuron function and hippocampal performance following antisense knockdown of micro <scp>RNA</scp> â€134. Epilepsia, 2018, 59, 1518-1526. | 2.6 | 17 |
| 142 | Advancing research toward faster diagnosis, better treatment, and end of stigma in epilepsy. Epilepsia, 2019, 60, 1281-1292. | 2.6 | 17 |
| 143 | High concordance between hippocampal transcriptome of the mouse intraâ€amygdala kainic acid model and human temporal lobe epilepsy. Epilepsia, 2020, 61, 2795-2810. | 2.6 | 17 |
| 144 | Epigenetics explained: a topic "primer―for the epilepsy community by the ILAE Genetics/Epigenetics Task Force. Epileptic Disorders, 2020, 22, 127-141. | 0.7 | 17 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 145 | Effects of transient focal cerebral ischemia in mice deficient in puma. Neuroscience Letters, 2009, 451, 237-240. | 1.0 | 16 |
| 146 | Elevated serum Bcl-2 in children with temporal lobe epilepsy. Seizure: the Journal of the British Epilepsy Association, 2012, 21, 250-253. | 0.9 | 16 |
| 147 | Hsp27 binding to the 3′UTR of∢i>bim⟨/i>mRNA prevents neuronal death during oxidative stress–induced injury: a novel cytoprotective mechanism. Molecular Biology of the Cell, 2014, 25, 3413-3423. | 0.9 | 16 |
| 148 | Transcriptional Response of Polycomb Group Genes to Status Epilepticus in Mice is Modified by Prior Exposure to Epileptic Preconditioning. Frontiers in Neurology, 2015, 6, 46. | 1.1 | 16 |
| 149 | RNA sequencing of synaptic and cytoplasmic Upf1-bound transcripts supports contribution of nonsense-mediated decay to epileptogenesis. Scientific Reports, 2017, 7, 41517. | 1.6 | 16 |
| 150 | The Anti-inflammatory Compound Candesartan Cilexetil Improves Neurological Outcomes in a Mouse Model of Neonatal Hypoxia. Frontiers in Immunology, 2019, 10, 1752. | 2.2 | 16 |
| 151 | MicroRNAs as biomarkers and treatment targets in status epilepticus. Epilepsy and Behavior, 2019, 101, 106272. | 0.9 | 16 |
| 152 | Can Genes Modify Stroke Outcome and By What Mechanisms?. Stroke, 2012, 43, 286-291. | 1.0 | 15 |
| 153 | Direct, non-amplified detection of microRNA-134 in plasma from epilepsy patients. RSC Advances, 2015, 5, 90071-90078. | 1.7 | 15 |
| 154 | Comparison of short-term effects of midazolam and lorazepam in the intra-amygdala kainic acid model of status epilepticus in mice. Epilepsy and Behavior, 2015, 51, 191-198. | 0.9 | 15 |
| 155 | Haploinsufficient TNAP Mice Display Decreased Extracellular ATP Levels and Expression of Pannexin-1 Channels. Frontiers in Pharmacology, 2018, 9, 170. | 1.6 | 14 |
| 156 | Epigenetics and noncoding RNA: Recent developments and future therapeutic opportunities. European Journal of Paediatric Neurology, 2020, 24, 30-34. | 0.7 | 14 |
| 157 | Poststatus Epilepticus Models: Focal Kainic Acid. , 2017, , 611-624. | | 13 |
| 158 | Enrichment of Circular RNA Expression Deregulation at the Transition to Recurrent Spontaneous Seizures in Experimental Temporal Lobe Epilepsy. Frontiers in Genetics, 2021, 12, 627907. | 1.1 | 13 |
| 159 | Antagomir-mediated suppression of microRNA-134 reduces kainic acid-induced seizures in immature mice. Scientific Reports, 2021, 11, 340. | 1.6 | 13 |
| 160 | Expression of neurogenesis genes in human temporal lobe epilepsy with hippocampal sclerosis. International Journal of Physiology, Pathophysiology and Pharmacology, 2011, 3, 38-47. | 0.8 | 13 |
| 161 | AntimiR targeting of microRNA-134 reduces seizures in a mouse model of Angelman syndrome. Molecular Therapy - Nucleic Acids, 2022, 28, 514-529. | 2.3 | 13 |
| 162 | Life-span characterization of epilepsy and comorbidities in Dravet syndrome mice carrying a targeted deletion of exon 1 of the Scn1a gene. Experimental Neurology, 2022, 354, 114090. | 2.0 | 13 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 163 | Detection of 14-3-3î¶ in cerebrospinal fluid following experimentally evoked seizures. Biomarkers, 2008, 13, 377-384. | 0.9 | 12 |
| 164 | Quantification of tRNA fragments by electrochemical direct detection in small volume biofluid samples. Scientific Reports, 2020, 10, 7516. | 1.6 | 12 |
| 165 | Temporally Altered miRNA Expression in a Piglet Model of Hypoxic Ischemic Brain Injury. Molecular Neurobiology, 2020, 57, 4322-4344. | 1.9 | 12 |
| 166 | Detection of spontaneous seizures in EEGs in multiple experimental mouse models of epilepsy. Journal of Neural Engineering, 2021, 18, 056060. | 1.8 | 12 |
| 167 | Epigenetic changes in status epilepticus. Epilepsia, 2018, 59, 82-86. | 2.6 | 11 |
| 168 | Polyadenylation of mRNA as a novel regulatory mechanism of gene expression in temporal lobe epilepsy. Brain, 2020, 143, 2139-2153. | 3.7 | 11 |
| 169 | Proteomic analysis of 14-3-3 zeta binding proteins in the mouse hippocampus. International Journal of Physiology, Pathophysiology and Pharmacology, 2012, 4, 74-83. | 0.8 | 11 |
| 170 | Electrical stimulation of the ventral hippocampal commissure delays experimental epilepsy and is associated with altered microRNA expression. Brain Stimulation, 2019, 12, 1390-1401. | 0.7 | 10 |
| 171 | Cell Death and Survival Mechanisms after Single and Repeated Brief Seizures. , 2012, , 362-376. | | 10 |
| 172 | Epilepsy Benchmarks Area III: Improved Treatment Options for Controlling Seizures and Epilepsy-Related Conditions Without Side Effects. Epilepsy Currents, 2020, 20, 23S-30S. | 0.4 | 9 |
| 173 | RNA-sequencing analysis of umbilical cord plasma microRNAs from healthy newborns. PLoS ONE, 2018, 13, e0207952. | 1.1 | 8 |
| 174 | Proteomic Analysis After Status Epilepticus Identifies UCHL1 as Protective Against Hippocampal Injury. Neurochemical Research, 2017, 42, 2033-2054. | 1.6 | 7 |
| 175 | Predictive modelling of hypoxic ischaemic encephalopathy risk following perinatal asphyxia. Heliyon, 2021, 7, e07411. | 1.4 | 7 |
| 176 | Distinct behavioral and epileptic phenotype differences in 129/P mice compared to C57BL/6 mice subject to intraamygdala kainic acid-induced status epilepticus. Epilepsy and Behavior, 2016, 64, 186-194. | 0.9 | 6 |
| 177 | GABA Regulation of Burst Firing in Hippocampal Astrocyte Neural Circuit: A Biophysical Model. Frontiers in Cellular Neuroscience, 2019, 13, 335. | 1.8 | 6 |
| 178 | P2X7 Receptor-Dependent microRNA Expression Profile in the Brain Following Status Epilepticus in Mice. Frontiers in Molecular Neuroscience, 2020, 13, 127. | 1.4 | 6 |
| 179 | Tubby-like protein 1 (Tulp1) is a target of microRNA-134 and is down-regulated in experimental epilepsy. International Journal of Physiology, Pathophysiology and Pharmacology, 2017, 9, 178-187. | 0.8 | 6 |
| 180 | Electroencephalographic and behavioral convulsant effects of hydrobromide and hydrochloride salts of bupropion in conscious rodents. Neuropsychiatric Disease and Treatment, 2009, 5, 189. | 1.0 | 5 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 181 | Suppression of TNF receptor-1 signaling in an in vitro model of epileptic tolerance. International Journal of Physiology, Pathophysiology and Pharmacology, 2011, 3, 120-32. | 0.8 | 5 |
| 182 | <scp>MicroRNA</scp> inhibition using <scp>antimiRs</scp> in acute human brain tissue sections. Epilepsia, 2022, 63, . | 2.6 | 5 |
| 183 | Bilateral intranigral NMDA infusion suppresses neuronal injury without affecting the duration of kainic acid-induced seizures in rats. Neuroscience Letters, 1998, 251, 69-71. | 1.0 | 4 |
| 184 | Overexpression of 14-3-3ζ Increases Brain Levels of C/EBP Homologous Protein CHOP. Journal of Molecular Neuroscience, 2015, 56, 255-262. | 1.1 | 4 |
| 185 | Generation of six induced pluripotent stem cell (iPSC) lines from two patients with amyotrophic lateral sclerosis (NUIGi043-A, NUIGi043-B, NUIGi043-C, NUIGi044-A, NUIGi044-B, NUIGi044-C). Stem Cell Research, 2019, 40, 101558. | 0.3 | 4 |
| 186 | Systemic delivery of selective EP1 and EP3 receptor antagonists attenuates pentylenetetrazole-induced seizures in mice. International Journal of Physiology, Pathophysiology and Pharmacology, 2018, 10, 47-59. | 0.8 | 4 |
| 187 | Targeting the proteasome in epilepsy. Oncotarget, 2017, 8, 45042-45043. | 0.8 | 3 |
| 188 | Investigating Gene Promoter Methylation in a Mouse Model of Status Epilepticus. Methods in Molecular Biology, 2013, 1067, 87-101. | 0.4 | 2 |
| 189 | Detection of MicroRNAs in Brain Slices Using In Situ Hybridization. Methods in Molecular Biology, 2017, 1509, 85-91. | 0.4 | 2 |
| 190 | Deletion of the BH3-only protein Noxa alters electrographic seizures but does not protect against hippocampal damage after status epilepticus in mice. Cell Death and Disease, 2018, 8, e2556-e2556. | 2.7 | 2 |
| 191 | Generation of twelve induced pluripotent stem cell lines from two healthy controls and two patients with sporadic amyotrophic lateral sclerosis. Stem Cell Research, 2020, 44, 101752. | 0.3 | 2 |
| 192 | Profiling of Argonaute-2-loaded microRNAs in a mouse model of frontotemporal dementia with parkinsonism-17. International Journal of Physiology, Pathophysiology and Pharmacology, 2018, 10, 172-183. | 0.8 | 2 |
| 193 | Neurogenic function in rats with unilateral hippocampal sclerosis that experienced early-life status epilepticus. International Journal of Physiology, Pathophysiology and Pharmacology, 2014, 6, 199-208. | 0.8 | 1 |
| 194 | BICS01 Mediates Reversible Anti-seizure Effects in Brain Slice Models of Epilepsy. Frontiers in Neurology, 2021, 12, 791608. | 1.1 | 1 |
| 195 | Meeting report: EpiXchange II brings together European epilepsy research projects to discuss latest advances. Epilepsy Research, 2021, 178, 106811. | 0.8 | 1 |
| 196 | Electrochemiluminescent detection of epilepsy biomarker miR-134 using a metal complex light switch. Bioelectrochemistry, 2022, 146, 108150. | 2.4 | 1 |
| 197 | Focally Applied Chemoconvulsants. , 2017, , 513-527. | | 0 |
| 198 | Could miR-134 be a marker of ionizing radiation toxicity?. Non-coding RNA Investigation, 2018, 2, 24-24. | 0.6 | 0 |

DAVIDÂC HENSHALL

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 199 | Brain delivery of a virus to block seizures helps mice get a silent NACHT. EBioMedicine, 2019, 47, 8-9. | 2.7 | O |
| 200 | Building a supportive framework for brain research in Ireland: Inaugural position paper of the Irish Brain Council. European Journal of Neuroscience, 2019, 49, 1362-1370. | 1.2 | 0 |
| 201 | Preconditioning for Epilepsy. , 2013, , 521-539. | | O |