Maria Elida Pirovani

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5926687/publications.pdf

Version: 2024-02-01

25 papers 574 citations

15 h-index 610482 24 g-index

27 all docs

 $\begin{array}{c} 27 \\ \text{docs citations} \end{array}$

times ranked

27

624 citing authors

#	Article	IF	CITATIONS
1	Intestinal and colonic bioaccessibility of phenolic compounds from fruit smoothies as affected by the thermal processing and the storage conditions. Food Research International, 2022, 155, 111086.	2.9	3
2	Kinetic modeling of the changes in bioactive compounds and quality attributes of fresh-cut strawberries stored in controlled atmospheres with high oxygen alone or with carbon dioxide. Postharvest Biology and Technology, 2022, 190, 111947.	2.9	5
3	Improving freshâ€cut apple quality and healthy potentialâ€related attributes through mild vacuum impregnation process. Journal of Food Processing and Preservation, 2021, 45, e15995.	0.9	3
4	Sous-Vide as a Technique for Preparing Healthy and High-Quality Vegetable and Seafood Products. Foods, 2020, 9, 1537.	1.9	42
5	Effect of enriched O2 and CO2 atmospheres on the overall quality and the bioactive potential of fresh blackberries. Postharvest Biology and Technology, 2020, 164, 111166.	2.9	15
6	Variation of Eugenia uniflora L. fruit quality attributes in the south of the Argentine Gran Chaco. Forests Trees and Livelihoods, 2020, 29, 130-142.	0.5	4
7	Changes in the bioactive properties of strawberries caused by the storage in oxygen―and carbon dioxideâ€enriched atmospheres. Food Science and Nutrition, 2019, 7, 2527-2536.	1.5	9
8	Changes due to high oxygen and high carbon dioxide atmospheres on the general quality and the polyphenolic profile of strawberries. Postharvest Biology and Technology, 2019, 148, 49-57.	2.9	18
9	Spray Washing Disinfection with Peracetic Acid in the Processing of Fresh-Cut Strawberries: An Alternative for Dipping Techniques. International Journal of Fruit Science, 2019, 19, 258-275.	1.2	2
10	Modeling the Impact of the Type of Cutting and Storage Temperature on the Bioactive Compound Content, Phenylpropanoid Metabolism Enzymes and Quality Attributes of Fresh-Cut Strawberries. Food and Bioprocess Technology, 2018, 11, 96-109.	2.6	15
11	Optimization of strawberry disinfection by fogging of a mixture of peracetic acid and hydrogen peroxide based on microbial reduction, color and phytochemicals retention. Food Science and Technology International, 2016, 22, 485-495.	1.1	22
12	Quantitative comparison of phytochemical profile, antioxidant, and anti-inflammatory properties of blackberry fruits adapted to Argentina. Journal of Food Composition and Analysis, 2016, 47, 82-91.	1.9	50
13	Impact of a new postharvest disinfection method based on peracetic acid fogging on the phenolic profile of strawberries. Postharvest Biology and Technology, 2016, 117, 197-205.	2.9	20
14	Kinetic Parameters of Changes in Sensory Characteristics of Minimally Processed Rambutan. International Journal of Fruit Science, 2016, 16, 159-170.	1.2	3
15	Modelling changes in anthocyanins, total vitamin C and colour as a consequence of peracetic acid washing disinfection of two cultivars of strawberries for freshâ€cut processing. International Journal of Food Science and Technology, 2013, 48, 954-961.	1.3	12
16	Bioactive Compounds and Antioxidant Capacity of Camarosa and Selva Strawberries (Fragaria x) Tj ETQq0 0 0 rg	gBT/Qverlo	ock ₈₀ 0 Tf 50 1
17	Modeling changes of sensory attributes for individual and mixed fresh-cut leafy vegetables. Postharvest Biology and Technology, 2005, 38, 202-212.	2.9	41
18	Reduction of chlorine concentration and microbial load during washing-disinfection of shredded lettuce. International Journal of Food Science and Technology, 2004, 39, 341-347.	1.3	47

#	ARTICLE	IF	CITATION
19	FRESH?CUT SPINACH QUALITY AS INFLUENCED BY SPIN DRYING PARAMETERS. Journal of Food Quality, 2003, 26, 231-242.	1.4	4
20	Mesophilic Aerobic Population of Fresh-cut Spinach as Affected by Chemical Treatment and Type of Packaging Film. Journal of Food Science, 2003, 68, 602-606.	1.5	15
21	Sensory Characteristics of Fresh-Cut Spinach Preserved by Combined Factors Methodology. Journal of Food Science, 2002, 67, 1544-1549.	1.5	29
22	Predictive Models for Available Chlorine Depletion and Total Microbial Count Reduction During Washing of Fresh-Cut Spinach. Journal of Food Science, 2001, 66, 860-864.	1.5	32
23	QUALITY OF MINIMALLY PROCESSED LETTUCE AS INFLUENCED BY PACKAGING AND CHEMICAL TREATMENT. Journal of Food Quality, 1998, 21, 475-484.	1.4	30
24	Survival and Growth of Salmonella hadar on Minimally Processed Cabbage as Influenced by Storage Abuse Conditions. Journal of Food Science, 1997, 62, 616-618.	1.5	32
25	STORAGE QUALITY OF MINIMALLY PROCESSED CABBAGE PACKAGED IN PLASTIC FILMS. Journal of Food Quality, 1997, 20, 381-389.	1.4	36