

# Yung-Feng Liao

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/5927097/yung-feng-liao-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

44  
papers

3,700  
citations

18  
h-index

47  
g-index

47  
ext. papers

3,985  
ext. citations

5.5  
avg, IF

3.49  
L-index

#	Paper	IF	Citations
44	Equilibrative nucleoside transporter 1 inhibition rescues energy dysfunction and pathology in a model of tauopathy. <i>Acta Neuropathologica Communications</i> , <b>2021</b> , 9, 112	7.3	2
43	Dual-Specificity Phosphatase 15 (DUSP15) Modulates Notch Signaling by Enhancing the Stability of Notch Protein. <i>Molecular Neurobiology</i> , <b>2021</b> , 58, 2204-2214	6.2	0
42	Piwi reduction in the aged niche eliminates germline stem cells via Toll-GSK3 signaling. <i>Nature Communications</i> , <b>2020</b> , 11, 3147	17.4	8
41	Phosphatidylinositol-4-phosphate 5-kinase type 1 $\beta$ attenuates A $\beta$ production by promoting non-amyloidogenic processing of amyloid precursor protein. <i>FASEB Journal</i> , <b>2020</b> , 34, 12127-12146	0.9	1
40	Novel Endogenous Ligands of Aryl Hydrocarbon Receptor Mediate Neural Development and Differentiation of Neuroblastoma. <i>ACS Chemical Neuroscience</i> , <b>2019</b> , 10, 4031-4042	5.7	18
39	Activation of Aryl Hydrocarbon Receptor by Kynurenine Impairs Progression and Metastasis of Neuroblastoma. <i>Cancer Research</i> , <b>2019</b> , 79, 5550-5562	10.1	18
38	Calreticulin regulates MYCN expression to control neuronal differentiation and stemness of neuroblastoma. <i>Journal of Molecular Medicine</i> , <b>2019</b> , 97, 325-339	5.5	2
37	Quantitative Measurement of $\beta$ Secretase-mediated Amyloid Precursor Protein and Notch Cleavage in Cell-based Luciferase Reporter Assay Platforms. <i>Journal of Visualized Experiments</i> , <b>2018</b>	1.6	1
36	ErbB2 regulates autophagic flux to modulate the proteostasis of APP-CTFs in Alzheimer's disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2017</b> , 114, E3129-E3138	11.5	40
35	VEGF expression correlates with neuronal differentiation and predicts a favorable prognosis in patients with neuroblastoma. <i>Scientific Reports</i> , <b>2017</b> , 7, 11212	4.9	5
34	[P2035]: AN HERBAL EXTRACT (HE238) SUPPRESSES AMYLOID- $\beta$ 2-ELICITED NEUROTOXICITY IN MICE THROUGH ACTIVATION OF AUTOPHAGY <b>2017</b> , 13, P658-P658		
33	Critical Roles of Dual-Specificity Phosphatases in Neuronal Proteostasis and Neurological Diseases. <i>International Journal of Molecular Sciences</i> , <b>2017</b> , 18,	6.3	14
32	A multidisciplinary team care approach improves outcomes in high-risk pediatric neuroblastoma patients. <i>Oncotarget</i> , <b>2017</b> , 8, 4360-4372	3.3	12
31	Multiple signaling factors and drugs alleviate neuronal death induced by expression of human and zebrafish tau proteins in vivo. <i>Journal of Biomedical Science</i> , <b>2016</b> , 23, 25	13.3	16
30	Diagnostic FDG and FDOPA positron emission tomography scans distinguish the genomic type and treatment outcome of neuroblastoma. <i>Oncotarget</i> , <b>2016</b> , 7, 18774-86	3.3	21
29	Epicatechin isolated from <i>Tripterygium wilfordii</i> extract reduces tau-GFP-induced neurotoxicity in zebrafish embryo through the activation of Nrf2. <i>Biochemical and Biophysical Research Communications</i> , <b>2016</b> , 477, 283-9	3.4	6
28	New Hydroxyquinoline-Based Derivatives as Potent Modulators of Amyloid- $\beta$ Aggregations. <i>Archiv Der Pharmazie</i> , <b>2016</b> , 349, 327-41	4.3	9

27	JARID1B Expression Plays a Critical Role in Chemoresistance and Stem Cell-Like Phenotype of Neuroblastoma Cells. <i>PLoS ONE</i> , <b>2015</b> , 10, e0125343	3.7	38
26	Calreticulin Regulates VEGF-A in Neuroblastoma Cells. <i>Molecular Neurobiology</i> , <b>2015</b> , 52, 758-70	6.2	12
25	Difluorophenylglycinols as new modulators of proteolytic processing of amyloid precursor proteins. <i>Archiv Der Pharmazie</i> , <b>2014</b> , 347, 161-73	4.3	1
24	Discovery of small molecular (D)-leucinamides as potent, Notch-sparing $\beta$ secretase modulators. <i>European Journal of Medicinal Chemistry</i> , <b>2014</b> , 79, 143-51	6.8	3
23	P1-084: LIGAND-ACTIVATED EPHA4 SIGNALING GOVERNS THE PROTEOSTASIS OF APP-BETA CTF TO CONTROL THE LEVEL OF AMYLOID-BETA <b>2014</b> , 10, P333-P333		
22	Aryl hydrocarbon receptor downregulates MYCN expression and promotes cell differentiation of neuroblastoma. <i>PLoS ONE</i> , <b>2014</b> , 9, e88795	3.7	22
21	Calreticulin activates $\alpha$ integrin via fucosylation by fucosyltransferase 1 in J82 human bladder cancer cells. <i>Biochemical Journal</i> , <b>2014</b> , 460, 69-78	3.8	18
20	Presenilin-1 regulates the expression of p62 to govern p62-dependent tau degradation. <i>Molecular Neurobiology</i> , <b>2014</b> , 49, 10-27	6.2	8
19	Ligand-dependent activation of EphA4 signaling regulates the proteolysis of amyloid precursor protein through a Lyn-mediated pathway. <i>Molecular Neurobiology</i> , <b>2014</b> , 49, 1055-68	6.2	4
18	The Nogo-C2/Nogo receptor complex regulates the morphogenesis of zebrafish lateral line primordium through modulating the expression of dkk1b, a Wnt signal inhibitor. <i>PLoS ONE</i> , <b>2014</b> , 9, e86345	3.7	7
17	Retinoic acid-elicited RAR/RXR signaling attenuates A $\beta$ production by directly inhibiting $\beta$ secretase-mediated cleavage of amyloid precursor protein. <i>ACS Chemical Neuroscience</i> , <b>2013</b> , 4, 1093-100	5.7	20
16	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , <b>2012</b> , 8, 445-544	4.2	2783
15	Calreticulin mediates nerve growth factor-induced neuronal differentiation. <i>Journal of Molecular Neuroscience</i> , <b>2012</b> , 47, 571-81	3.3	13
14	Autophagy: a double-edged sword in Alzheimer's disease. <i>Journal of Biosciences</i> , <b>2012</b> , 37, 157-65	2.3	69
13	Corrigendum to Silencing of miR-124 induces neuroblastoma SK-N-SH cell differentiation, cell cycle arrest and apoptosis through promoting AHR [FEBS Lett. 585 (2011) 3582-3586]. <i>FEBS Letters</i> , <b>2012</b> , 586, 107-107	3.8	
12	Nuclear GRP75 binds retinoic acid receptors to promote neuronal differentiation of neuroblastoma. <i>PLoS ONE</i> , <b>2011</b> , 6, e26236	3.7	20
11	Nuclear localized GRP75 binds to retinoic acid receptors (RAR/RXR) and promotes retinoic acid-induced neuronal differentiation of neuroblastoma. <i>FASEB Journal</i> , <b>2011</b> , 25, 1001.12	0.9	
10	Notch1 expression predicts an unfavorable prognosis and serves as a therapeutic target of patients with neuroblastoma. <i>Clinical Cancer Research</i> , <b>2010</b> , 16, 4411-20	12.9	35

9	Caveolin-1 regulates $\beta$ -secretase-mediated APP processing by modulating spatial distribution of $\beta$ -secretase in membrane. <i>Journal of Alzheimer's Disease</i> , <b>2010</b> , 22, 423-42	4.3	22
8	The evolutionarily conserved interaction between LC3 and p62 selectively mediates autophagy-dependent degradation of mutant huntingtin. <i>Cellular and Molecular Neurobiology</i> , <b>2010</b> , 30, 795-806	4.6	35
7	Sodium selenite inhibits gamma-secretase activity through activation of ERK. <i>Neuroscience Letters</i> , <b>2008</b> , 440, 38-43	3.3	31
6	Identification of GRP75 as an independent favorable prognostic marker of neuroblastoma by a proteomics analysis. <i>Clinical Cancer Research</i> , <b>2008</b> , 14, 6237-45	12.9	26
5	Tumor necrosis factor-alpha-elicited stimulation of gamma-secretase is mediated by c-Jun N-terminal kinase-dependent phosphorylation of presenilin and nicastrin. <i>Molecular Biology of the Cell</i> , <b>2008</b> , 19, 4201-12	3.5	49
4	New 1,2,3,4-tetrahydroisoquinoline derivatives as modulators of proteolytic cleavage of amyloid precursor proteins. <i>Bioorganic and Medicinal Chemistry</i> , <b>2008</b> , 16, 1957-65	3.4	15
3	Unnatural amino acid-substituted (hydroxyethyl)urea peptidomimetics inhibit gamma-secretase and promote the neuronal differentiation of neuroblastoma cells. <i>Molecular Pharmacology</i> , <b>2007</b> , 71, 588-601	4.3	21
2	A high-throughput screen to identify inhibitors of amyloid beta-protein precursor processing. <i>Journal of Biomolecular Screening</i> , <b>2005</b> , 10, 1-12		26
1	Tumor necrosis factor-alpha, interleukin-1beta, and interferon-gamma stimulate gamma-secretase-mediated cleavage of amyloid precursor protein through a JNK-dependent MAPK pathway. <i>Journal of Biological Chemistry</i> , <b>2004</b> , 279, 49523-32	5.4	247