

Hiroaki Ishii

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5941381/hiroaki-ishii-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

80
papers

1,604
citations

20
h-index

37
g-index

83
ext. papers

1,773
ext. citations

2.6
avg, IF

4.54
L-index

#	Paper	IF	Citations
80	Tolerance and acclimation of photosynthesis of nine urban tree species to warmer growing conditions. <i>Trees - Structure and Function</i> , 2021 , 35, 1793	2.6	0
79	Predicting effects of climate change on productivity and persistence of forest trees. <i>Ecological Research</i> , 2020 , 35, 562-574	1.9	2
78	Regional and topographic growth variation among 45-year-old clonal plantations of <i>Cryptomeria japonica</i> : effects of genotype and phenotypic plasticity. <i>Journal of Forest Research</i> , 2020 , 25, 329-338	1.4	1
77	Acclimation potential of three evergreen tree species to wind-induced water stress in an urban green-roof environment. <i>Urban Forestry and Urban Greening</i> , 2019 , 43, 126386	5.4	2
76	Comparison of Leaf Functional Trait Plasticity between Exotic and Native <i>Ligustrum</i> Species in Japan. <i>Journal of the Japanese Forest Society</i> , 2019 , 101, 221-226	0.2	
75	Height-related variations of leaf traits reflect strategies for maintaining photosynthetic and hydraulic homeostasis in mature and old <i>Pinus densiflora</i> trees. <i>Oecologia</i> , 2019 , 189, 317-328	2.9	8
74	Large, retained trees of <i>Cryptomeria japonica</i> functioned as refugia for canopy woody plants after logging 350 years ago in Yakushima, Japan. <i>Forest Ecology and Management</i> , 2018 , 409, 457-467	3.9	16
73	Evaluating restoration success of a 40-year-old urban forest in reference to mature natural forest. <i>Urban Forestry and Urban Greening</i> , 2018 , 32, 123-132	5.4	12
72	Variation of intra-crown leaf plasticity of <i>Fagus crenata</i> across its geographical range in Japan. <i>Forest Ecology and Management</i> , 2018 , 429, 437-448	3.9	10
71	Crown dynamics and wood production of Douglas-fir trees in an old-growth forest. <i>Forest Ecology and Management</i> , 2017 , 384, 157-168	3.9	14
70	Hydraulic Architecture and Function of Tall Trees. <i>Journal of the Japanese Forest Society</i> , 2017 , 99, 74-83	0.2	3
69	Physiological and morphological acclimation to height in cupressoid leaves of 100-year-old <i>Chamaecyparis obtusa</i> . <i>Tree Physiology</i> , 2017 , 37, 1327-1336	4.2	2
68	Water retained in tall <i>Cryptomeria japonica</i> leaves as studied by infrared micro-spectroscopy. <i>Tree Physiology</i> , 2017 , 37, 1367-1378	4.2	6
67	Leaf water maintains daytime transpiration in young <i>Cryptomeria japonica</i> trees. <i>Tree Physiology</i> , 2017 , 37, 1394-1403	4.2	7
66	Oviposition site selection by Japanese gypsy moth (<i>Lymatria dispar japonica</i>) in a warm-temperate secondary forest in western Japan. <i>Forest Science and Technology</i> , 2016 , 12, 130-136	1.5	3
65	Vegetation recovery after removal of invasive <i>Trachycarpus fortunei</i> in a fragmented urban shrine forest. <i>Urban Forestry and Urban Greening</i> , 2016 , 15, 53-57	5.4	8
64	Function and structure of leaves contributing to increasing water storage with height in the tallest <i>Cryptomeria japonica</i> trees of Japan. <i>Trees - Structure and Function</i> , 2016 , 30, 141-152	2.6	19

63	Urban forest management in Melbourne city based on “Water-Sensitive Urban Design: Urban design considering the water cycle”; <i>Journal of the Japanese Society of Revegetation Technology</i> , 2016 , 42, 455-459	0.1	
62	Twenty-one years of stand dynamics in a 33-year-old urban forest restoration site at Kobe Municipal Sports Park, Japan. <i>Urban Forestry and Urban Greening</i> , 2015 , 14, 309-314	5.4	8
61	Intraspecific variation in fine root respiration and morphology in response to in situ soil nitrogen fertility in a 100-year-old <i>Chamaecyparis obtusa</i> forest. <i>Oecologia</i> , 2015 , 179, 959-67	2.9	15
60	Urban forestry: toward creation of diverse urban green space. <i>Journal of the Japanese Society of Revegetation Technology</i> , 2015 , 40, 505-507	0.1	1
59	Pushing the limits to tree height: could foliar water storage compensate for hydraulic constraints in <i>Sequoia sempervirens</i> ?. <i>Functional Ecology</i> , 2014 , 28, 1087-1093	5.6	44
58	Stand Structure of an Abandoned Deciduous Broadleaf Secondary Forest Adjacent to Lucidophyllous Forest and Agricultural Fields.. <i>Journal of the Japanese Forest Society</i> , 2014 , 96, 75-82	0.2	8
57	Application of Laser Remote Sensing to Forest Ecological Research. <i>Journal of the Japanese Forest Society</i> , 2014 , 96, 168-181	0.2	4
56	Comparison of branch growth estimates in <i>Larix gmelinii</i> by several methods. <i>Journal of Forest Research</i> , 2013 , 18, 345-352	1.4	
55	The need for a canopy perspective to understand the importance of phenotypic plasticity for promoting species coexistence and light-use complementarity in forest ecosystems. <i>Ecological Research</i> , 2013 , 28, 191-198	1.9	26
54	Response of Ant Community Structure to Understory Removal in a Line-thinned Japanese Cedar (<i>Cryptomeria japonica</i>) Plantation. <i>Journal of the Japanese Forest Society</i> , 2013 , 95, 95-100	0.2	1
53	Variation in light-intercepting area and photosynthetic rate of sun and shade shoots of two <i>Picea</i> species in relation to the angle of incoming light. <i>Tree Physiology</i> , 2012 , 32, 1227-36	4.2	12
52	Ant Community Structure and Related Environmental Factors after Line Thinning in Japanese Cedar (<i>Cryptomeria japonica</i>) Plantations. <i>Journal of the Japanese Forest Society</i> , 2012 , 94, 36-41	0.2	
51	How Do Changes in Leaf/Shoot Morphology and Crown Architecture Affect Growth and Physiological Function of Tall Trees?. <i>Tree Physiology</i> , 2011 , 215-232		7
50	Very fine roots respond to soil depth: biomass allocation, morphology, and physiology in a broad-leaved temperate forest. <i>Ecological Research</i> , 2011 , 26, 95-104	1.9	68
49	Aboveground productivity of an unsuccessful 140-year-old <i>Cryptomeria japonica</i> plantation in northern Kyushu, Japan. <i>Journal of Forest Research</i> , 2011 , 16, 268-274	1.4	5
48	Light acclimation potential and carry-over effects vary among three evergreen tree species with contrasting patterns of leaf emergence and maturation. <i>Tree Physiology</i> , 2011 , 31, 819-30	4.2	15
47	Physiological and morphological properties of <i>Cinnamomum camphora</i> cuttings: comparison with seedlings. <i>Journal of the Japanese Society of Revegetation Technology</i> , 2011 , 37, 21-25	0.1	
46	Climatic factors affecting radial growth of <i>Betula ermanii</i> and <i>Betula platyphylla</i> in Kamchatka. <i>Canadian Journal of Forest Research</i> , 2010 , 40, 273-285	1.9	11

45	Seasonal prevalence of arthropods after line thinning of overstocked Japanese cedar (<i>Cryptomeria japonica</i> D. Don) plantations in central Japan. <i>Landscape and Ecological Engineering</i> , 2010 , 6, 43-52	2	5
44	Integrating ecological and cultural values toward conservation and utilization of shrine/temple forests as urban green space in Japanese cities. <i>Landscape and Ecological Engineering</i> , 2010 , 6, 307-315	2	39
43	Vertical stratification and effects of crown damage on maximum tree height in mixed coniferBroadleaf forests of Yakushima Island, southern Japan. <i>Plant Ecology</i> , 2010 , 211, 27-36	1.7	8
42	The role of crown architecture, leaf phenology and photosynthetic activity in promoting complementary use of light among coexisting species in temperate forests. <i>Ecological Research</i> , 2010 , 25, 715-722	1.9	93
41	Branch and Foliage Mass and their Vertical Distribution in a 90-year-old <i>Chamaecyparis obtusa</i> Plantation.. <i>Journal of the Japanese Forest Society</i> , 2010 , 92, 63-71	0.2	1
40	Field Measurement of Heterotrophic Respiration of Root Litter Using a Small Chamber System.. <i>Journal of the Japanese Forest Society</i> , 2010 , 92, 269-272	0.2	
39	Fine root morphological traits determine variation in root respiration of <i>Quercus serrata</i> . <i>Tree Physiology</i> , 2009 , 29, 579-85	4.2	81
38	Convergence of leaf display and photosynthetic characteristics of understory <i>Abies amabilis</i> and <i>Tsuga heterophylla</i> in an old-growth forest in southwestern Washington State, USA. <i>Tree Physiology</i> , 2009 , 29, 989-98	4.2	7
37	Basal reiteration improves the hydraulic functional status of mature <i>Cinnamomum camphora</i> trees. <i>Trees - Structure and Function</i> , 2009 , 23, 317-323	2.6	4
36	Arthropods as bioindicators of sustainable forest management, with a focus on plantation forests. <i>Applied Entomology and Zoology</i> , 2009 , 44, 1-11	1.5	98
35	Different physiological and morphological responses of leaves and branches of <i>Ligustrum japonicum</i> and invasive <i>L. lucidum</i> to the light environment.. <i>Journal of the Japanese Society of Revegetation Technology</i> , 2009 , 35, 45-50	0.1	2
34	Simulation Study of Size-structure Dynamics with Changing Spatial Pattern of Tree Sizes in a Lattice-planted Japanese Cedar (<i>Cryptomeria japonica</i> D. Don) Plantation. <i>Journal of Forest Planning</i> , 2009 , 15, 11-19	0	
33	Ecological restoration of a fragmented urban shrine forest in southeastern Hyogo Prefecture, Japan: Initial effects of the removal of invasive <i>Trachycarpus fortunei</i> . <i>Urban Ecosystems</i> , 2008 , 11, 309-316	2.8	13
32	Establishment and growth pattern of <i>Pinus pumila</i> under a forest canopy in central Kamchatka. <i>Ecological Research</i> , 2008 , 23, 831-840	1.9	6
31	Effects of individual size, local competition and canopy closure on the stem volume growth in a monoclonal Japanese cedar (<i>Cryptomeria japonica</i> D. Don) plantation. <i>Ecological Research</i> , 2008 , 23, 953-964	1.9	19
30	Line thinning promotes stand growth and understory diversity in Japanese cedar (<i>Cryptomeria japonica</i> D. Don) plantations. <i>Journal of Forest Research</i> , 2008 , 13, 73-78	1.4	28
29	Removal of understory dwarf bamboo (<i>Sasa kurilensis</i>) induces changes in water-relations characteristics of overstory <i>Betula ermanii</i> trees. <i>Journal of Forest Research</i> , 2008 , 13, 101-109	1.4	17
28	Hydrostatic constraints on morphological exploitation of light in tall <i>Sequoia sempervirens</i> trees. <i>Oecologia</i> , 2008 , 156, 751-63	2.9	79

27	Line thinning fosters the abundance and diversity of understory Hymenoptera (Insecta) in Japanese cedar (<i>Cryptomeria japonica</i> D. Don) plantations. <i>Journal of Forest Research</i> , 2007 , 12, 14-23	1.4	20
26	Line thinning enhances diversity of Coleoptera in overstocked <i>Cryptomeria japonica</i> plantations in central Japan. <i>Arthropod-Plant Interactions</i> , 2007 , 1, 175-185	2.2	16
25	Plasticity of shoot and needle morphology and photosynthesis of two <i>Picea</i> species with different site preferences in northern Japan. <i>Tree Physiology</i> , 2007 , 27, 1595-605	4.2	21
24	Physiological and ecological implications of adaptive reiteration as a mechanism for crown maintenance and longevity. <i>Tree Physiology</i> , 2007 , 27, 455-62	4.2	39
23	Evaluation of shrine forests as urban green space. <i>Landscape Ecology and Management</i> , 2007 , 12, 1-7	0	8
22	Removal of invasive <i>Trachycarpus fortunei</i> at Nishinomiya Shrine, Hyogo Prefecture, Japan. <i>Landscape Ecology and Management</i> , 2007 , 12, 35-43	0	
21	Biomass and dynamics of attached dead branches in the canopy of 450-year-old Douglas-fir trees. <i>Canadian Journal of Forest Research</i> , 2006 , 36, 378-389	1.9	10
20	Biomass Cycling and Soil Properties in an Agroforestry-based Plantation System of kayu putih (<i>Melaleuca leucadendron</i> LINN) in East Java, Indonesia. <i>Agroforestry Systems</i> , 2006 , 67, 135-145	2	9
19	Productivity of kayu putih (<i>Melaleuca leucadendron</i> LINN) tree plantation managed in non-timber forest production systems in Java, Indonesia. <i>Agroforestry Systems</i> , 2005 , 64, 143-155	2	5
18	Tree growth and competition in a <i>Betula platyphylla</i> - <i>Larix cajanderi</i> post-fire forest in central Kamchatka. <i>Annals of Botany</i> , 2004 , 94, 333-43	4.1	34
17	Model analysis of the importance of reiteration for branch longevity in <i>Pseudotsuga menziesii</i> compared with <i>Abies grandis</i> . <i>Canadian Journal of Botany</i> , 2004 , 82, 892-909		14
16	Acclimation of shoot and needle morphology and photosynthesis of two <i>Picea</i> species to differences in soil nutrient availability. <i>Tree Physiology</i> , 2003 , 23, 453-61	4.2	12
15	Persistence of <i>Pseudotsuga menziesii</i> (Douglas-fir) in temperate coniferous forests of the Pacific Northwest Coast, USA. <i>Folia Geobotanica</i> , 2002 , 37, 63-69	1.4	19
14	The relationship between tree height and leaf area: sapwood area ratio. <i>Oecologia</i> , 2002 , 132, 12-20	2.9	246
13	Variation in specific needle area of old-growth Douglas-fir in relation to needle age, within-crown position and epicormic shoot production. <i>Tree Physiology</i> , 2002 , 22, 31-40	4.2	36
12	The role of epicormic shoot production in maintaining foliage in old <i>Pseudotsuga menziesii</i> (Douglas-fir) trees II. Basal reiteration from older branch axes. <i>Canadian Journal of Botany</i> , 2002 , 80, 916-926		26
11	Age-related development of crown structure in coastal Douglas-fir trees. <i>Forest Ecology and Management</i> , 2002 , 169, 257-270	3.9	53
10	The method of synthesis in ecology. <i>Oikos</i> , 2001 , 93, 153-160	4	14

9	Crown structure of old-growth Douglas-fir in the western Cascade Range, Washington. <i>Canadian Journal of Forest Research</i> , 2001 , 31, 1250-1261	1.9	35
8	The role of epicormic shoot production in maintaining foliage in old <i>Pseudotsuga menziesii</i> (Douglas-fir) trees. <i>Canadian Journal of Botany</i> , 2001 , 79, 251-264		50
7	Height growth and vertical development of an old-growth <i>Pseudotsuga-Tsuga</i> forest in southwestern Washington State, U.S.A.. <i>Canadian Journal of Forest Research</i> , 2000 , 30, 17-24	1.9	32
6	Branch growth and crown form in old coastal Douglas-fir. <i>Forest Ecology and Management</i> , 2000 , 131, 81-91	3.9	34
5	Height growth and vertical development of an old-growth <i>Pseudotsuga-Tsuga</i> forest in southwestern Washington State, U.S.A.. <i>Canadian Journal of Forest Research</i> , 2000 , 30, 17-24	1.9	17
4	Effects of the spatial arrangement of aerial stems and current-year shoots on the demography and growth of <i>Hydrangea hirta</i> in a light-limited environment. <i>New Phytologist</i> , 1997 , 136, 443-453	9.8	15
3	Tree Architecture in <i>Ocotea</i> (Lauraceae): Do Ant-Occupied Species Differ Structurally from Non-Ant-Occupied Species?. <i>Tropics</i> , 1995 , 4, 239-245	0.9	
2	One large tree crown can be defined as a local hotspot for plant species diversity in a forest ecosystem: a case study in temperate old-growth forest. <i>Plant Ecology</i> , 1	1.7	0
1	Spatial and functional niche overlap between invasive <i>Ligustrum lucidum</i> and native woody species in an urban shrine forest in Japan. <i>Landscape and Ecological Engineering</i> , 1		2