

Franz Bucar

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5945185/franz-bucar-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

60
papers

1,846
citations

26
h-index

41
g-index

65
ext. papers

2,121
ext. citations

4.9
avg, IF

5
L-index

#	Paper	IF	Citations
60	Globularia alypum L. and Related Species: LC-MS Profiles and Antidiabetic, Antioxidant, Anti-Inflammatory, Antibacterial and Anticancer Potential. <i>Pharmaceuticals</i> , 2022 , 15, 506	5.2	0
59	Flavonoids as Inhibitors of Bacterial Efflux Pumps. <i>Molecules</i> , 2021 , 26,	4.8	2
58	Anti-adhesion activity of phytochemicals to prevent Campylobacter jejuni biofilm formation on abiotic surfaces. <i>Phytochemistry Reviews</i> , 2021 , 20, 55-84	7.7	20
57	Efflux Pump Inhibition and Resistance Modulation in by and Its Coumarins. <i>Antibiotics</i> , 2021 , 10,	4.9	3
56	Antibiofilm Potential of Preparations against Campylobacter jejuni. <i>Applied and Environmental Microbiology</i> , 2021 , 87, e0109921	4.8	1
55	Flavonoids as Novel Efflux Pump Inhibitors and Antimicrobials Against Both Environmental and Pathogenic Intracellular Mycobacterial Species. <i>Molecules</i> , 2020 , 25,	4.8	24
54	In Vitro Effect of the Common Culinary Herb Winter Savory () against the Infamous Food Pathogen. <i>Foods</i> , 2020 , 9,	4.9	7
53	Targeting fish spoilers Pseudomonas and Shewanella with oregano and nettle extracts. <i>International Journal of Food Microbiology</i> , 2020 , 328, 108664	5.8	15
52	Palladium-catalysed synthesis of arylnaphthoquinones as antiprotozoal and antimycobacterial agents. <i>European Journal of Medicinal Chemistry</i> , 2020 , 207, 112837	6.8	2
51	Antimicrobial and Efflux Pump Inhibitory Activity of Carvotacetones from Against Mycobacteria. <i>Antibiotics</i> , 2020 , 9,	4.9	3
50	A new tetracyclic saponin from L. and its neuroprotective and hMAO-B inhibiting activity. <i>Natural Product Research</i> , 2020 , 34, 511-517	2.3	9
49	Qualitative and Quantitative Analysis of Different Rhodiola rosea Rhizome Extracts by UHPLC-DAD-ESI-MSn. <i>Scientia Pharmaceutica</i> , 2019 , 87, 8	4.3	14
48	Phenolic compounds of Iris adriatica and their antimycobacterial effects. <i>Acta Pharmaceutica</i> , 2019 , 69, 673-681	3.2	4
47	Resistance modulatory and efflux-inhibitory activities of capsaicinoids and capsinoids. <i>Bioorganic Chemistry</i> , 2019 , 82, 378-384	5.1	11
46	Metabolic profiling of rhizomes of native populations of the strictly endemic Croatian species Iris adriatica. <i>Plant Biosystems</i> , 2019 , 153, 317-324	1.6	11
45	Antimicrobial Natural Products Against Campylobacter. <i>Sustainable Development and Biodiversity</i> , 2018 , 3-30	2.1	1
44	Antiadhesion activity of juniper (Juniperus communis L.) preparations against Campylobacter jejuni evaluated with PCR-based methods. <i>Phytotherapy Research</i> , 2018 , 32, 542-550	6.7	10

43	Aqueous Extracts of Wild Mushrooms Show Antimicrobial and Antiadhesion Activities against Bacteria and Fungi. <i>Phytotherapy Research</i> , 2017 , 31, 1971-1976	6.7	10
42	Efficient identification of flavones, flavanones and their glycosides in routine analysis via off-line combination of sensitive NMR and HPLC experiments. <i>Food Chemistry</i> , 2017 , 218, 600-609	8.5	33
41	LC-PDA-ESI-MS analysis of phenolic and iridoid compounds from <i>Globularia</i> spp. <i>Journal of Mass Spectrometry</i> , 2016 , 51, 1211-1236	2.2	32
40	Anti-adhesion activity of thyme (<i>Thymus vulgaris</i> L.) extract, thyme post-distillation waste, and olive (<i>Olea europea</i> L.) leaf extract against <i>Campylobacter jejuni</i> on polystyrene and intestine epithelial cells. <i>Journal of the Science of Food and Agriculture</i> , 2016 , 96, 2723-30	4.3	24
39	Pumpkin seed extract: Cell growth inhibition of hyperplastic and cancer cells, independent of steroid hormone receptors. <i>Phytotherapy Research</i> , 2016 , 30, 150-6	3.2	38
38	Attenuation of Adhesion, Biofilm Formation and Quorum Sensing of <i>Campylobacter jejuni</i> by <i>Euodia ruticarpa</i> . <i>Phytotherapy Research</i> , 2016 , 30, 1527-32	6.7	32
37	Content of phenolic compounds in wild populations of <i>Epilobium angustifolium</i> growing at different altitudes. <i>Pharmaceutical Biology</i> , 2015 , 53, 1576-82	3.8	20
36	Plant derived inhibitors of bacterial efflux pumps: an update. <i>Phytochemistry Reviews</i> , 2015 , 14, 961-974	7.7	39
35	<i>Alpinia katsumadai</i> Extracts Inhibit Adhesion and Invasion of <i>Campylobacter jejuni</i> in Animal and Human Foetal Small Intestine Cell Lines. <i>Phytotherapy Research</i> , 2015 , 29, 1585-9	6.7	14
34	Antibiotic resistance modulation and modes of action of (-)- β -pinene in <i>Campylobacter jejuni</i> . <i>PLoS ONE</i> , 2015 , 10, e0122871	3.7	63
33	1,2-substituted 4-(1H)-quinolones: synthesis, antimalarial and antitrypanosomal activities in vitro. <i>Molecules</i> , 2014 , 19, 14204-20	4.8	12
32	Reduction of microbiological risk in minced meat by a combination of natural antimicrobials. <i>Journal of the Science of Food and Agriculture</i> , 2014 , 94, 2758-65	4.3	11
31	Natural product isolation--how to get from biological material to pure compounds. <i>Natural Product Reports</i> , 2013 , 30, 525-45	15.1	232
30	Compounds of <i>Alpinia katsumadai</i> as potential efflux inhibitors in <i>Mycobacterium smegmatis</i> . <i>Bioorganic and Medicinal Chemistry</i> , 2012 , 20, 2701-6	3.4	43
29	The volatile fraction of herbal teas. <i>Phytochemistry Reviews</i> , 2012 , 11, 245-254	7.7	8
28	Putative mycobacterial efflux inhibitors from the seeds of <i>Aframomum melegueta</i> . <i>Journal of Natural Products</i> , 2012 , 75, 1393-9	4.9	43
27	Guaianolides and volatile compounds in chamomile tea. <i>Plant Foods for Human Nutrition</i> , 2012 , 67, 129-35	5.9	24
26	Synthesis and antibacterial evaluation of a new series of N-Alkyl-2-alkynyl/(E)-alkenyl-4-(1H)-quinolones. <i>Molecules</i> , 2012 , 17, 8217-40	4.8	14

25	Influence of saponin plants on the volatile fraction of thyme in herbal teas. <i>Phytotherapy Research</i> , 2011 , 82, 903-10	3.2	14
24	Design, synthesis and antimycobacterial activities of 1-methyl-2-alkenyl-4(1H)-quinolones. <i>Bioorganic and Medicinal Chemistry</i> , 2011 , 19, 567-79	3.4	55
23	Synthesis of N-substituted 2-[(1E)-alkenyl]-4-(1H)-quinolone derivatives as antimycobacterial agents against non-tubercular mycobacteria. <i>European Journal of Medicinal Chemistry</i> , 2011 , 46, 2091-101	6.8	18
22	Interaction of N-methyl-2-alkenyl-4-quinolones with ATP-dependent MurE ligase of Mycobacterium tuberculosis: antibacterial activity, molecular docking and inhibition kinetics. <i>Journal of Antimicrobial Chemotherapy</i> , 2011 , 66, 1766-72	5.1	33
21	Investigation of the volatile fraction of rosemary infusion extracts. <i>Scientia Pharmaceutica</i> , 2010 , 78, 483-92	4.3	26
20	Antiprotozoal activity of drimane and coloratane sesquiterpenes towards Trypanosoma brucei rhodesiense and Plasmodium falciparum in vitro. <i>Phytotherapy Research</i> , 2010 , 24, 1468-72	6.7	33
19	Volatile fraction of lavender and bitter fennel infusion extracts. <i>Natural Product Communications</i> , 2010 , 5, 1431-6	0.9	12
18	Influence of altitudinal variation on the content of phenolic compounds in wild populations of Calluna vulgaris, Sambucus nigra, and Vaccinium myrtillus. <i>Journal of Agricultural and Food Chemistry</i> , 2008 , 56, 9080-6	5.7	100
17	Plant phenolic compounds as ethidium bromide efflux inhibitors in Mycobacterium smegmatis. <i>Journal of Antimicrobial Chemotherapy</i> , 2008 , 62, 345-8	5.1	113
16	Constituents of the stem bark of Discopodium penninervium and their LTB4 and COX-1 and -2 inhibitory activities. <i>Phytochemistry</i> , 2008 , 69, 982-7	4	38
15	Modulation of isoniazid susceptibility by flavonoids in Mycobacterium. <i>Phytochemistry Letters</i> , 2008 , 1, 71-75	1.9	31
14	Antimycobacterial polyacetylenes from Levisticum officinale. <i>Phytotherapy Research</i> , 2008 , 22, 681-4	6.7	46
13	Quantum Chemical Studies on Structure Activity Relationship of Natural Product Polyacetylenes. <i>Theoretical Chemistry Accounts</i> , 2007 , 117, 247-252	1.9	7
12	Knipholone, a selective inhibitor of leukotriene metabolism. <i>Phytomedicine</i> , 2006 , 13, 452-6	6.5	26
11	In vitro 12(S)-HETE and leukotriene metabolism inhibitory activity of sesquiterpenes of Warburgia ugandensis. <i>Planta Medica</i> , 2006 , 72, 754-6	3.1	9
10	In vitro 12(S)-HETE inhibitory activities of naphthoquinones isolated from the root bark of Euclea racemosa ssp. schimperii. <i>Journal of Ethnopharmacology</i> , 2005 , 102, 191-6	5	14
9	Quinolone alkaloids from Evodia rutaecarpa: a potent new group of antimycobacterial compounds. <i>International Journal of Antimicrobial Agents</i> , 2005 , 26, 262-4	14.3	35
8	Bioactive constituents of Artemisia monosperma. <i>Phytochemistry</i> , 2005 , 66, 233-9	4	43

7	Sesquiterpenes from <i>Warburgia ugandensis</i> and their antimycobacterial activity. <i>Phytochemistry</i> , 2005 , 66, 2309-15	4	50
6	Lipoxygenase inhibitors from natural plant sources. Part 1: Medicinal plants with inhibitory activity on arachidonate 5-lipoxygenase and 5-lipoxygenase[sol]cyclooxygenase. <i>Phytotherapy Research</i> , 2005 , 19, 81-102	6.7	98
5	Lipoxygenase inhibitors from natural plant sources. Part 2: medicinal plants with inhibitory activity on arachidonate 12-lipoxygenase, 15-lipoxygenase and leukotriene receptor antagonists. <i>Phytotherapy Research</i> , 2005 , 19, 263-72	6.7	50
4	Antimalarial compounds from <i>Kniphofia foliosa</i> roots. <i>Phytotherapy Research</i> , 2005 , 19, 472-6	6.7	37
3	Inhibitory activity of <i>Juniperus communis</i> on 12(S)-HETE production in human platelets. <i>Planta Medica</i> , 2004 , 70, 471-4	3.1	18
2	The antimycobacterial components of hops (<i>Humulus lupulus</i>) and their dereplication. <i>Phytotherapy Research</i> , 2004 , 18, 774-6	6.7	38
1	Ostruthin: an antimycobacterial coumarin from the roots of <i>Peucedanum ostruthium</i> . <i>Planta Medica</i> , 2003 , 69, 369-71	3.1	67