## Frederik Persson

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/594689/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. European Heart Journal, 2020, 41, 255-323.	1.0	2,811
2	Cardiorenal End Points in a Trial of Aliskiren for Type 2 Diabetes. New England Journal of Medicine, 2012, 367, 2204-2213.	13.9	1,145
3	Aliskiren Combined with Losartan in Type 2 Diabetes and Nephropathy. New England Journal of Medicine, 2008, 358, 2433-2446.	13.9	1,051
4	Cardiovascular mortality and morbidity in patients with type 2 diabetes following initiation of sodium-glucose co-transporter-2 inhibitors versus other glucose-lowering drugs (CVD-REAL Nordic): a multinational observational analysis. Lancet Diabetes and Endocrinology,the, 2017, 5, 709-717.	5.5	285
5	Urinary Proteomics for Early Diagnosis in Diabetic Nephropathy. Diabetes, 2012, 61, 3304-3313.	0.3	221
6	Diagnosis and Prediction of CKD Progression by Assessment of Urinary Peptides. Journal of the American Society of Nephrology: JASN, 2015, 26, 1999-2010.	3.0	205
7	Diagnosis of diabetic kidney disease: state of the art and future perspective. Kidney International Supplements, 2018, 8, 2-7.	4.6	202
8	Early detection of diabetic kidney disease by urinary proteomics and subsequent intervention with spironolactone to delay progression (PRIORITY): a prospective observational study and embedded randomised placebo-controlled trial. Lancet Diabetes and Endocrinology,the, 2020, 8, 301-312.	5.5	166
9	Dapagliflozin is associated with lower risk of cardiovascular events and allâ€cause mortality in people with type 2 diabetes ( <scp>CVDâ€REAL Nordic</scp> ) when compared with dipeptidyl peptidaseâ€4 inhibitor therapy: <scp>A</scp> multinational observational study. Diabetes, Obesity and Metabolism, 2018, 20, 344-351.	2.2	164
10	Implementation of proteomic biomarkers: making it work. European Journal of Clinical Investigation, 2012, 42, 1027-1036.	1.7	151
11	Renal Effects of Aliskiren Compared With and in Combination With Irbesartan in Patients With Type 2 Diabetes, Hypertension, and Albuminuria. Diabetes Care, 2009, 32, 1873-1879.	4.3	147
12	A urinary peptide biomarker set predicts worsening of albuminuria in type 2 diabetes mellitus. Diabetologia, 2013, 56, 259-267.	2.9	128
13	Time course of the antiproteinuric and antihypertensive effects of direct renin inhibition in type 2 diabetes. Kidney International, 2008, 73, 1419-1425.	2.6	121
14	YKL-40, a Marker of Inflammation and Endothelial Dysfunction, Is Elevated in Patients With Type 1 Diabetes and Increases With Levels of Albuminuria. Diabetes Care, 2009, 32, 323-328.	4.3	117
15	Differential Effects of Dapagliflozin on Cardiovascular Risk Factors at Varying Degrees of Renal Function. Clinical Journal of the American Society of Nephrology: CJASN, 2017, 12, 751-759.	2.2	114
16	The dapagliflozin and prevention of adverse outcomes in chronic kidney disease (DAPA-CKD) trial: baseline characteristics. Nephrology Dialysis Transplantation, 2020, 35, 1700-1711.	0.4	107
17	Multicentre prospective validation of a urinary peptidome-based classifier for the diagnosis of type 2 diabetic nephropathy. Nephrology Dialysis Transplantation, 2014, 29, 1563-1570.	0.4	106
18	Proteomic prediction and Renin angiotensin aldosterone system Inhibition prevention Of early diabetic nephRopathy in TYpe 2 diabetic patients with normoalbuminuria (PRIORITY): essential study design and rationale of a randomised clinical multicentre trial. BMI Open, 2016, 6, e010310.	0.8	103

#	Article	IF	CITATIONS
19	Arterial Stiffness Is Associated With Cardiovascular, Renal, Retinal, and Autonomic Disease in Type 1 Diabetes. Diabetes Care, 2013, 36, 715-721.	4.3	99
20	Progressive Decline in Estimated Glomerular Filtration Rate in Patients With Diabetes After Moderate Loss in Kidney Function—Even Without Albuminuria. Diabetes Care, 2019, 42, 1886-1894.	4.3	99
21	Intensified multifactorial intervention in type 2 diabetics with microalbuminuria leads to long-term renal benefits. Kidney International, 2017, 91, 982-988.	2.6	80
22	Irbesartan Treatment Reduces Biomarkers of Inflammatory Activity in Patients With Type 2 Diabetes and Microalbuminuria: An IRMA 2 Substudy. Diabetes, 2006, 55, 3550-3555.	0.3	77
23	The effect of liraglutide on renal function: A randomized clinical trial. Diabetes, Obesity and Metabolism, 2017, 19, 239-247.	2.2	77
24	Microalbuminuria: A parameter that has changed diabetes care. Diabetes Research and Clinical Practice, 2015, 107, 1-8.	1.1	75
25	Endothelial dysfunction and inflammation predict development of diabetic nephropathy in the Irbesartan in Patients with Type 2 Diabetes and Microalbuminuria (IRMA 2) study. Scandinavian Journal of Clinical and Laboratory Investigation, 2008, 68, 731-738.	0.6	74
26	Spironolactone diminishes urinary albumin excretion in patients with type 1 diabetes and microalbuminuria: a randomized placeboâ€controlled crossover study. Diabetic Medicine, 2012, 29, e184-90.	1.2	71
27	Urinary proteomics predict onset of microalbuminuria in normoalbuminuric type 2 diabetic patients, a sub-study of the DIRECT-Protect 2 study. Nephrology Dialysis Transplantation, 2017, 32, gfw292.	0.4	66
28	Tissue Renin–Angiotensin Systems: A Unifying Hypothesis of Metabolic Disease. Frontiers in Endocrinology, 2014, 5, 23.	1.5	65
29	Noninvasive diagnosis of chronic kidney diseases using urinary proteome analysis. Nephrology Dialysis Transplantation, 2017, 32, gfw337.	0.4	62
30	Time course and mechanisms of the antiâ€hypertensive and renal effects of liraglutide treatment. Diabetic Medicine, 2015, 32, 343-352.	1.2	61
31	Effects of Dapagliflozin on Volume Status When Added to Renin–Angiotensin System Inhibitors. Journal of Clinical Medicine, 2019, 8, 779.	1.0	61
32	Impact of Baseline Renal Function on the Efficacy and Safety of Aliskiren Added to Losartan in Patients With Type 2 Diabetes and Nephropathy. Diabetes Care, 2010, 33, 2304-2309.	4.3	58
33	Initial Angiotensin Receptor Blockade–Induced Decrease in Albuminuria Is Associated With Long-Term Renal Outcome in Type 2 Diabetic Patients With Microalbuminuria. Diabetes Care, 2011, 34, 2078-2083.	4.3	58
34	Glucagon-like peptide 1 receptor agonist (GLP-1 RA): long-term effect on kidney function in patients with type 2 diabetes. Journal of Diabetes and Its Complications, 2015, 29, 670-674.	1.2	58
35	Baseline characteristics in the Aliskiren Trial in Type 2 Diabetes Using Cardio-Renal Endpoints (ALTITUDE). JRAAS - Journal of the Renin-Angiotensin-Aldosterone System, 2012, 13, 387-393.	1.0	56
36	Aliskiren in Combination with Losartan Reduces Albuminuria Independent of Baseline Blood Pressure in Patients with Type 2 Diabetes and Nephropathy. Clinical Journal of the American Society of Nephrology: CJASN, 2011, 6, 1025-1031.	2.2	53

#	Article	IF	CITATIONS
37	Is a reduction in albuminuria associated with renal and cardiovascular protection? A <i>post hoc</i> analysis of the <scp>ALTITUDE</scp> trial. Diabetes, Obesity and Metabolism, 2016, 18, 169-177.	2.2	49
38	The Impact of Sotagliflozin on Renal Function, Albuminuria, Blood Pressure, and Hematocrit in Adults With Type 1 Diabetes. Diabetes Care, 2019, 42, 1921-1929.	4.3	47
39	Efficacy and Safety of Dapagliflozin by Baseline Glycemic Status: A Prespecified Analysis From the DAPA-CKD Trial. Diabetes Care, 2021, 44, 1894-1897.	4.3	47
40	Symmetric and asymmetric dimethylarginine as risk markers of cardiovascular disease, all-cause mortality and deterioration in kidney function in persons with type 2 diabetes and microalbuminuria. Cardiovascular Diabetology, 2017, 16, 88.	2.7	41
41	High YKL-40 levels predict mortality in patients with type 2 diabetes. Diabetes Research and Clinical Practice, 2012, 96, 84-89.	1.1	39
42	Renal outcomes with aliskiren in patients with type 2 diabetes: a prespecified secondary analysis of the ALTITUDE randomised controlled trial. Lancet Diabetes and Endocrinology,the, 2016, 4, 309-317.	5.5	39
43	Effects of liraglutide on cardiovascular risk biomarkers in patients with type 2 diabetes and albuminuria: <scp>A</scp> subâ€analysis of a randomized, placeboâ€controlled, doubleâ€blind, crossover trial. Diabetes, Obesity and Metabolism, 2017, 19, 901-905.	2.2	39
44	Soluble Urokinase Plasminogen Activator Receptor Predicts Cardiovascular Events, Kidney Function Decline, and Mortality in Patients With Type 1 Diabetes. Diabetes Care, 2019, 42, 1112-1119.	4.3	38
45	Prevention of microalbuminuria using early intervention with renin-angiotensin system inhibitors in patients with type 2 diabetes: A systematic review. JRAAS - Journal of the Renin-Angiotensin-Aldosterone System, 2016, 17, 147032031665204.	1.0	36
46	Prognosis and treatment of diabetic nephropathy: Recent advances and perspectives. Nephrologie Et Therapeutique, 2018, 14, S31-S37.	0.2	35
47	Effect of large weight reductions on measured and estimated kidney function. BMC Nephrology, 2017, 18, 52.	0.8	34
48	Optimal dose of lisinopril for renoprotection in type 1 diabetic patients with diabetic nephropathy: a randomised crossover trial. Diabetologia, 2009, 52, 46-49.	2.9	32
49	Optimal antiproteinuric dose of aliskiren in type 2 diabetes mellitus: a randomised crossover trial. Diabetologia, 2010, 53, 1576-1580.	2.9	30
50	24-hour central aortic systolic pressure and 24-hour central pulse pressure are related to diabetic complications in type 1 diabetes – a cross-sectional study. Cardiovascular Diabetology, 2013, 12, 122.	2.7	30
51	Changes in Albuminuria Predict Cardiovascular and Renal Outcomes in Type 2 Diabetes: A Post Hoc Analysis of the LEADER Trial. Diabetes Care, 2021, 44, 1020-1026.	4.3	30
52	Predicting albuminuria response to spironolactone treatment with urinary proteomics in patients with type 2 diabetes and hypertension. Nephrology Dialysis Transplantation, 2018, 33, gfw406.	0.4	29
53	Growth differentiation factor-15 and fibroblast growth factor-23 are associated with mortality in type 2 diabetes $\hat{a} \in An$ observational follow-up study. PLoS ONE, 2018, 13, e0196634.	1.1	29
54	The effects of dapagliflozin, metformin or exercise on glycaemic variability in overweight or obese individuals with prediabetes (the PRE-D Trial): a multi-arm, randomised, controlled trial. Diabetologia, 2021, 64, 42-55.	2.9	29

#	Article	IF	CITATIONS
55	Incidence of Ketoacidosis in the Danish Type 2 Diabetes Population Before and After Introduction of Sodium–Glucose Cotransporter 2 Inhibitors—A Nationwide, Retrospective Cohort Study, 1995–2014. Diabetes Care, 2017, 40, e57-e58.	4.3	26
56	Ambulatory Tonometric Blood Pressure Measurements in Patients with Diabetes. Diabetes Technology and Therapeutics, 2012, 14, 453-456.	2.4	25
57	Development and validation of a brief, descriptive Danish pain questionnaire (BDDPQ). Acta Anaesthesiologica Scandinavica, 2004, 48, 486-490.	0.7	24
58	Elevated NT-proBNP and coronary calcium score in relation to coronary artery disease in asymptomatic type 2 diabetic patients with elevated urinary albumin excretion rate. Nephrology Dialysis Transplantation, 2011, 26, 3242-3249.	0.4	24
59	Plasma high-sensitivity troponin T predicts end-stage renal disease and cardiovascular and all-cause mortality in patients with type 1 diabetes and diabetic nephropathy. Kidney International, 2017, 92, 1242-1248.	2.6	24
60	Different patterns of secondâ€line treatment in type 2 diabetes after metformin monotherapy in Denmark, Finland, Norway and Sweden (D360 Nordic): A multinational observational study. Endocrinology, Diabetes and Metabolism, 2018, 1, e00036.	1.0	24
61	Characteristics of high―and lowâ€risk individuals in the <scp>PRIORITY</scp> study: urinary proteomics and mineralocorticoid receptor antagonism for prevention of diabetic nephropathy in Type 2 diabetes. Diabetic Medicine, 2018, 35, 1375-1382.	1.2	24
62	Effect of dapagliflozin on cardiac function in people with type 2 diabetes and albuminuria – A double blind randomized placebo-controlled crossover trial. Journal of Diabetes and Its Complications, 2020, 34, 107590.	1.2	24
63	Urinary renin and angiotensinogen in type 2 diabetes. Journal of Hypertension, 2013, 31, 1646-1652.	0.3	23
64	Urinary tubular biomarkers as predictors of kidney function decline, cardiovascular events and mortality in microalbuminuric type 2 diabetic patients. Acta Diabetologica, 2018, 55, 1143-1150.	1.2	23
65	Variability in response to albuminuriaâ€lowering drugs: true or random?. British Journal of Clinical Pharmacology, 2017, 83, 1197-1204.	1.1	22
66	Improving peptide relative quantification in MALDI-TOF MS for biomarker assessment. Proteomics, 2013, 13, 2967-2975.	1.3	21
67	Levels of NT-proBNP, markers of low-grade inflammation, and endothelial dysfunction during spironolactone treatment in patients with diabetic kidney disease. JRAAS - Journal of the Renin-Angiotensin-Aldosterone System, 2013, 14, 161-166.	1.0	21
68	Number and Frequency of Albuminuria Measurements in Clinical Trials in Diabetic Nephropathy. Clinical Journal of the American Society of Nephrology: CJASN, 2015, 10, 410-416.	2.2	21
69	Impact of aliskiren treatment on urinary aldosterone levels in patients with type 2 diabetes and nephropathy: an AVOID substudy. JRAAS - Journal of the Renin-Angiotensin-Aldosterone System, 2012, 13, 118-121.	1.0	20
70	A prediction of the renal and cardiovascular efficacy of aliskiren in ALTITUDE using short-term changes in multiple risk markers. European Journal of Preventive Cardiology, 2014, 21, 434-441.	0.8	19
71	Interpretation of HbA <sub>1c</sub> in primary care and potential influence of anaemia and chronic kidney disease: an analysis from the Copenhagen Primary Care Laboratory (CopLab) Database. Diabetic Medicine, 2018, 35, 1700-1706.	1.2	19
72	Mitigating risk of aldosterone in diabetic kidney disease. Current Opinion in Nephrology and Hypertension, 2020, 29, 145-151.	1.0	19

5

#	Article	IF	CITATIONS
73	Toe–brachial index as a predictor of cardiovascular disease and all-cause mortality in people with type 2 diabetes and microalbuminuria. Diabetologia, 2017, 60, 1883-1891.	2.9	18
74	Protocol for a randomised controlled trial of the effect of dapagliflozin, metformin and exercise on glycaemic variability, body composition and cardiovascular risk in prediabetes (the PRE-D Trial). BMJ Open, 2017, 7, e013802.	0.8	17
75	Urinary albumin and 8-oxo-7,8-dihydroguanosine as markers of mortality and cardiovascular disease during 19 years after diagnosis of type 2 diabetes – A comparative study of two markers to identify high risk patients. Redox Biology, 2017, 13, 363-369.	3.9	17
76	Uric acid is not associated with diabetic nephropathy and other complications in type 1 diabetes. Nephrology Dialysis Transplantation, 2019, 34, 659-666.	0.4	17
77	Urinary Proteomics and Precision Medicine for Chronic Kidney Disease: Current Status and Future Perspectives. Proteomics - Clinical Applications, 2019, 13, 1800176.	0.8	16
78	A Validated Prediction Model for End-Stage Kidney Disease in Type 1 Diabetes. Diabetes Care, 2021, 44, 901-907.	4.3	16
79	Drugâ€drug interaction between warfarin and statins: A Danish cohort study. British Journal of Clinical Pharmacology, 2021, 87, 694-699.	1.1	15
80	Pleiotropic effects of liraglutide treatment on renal risk factors in type 2 diabetes: Individual effects of treatment. Journal of Diabetes and Its Complications, 2017, 31, 162-168.	1.2	13
81	Omics research in diabetic kidney disease: new biomarker dimensions and new understandings?. Journal of Nephrology, 2020, 33, 931-948.	0.9	13
82	Assessment of the sublingual microcirculation with the GlycoCheck system: Reproducibility and examination conditions. PLoS ONE, 2020, 15, e0243737.	1.1	13
83	Comments on the 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases. European Heart Journal, 2020, 41, 328-328.	1.0	12
84	Linking Kidney and Cardiovascular Complications in Diabetes—Impact on Prognostication and Treatment: The 2019 Edwin Bierman Award Lecture. Diabetes, 2021, 70, 39-50.	0.3	12
85	Sequential RAAS Blockade: Is It Worth the Risk?. Advances in Chronic Kidney Disease, 2014, 21, 159-165.	0.6	11
86	Multifactorial intervention to prevent cardiovascular disease in patients with early rheumatoid arthritis: protocol for a multicentre randomised controlled trial. BMJ Open, 2016, 6, e009134.	0.8	11
87	Direct renin inhibition in chronic kidney disease. British Journal of Clinical Pharmacology, 2012, 76, n/a-n/a.	1.1	10
88	The influence of pharmaceutically induced weight changes on estimates of renal function: A patient-level pooled analysis of seven randomised controlled trials of glucose lowering medication. Journal of Diabetes and Its Complications, 2015, 29, 1146-1151.	1.2	10
89	Nâ€ŧerminal proâ€brain natriuretic peptide (NTâ€proBNP) predicts the cardioâ€renal response to aliskiren in patients with type 2 diabetes at high renal and cardiovascular risk. Diabetes, Obesity and Metabolism, 2018, 20, 2899-2904.	2.2	10
90	Study rationale and design of the EANITIATE study (EmpAgliflozin compared to NPH Insulin for sTeroId) Tj ETQqQ empagliflozin compared with NPH-insulin in patients with newly onset diabetes following initiation of glucocorticoid treatment. BMC Endocrine Disorders, 2020, 20, 86.	0 0 rgBT ( 0.9	Overlock 10 10

#	Article	IF	CITATIONS
91	Cardiovascular and renal outcomes by baseline albuminuria status and renal function: Results from the <scp>LEADER</scp> randomized trial. Diabetes, Obesity and Metabolism, 2020, 22, 2077-2088.	2.2	10
92	Postural Stability after Inguinal Herniorrhaphy under Local Infiltration Anaesthesia. The European Journal of Surgery, 2001, 167, 449-452.	1.0	9
93	Serum uric acid and progression of diabetic nephropathy in type 1 diabetes. Journal of Diabetes and Its Complications, 2018, 32, 470-473.	1.2	9
94	Application of urinary proteomics as possible risk predictor of renal and cardiovascular complications in patients with type 2-diabetes and microalbuminuria. Journal of Diabetes and Its Complications, 2018, 32, 1133-1140.	1.2	9
95	Pleiotropic effects of liraglutide in patients with type 2 diabetes and moderate renal impairment: Individual effects of treatment. Diabetes, Obesity and Metabolism, 2019, 21, 1261-1265.	2.2	9
96	No effects of dapagliflozin, metformin or exercise on plasma glucagon concentrations in individuals with prediabetes: A post hoc analysis from the randomized controlled <scp>PREâ€D</scp> trial. Diabetes, Obesity and Metabolism, 2021, 23, 530-539.	2.2	9
97	Risk factor management of type 2 diabetic patients in primary care in the Scandinavian countries between 2003 and 2015. Primary Care Diabetes, 2021, 15, 262-268.	0.9	9
98	Long-Term Effects of Irbesartan Treatment and Smoking on Nucleic Acid Oxidation in Patients With Type 2 Diabetes and Microalbuminuria. Diabetes Care, 2011, 34, 1192-1198.	4.3	8
99	Renin angiotensin system blockade reduces urinary levels of soluble urokinase plasminogen activator receptor (suPAR) in patients with type 2 diabetes. Journal of Diabetes and Its Complications, 2016, 30, 1440-1442.	1.2	8
100	The effect of liraglutide and sitagliptin on oxidative stress in persons with type 2 diabetes. Scientific Reports, 2021, 11, 10624.	1.6	8
101	Pooled Analysis of Multiple Crossover Trials To Optimize Individual Therapy Response to Renin-Angiotensin-Aldosterone System Intervention. Clinical Journal of the American Society of Nephrology: CJASN, 2017, 12, 1804-1813.	2.2	7
102	Visitâ€ŧoâ€visit variability of clinical risk markers in relation to longâ€ŧerm complications in type 1 diabetes. Diabetic Medicine, 2021, 38, e14459.	1.2	7
103	Is Time-Restricted Eating Safe in the Treatment of Type 2 Diabetes?—A Review of Intervention Studies. Nutrients, 2022, 14, 2299.	1.7	7
104	Incidence of New-Onset Type 2 Diabetes After Cancer: A Danish Cohort Study. Diabetes Care, 2022, 45, e105-e106.	4.3	7
105	Irbesartan treatment does not influence plasma levels of the advanced glycation end products NÂ(1-carboxymethyl)lysine and NÂ(1-carboxyethyl)lysine in patients with type 2 diabetes and microalbuminuria. A randomized controlled trial. Nephrology Dialysis Transplantation, 2011, 26, 3573-3577.	0.4	6
106	Improving the effectiveness of shortâ€ŧerm courses for multidisciplinary health care professionals. Practical Diabetes, 2015, 32, 180-185.	0.1	6
107	Effect of weight reductions on estimated kidney function: Post-hoc analysis of two randomized trials. Journal of Diabetes and Its Complications, 2017, 31, 1164-1168.	1.2	6
108	The effect of uric acid lowering treatment on albuminuria and renal function in Type 1 diabetes: a randomized clinical trial. Diabetic Medicine, 2018, 35, 392-393.	1.2	6

#	Article	IF	CITATIONS
109	Achieving a useful and personâ€centred diabetes consultation is a shared responsibility between diabetologists and people with diabetes: a qualitative study of perspectives from people with type 1 diabetes. Diabetic Medicine, 2021, 38, e14382.	1.2	6
110	Nordic Longitudinal Data from Electronic Medical Records and Full Population National Registers: Unique Opportunities for New Insights in Benefit of Diabetes Patients. Value in Health, 2015, 18, A726.	0.1	5
111	YKL-40 in dialysis patients: another candidate in the quest for useful biomarkers in nephrology. Kidney International, 2018, 93, 21-22.	2.6	5
112	Inflammation Leads the Way on the ROADMAP to Diabetic Kidney Disease. Kidney International Reports, 2019, 4, 1362-1365.	0.4	5
113	Irbesartan treatment does not influence plasma levels of the dicarbonyls methylglyoxal, glyoxal and 3â€deoxyglucosone in participants with type 2 diabetes and microalbuminuria: An IRMA2 subâ€study. Diabetic Medicine, 2021, 38, e14405.	1.2	5
114	Importance of standardizing renal outcomes in clinical trials: illustration by recent sodium glucose cotransporter 2 inhibitor studies. Kidney International, 2021, 99, 768-770.	2.6	5
115	A narrative review of new treatment options for chronic kidney disease in type 2 diabetes. Annals of Translational Medicine, 2021, 9, 716-716.	0.7	5
116	The importance of addressing multiple risk markers in type 2 diabetes: Results from the <scp>LEADER</scp> and <scp>SUSTAIN</scp> 6 trials. Diabetes, Obesity and Metabolism, 2022, 24, 281-288.	2.2	5
117	Copeptin and renal function decline, cardiovascular events and mortality in type 1 diabetes. Nephrology Dialysis Transplantation, 2020, , .	0.4	5
118	Impact of random variation in albuminuria and estimated glomerular filtration rate on patient enrolment and duration of clinical trials in nephrology. Diabetes, Obesity and Metabolism, 2022, 24, 983-990.	2.2	5
119	Sodium–glucose cotransporter 2 inhibitors as adjunct therapy for type 1 diabetes and the benefit on cardiovascular and renal disease evaluated by Steno risk engines. Journal of Diabetes and Its Complications, 2022, 36, 108257.	1.2	5
120	Higher Parathyroid Hormone Level Is Associated With Increased Arterial Stiffness in Type 1 Diabetes. Diabetes Care, 2017, 40, e32-e33.	4.3	4
121	Lipoprotein(a)and renal function decline, cardiovascular disease and mortality in type 2 diabetes and microalbuminuria. Journal of Diabetes and Its Complications, 2020, 34, 107593.	1.2	4
122	Improving frequency of urinary albumin testing in type 2 diabetes in primary care — An analysis of cross-sectional studies in Denmark. Primary Care Diabetes, 2021, 15, 1007-1011.	0.9	4
123	Impact of glycaemic control on the effect of direct renin inhibition in the AVOID study. JRAAS - Journal of the Renin-Angiotensin-Aldosterone System, 2012, 13, 250-253.	1.0	3
124	Discrepancy Between Tonometric Ambulatory and Cuffâ€Based Office Blood Pressure Measurements in Patients With Type 1 Diabetes. Journal of Clinical Hypertension, 2012, 14, 686-693.	1.0	3
125	Effects of RAS inhibitors on diabetic retinopathy. Lancet Diabetes and Endocrinology,the, 2015, 3, 315-316.	5.5	3
126	Increasing ESKD in Diabetes in the Land Down Under: What Can Be Done, We Must Wonder. American Journal of Kidney Diseases, 2019, 73, 293-296.	2.1	3

Frederik Persson

#	Article	IF	CITATIONS
127	Oral treatment of glucocorticoidâ€induced diabetes mellitus: A systematic review. International Journal of Clinical Practice, 2020, 74, e13529.	0.8	3
128	Endothelial glycocalyx and cardio-renal risk factors in type 1 diabetes. PLoS ONE, 2021, 16, e0254859.	1.1	3
129	Acute and Long-Term Treatment With Dapagliflozin and Association With Serum Soluble Urokinase Plasminogen Activator Receptor. Frontiers in Pharmacology, 2022, 13, 799915.	1.6	3
130	Improvements in albuminuria and chronic kidney disease progression with the appetite suppressant lorcaserin. Kidney International, 2019, 95, 1287-1288.	2.6	2
131	A large remaining potential in lipidâ€lowering drug treatment in the type 2 diabetes population: A Danish nationwide cohort study. Diabetes, Obesity and Metabolism, 2021, 23, 2354-2363.	2.2	2
132	Empagliflozin reduces albuminuria—a promise for better cardiorenal protection from the EMPA-REG OUTCOME trial. Annals of Translational Medicine, 2017, 5, 478-478.	0.7	1
133	FC 058THE IMPORTANCE OF ADDRESSING MULTIPLE RISK MARKERS IN TYPE 2 DIABETES: RESULTS FROM THE LEADER AND SUSTAIN 6 TRIALS. Nephrology Dialysis Transplantation, 2021, 36, .	0.4	1
134	What Have We Learned so Far From the Use of Sodium-Glucose Cotransporter 2 Inhibitors in Clinical Practice?. Advances in Chronic Kidney Disease, 2021, 28, 290-297.	0.6	1
135	Aldosterone Blockade Added to Renin-Angiotensin System Blockade to Reduce Albuminuria—A Path for Improved Renoprotection?. American Journal of the Medical Sciences, 2018, 355, 407-408.	0.4	Ο
136	Title is missing!. , 2020, 15, e0243737.		0
137	Title is missing!. , 2020, 15, e0243737.		0
138	Title is missing!. , 2020, 15, e0243737.		0
139	Title is missing!. , 2020, 15, e0243737.		0
140	Title is missing!. , 2020, 15, e0243737.		0
141	Title is missing!. , 2020, 15, e0243737.		0
142	Title is missing!. , 2020, 15, e0243737.		0
143	Title is missing!. , 2020, 15, e0243737.		0