
Mona Koosha

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5957846/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Molecular and Biochemical Detection of Insecticide Resistance in the <i>Leishmania</i> Vector, <i>Phlebotomus papatasi</i> (Diptera: Psychodidae) to Dichlorodiphenyltrichloroethane and Pyrethroids, in Central Iran. Journal of Medical Entomology, 2022, 59, 1347-1354.	0.9	2
2	Aerobic Bacterial Community of the Rice Striped Stem Borer: A Step towards Finding Bacterial Candidates for Paratransgenesis and RNAi Control. International Journal of Plant Animal and Environmental Sciences, 2021, 11, .	0.2	0
3	An integrated overview of the bacterial flora composition of Hyalomma anatolicum, the main vector of CCHF. PLoS Neglected Tropical Diseases, 2021, 15, e0009480.	1.3	9
4	Hyalomma spp. ticks and associated Anaplasma spp. and Ehrlichia spp. on the Iran-Pakistan border. Parasites and Vectors, 2021, 14, 469.	1.0	17
5	Molecular detection of Dirofilaria spp. and host blood-meal identification in the Simulium turgaicum complex (Diptera: Simuliidae) in the Aras River Basin, northwestern Iran. Parasites and Vectors, 2020, 13, 548.	1.0	7
6	Identification of Forensically Important Flesh Flies Using the Cytochrome C Oxidase Subunits I and II Genes. Journal of Medical Entomology, 2019, 56, 1253-1259.	0.9	13
7	Effect of Serratia AS1 (Enterobacteriaceae: Enterobacteriales) on the Fitness of Culex pipiens (Diptera:) Tj ETQq1	1 0.7843	l4.rgBT /Ov
8	Delivery of a Genetically Marked Serratia AS1 to Medically Important Arthropods for Use in RNAi and Paratransgenic Control Strategies. Microbial Ecology, 2019, 78, 185-194.	1.4	29
9	Aerobic midgut microbiota of sand fly vectors of zoonotic visceral leishmaniasis from northern Iran, a step toward finding potential paratransgenic candidates. Parasites and Vectors, 2019, 12, 10.	1.0	25
10	Effect of Meteorological Factors on Species Composition and Their Host Preference, Seasonal Prevalence and Infection Status to Crimean-Congo Haemorrhagic Fever in Iran. Journal of Arthropod-Borne Diseases, 2019, 13, 268-283.	0.9	9
11	wsp-based analysis of <i>Wolbachia</i> strains associated with <i>Phlebotomus papatasi</i> and <i>P. sergenti</i> (Diptera: Psychodidae) main cutaneous leishmaniasis vectors, introduction of a new subgroup wSerg. Pathogens and Global Health, 2018, 112, 152-160.	1.0	25
12	Sequence analysis of mtDNA COI barcode region revealed three haplotypes within Culex pipiens assemblage. Experimental Parasitology, 2017, 181, 102-110.	0.5	12
13	Molecular Characterization and Phylogenetic Congruence of (Tricoptera: Hydropsychidae) Using Mitochondrial and Nuclear Markers. Journal of Arthropod-Borne Diseases, 2017, 11, 60-77.	0.9	2
14	Assessing the Fauna of Aquatic Insects for Possible Use for Malaria Vector Control in Large River, Central Iran. Acta Medica Iranica, 2015, 53, 523-32.	0.8	9
15	Lectin Activity in Gut Extract of Culex pipiens. Journal of Arthropod-Borne Diseases, 2013, 7, 23-30.	0.9	0