

Patrick Cyriel Souverein

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5961779/patrick-cyriel-souverein-publications-by-citations.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

134 papers	2,325 citations	26 h-index	41 g-index
139 ext. papers	2,744 ext. citations	4.3 avg, IF	4.78 L-index

#	Paper	IF	Citations
134	Incidence of fractures among epilepsy patients: a population-based retrospective cohort study in the General Practice Research Database. <i>Epilepsia</i> , 2005 , 46, 304-10	6.4	167
133	Antipsychotic drug use and risk of pneumonia in elderly people. <i>Journal of the American Geriatrics Society</i> , 2008 , 56, 661-6	5.6	145
132	Increased bleeding risk with concurrent use of selective serotonin reuptake inhibitors and coumarins. <i>Archives of Internal Medicine</i> , 2008 , 168, 180-5		98
131	Use of antidepressant drugs and risk of osteoporotic and non-osteoporotic fractures. <i>Bone</i> , 2010 , 47, 604-9	4.7	76
130	Exposure to benzodiazepines (anxiolytics, hypnotics and related drugs) in seven European electronic healthcare databases: a cross-national descriptive study from the PROTECT-EU Project. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 56-65	2.6	73
129	Construction of drug treatment episodes from drug-dispensing histories is influenced by the gap length. <i>Journal of Clinical Epidemiology</i> , 2010 , 63, 422-7	5.7	69
128	Statin treatment and the risk of recurrent pulmonary embolism. <i>European Heart Journal</i> , 2013 , 34, 1800-6	5.5	63
127	Risk of aplastic anemia in patients using antiepileptic drugs. <i>Epilepsia</i> , 2006 , 47, 1232-6	6.4	60
126	Is aspirin useful in patients on lithium? A pharmacoepidemiological study related to bipolar disorder. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2010 , 82, 9-14	2.8	56
125	Use of oral glucocorticoids and the risk of pulmonary embolism: a population-based case-control study. <i>Chest</i> , 2013 , 143, 1337-1342	5.3	52
124	Increased use of antimicrobial agents and hospital admission for infections in patients with primary adrenal insufficiency: a cohort study. <i>European Journal of Endocrinology</i> , 2013 , 168, 609-14	6.5	47
123	Hyponatraemia as an adverse drug reaction of antipsychotic drugs: a case-control study in VigiBase. <i>Drug Safety</i> , 2010 , 33, 569-78	5.1	44
122	Prevalence of antibiotic use: a comparison across various European health care data sources. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 11-20	2.6	40
121	Trends in incidence and characteristics of children, adolescents, and adults initiating immediate- or extended-release methylphenidate or atomoxetine in the Netherlands during 2001-2006. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2010 , 20, 55-61	2.9	35
120	Use of organic nitrates and the risk of hip fracture: a population-based case-control study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 1924-31	5.6	34
119	Increasing trends in the incidence and prevalence rates of type 1 diabetes among children and adolescents in the Netherlands. <i>Pediatric Diabetes</i> , 2016 , 17, 44-52	3.6	31
118	Less discontinuation of ADHD drug use since the availability of long-acting ADHD medication in children, adolescents and adults under the age of 45 years in the Netherlands. <i>ADHD Attention Deficit and Hyperactivity Disorders</i> , 2010 , 2, 213-20	3.1	31

117	Antiepileptic drug use in seven electronic health record databases in Europe: a methodologic comparison. <i>Epilepsia</i> , 2014 , 55, 666-673	6.4	30
116	Association between ADHD drug use and injuries among children and adolescents. <i>European Child and Adolescent Psychiatry</i> , 2014 , 23, 95-102	5.5	30
115	Sudden cardiac arrest associated with use of a non-cardiac drug that reduces cardiac excitability: evidence from bench, bedside, and community. <i>European Heart Journal</i> , 2013 , 34, 1506-16	9.5	30
114	Hip/femur fractures associated with the use of benzodiazepines (anxiolytics, hypnotics and related drugs): a methodological approach to assess consistencies across databases from the PROTECT-EU project. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 66-78	2.6	30
113	Incidence of direct oral anticoagulant use in patients with nonvalvular atrial fibrillation and characteristics of users in 6 European countries (2008-2015): A cross-national drug utilization study. <i>British Journal of Clinical Pharmacology</i> , 2019 , 85, 2524-2539	3.8	29
112	Adverse Drug Reactions Reported With Cholinesterase Inhibitors: An Analysis of 16 Years of Individual Case Safety Reports From VigiBase. <i>Annals of Pharmacotherapy</i> , 2015 , 49, 1197-206	2.9	28
111	Effect of oral antiplatelet agents on major bleeding in users of coumarins. <i>Thrombosis and Haemostasis</i> , 2008 , 100, 1076-83	7	28
110	Evaluation of risk profiles for gastrointestinal and cardiovascular adverse effects in nonselective NSAID and COX-2 inhibitor users: a cohort study using pharmacy dispensing data in The Netherlands. <i>Drug Safety</i> , 2008 , 31, 143-58	5.1	27
109	Reimbursement restriction and moderate decrease in benzodiazepine use in general practice. <i>Annals of Family Medicine</i> , 2012 , 10, 42-9	2.9	26
108	Society already achieves economic benefits from generic substitution but fails to do the same for therapeutic substitution. <i>British Journal of Clinical Pharmacology</i> , 2007 , 64, 680-5	3.8	25
107	Angiotensin-converting enzyme inhibitor use and protection against pneumonia in patients with diabetes. <i>Journal of Hypertension</i> , 2007 , 25, 235-9	1.9	25
106	Long-term use of 5 α -reductase inhibitors and the risk of male breast cancer. <i>Cancer Causes and Control</i> , 2014 , 25, 1577-82	2.8	24
105	Association between statin use and lupus-like syndrome using spontaneous reports. <i>Seminars in Arthritis and Rheumatism</i> , 2011 , 41, 373-81	5.3	24
104	Determinants of angiotensin-converting enzyme inhibitor (ACEI) intolerance and angioedema in the UK Clinical Practice Research Datalink. <i>British Journal of Clinical Pharmacology</i> , 2016 , 82, 1647-1659	3.8	23
103	Drug-induced thrombocytopenia: a population study. <i>Drug Safety</i> , 2006 , 29, 713-21	5.1	23
102	Risk of major bleeding among users of direct oral anticoagulants combined with interacting drugs: A population-based nested case-control study. <i>British Journal of Clinical Pharmacology</i> , 2020 , 86, 1150-1164	3.8	22
101	Statin-associated polymyalgia rheumatica. An analysis using WHO global individual case safety database: a case/non-case approach. <i>PLoS ONE</i> , 2012 , 7, e41289	3.7	22
100	Are ECG monitoring recommendations before prescription of QT-prolonging drugs applied in daily practice? The example of haloperidol. <i>Pharmacoepidemiology and Drug Safety</i> , 2015 , 24, 701-8	2.6	21

99	Discontinuation of cholinesterase inhibitor treatment and determinants thereof in the Netherlands: A retrospective cohort study. <i>Drugs and Aging</i> , 2010 , 27, 663-75	4.7	21
98	Understanding inconsistency in the results from observational pharmacoepidemiological studies: the case of antidepressant use and risk of hip/femur fractures. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 88-102	2.6	21
97	Association between hospitalization and discontinuity of medication therapy used in the community setting in the Netherlands. <i>Annals of Pharmacotherapy</i> , 2008 , 42, 933-9	2.9	20
96	Gastrointestinal toxicity among patients taking selective COX-2 inhibitors or conventional NSAIDs, alone or combined with proton pump inhibitors: a case-control study. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 1141-1148	2.6	19
95	Chronic comorbidities in children with type 1 diabetes: a population-based cohort study. <i>Archives of Disease in Childhood</i> , 2015 , 100, 763-8	2.2	19
94	Medication adherence among gout patients initiated allopurinol: a retrospective cohort study in the Clinical Practice Research Datalink (CPRD). <i>Rheumatology</i> , 2018 , 57, 1641-1650	3.9	19
93	Linking community pharmacy dispensing data to prescribing data of general practitioners. <i>BMC Medical Informatics and Decision Making</i> , 2006 , 6, 18	3.6	18
92	Reported Adverse Drug Reactions in Children and Adolescents Treated with Antipsychotics. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2019 , 29, 124-132	2.9	17
91	The impact of age and sex on the reporting of cough and angioedema with renin-angiotensin system inhibitors: a case/noncase study in Vigibase. <i>Fundamental and Clinical Pharmacology</i> , 2017 , 31, 676-684	3.1	17
90	Time-trends in treatment and cardiovascular events in patients with heart failure: a pharmacosurveillance study. <i>European Journal of Heart Failure</i> , 2011 , 13, 489-95	12.3	17
89	Undertreatment of hypertension and hypercholesterolaemia in children and adolescents with type 1 diabetes: long-term follow-up on time trends in the occurrence of cardiovascular disease, risk factors and medications use. <i>British Journal of Clinical Pharmacology</i> , 2018 , 84, 776-785	3.8	17
88	Risk of Nephrotic Syndrome for Non-Steroidal Anti-Inflammatory Drug Users. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2019 , 14, 1355-1362	6.9	16
87	Increased risk of sudden cardiac arrest in obstructive pulmonary disease: a case-control study. <i>PLoS ONE</i> , 2013 , 8, e65638	3.7	16
86	Methodological comparison of marginal structural model, time-varying Cox regression, and propensity score methods: the example of antidepressant use and the risk of hip fracture. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 114-21	2.6	16
85	Predictive value of amplitude spectrum area of ventricular fibrillation waveform in patients with acute or previous myocardial infarction in out-of-hospital cardiac arrest. <i>Resuscitation</i> , 2017 , 120, 125-134	4	15
84	Inhaled Corticosteroid Adherence Patterns in a Longitudinal Asthma Cohort. <i>Journal of Allergy and Clinical Immunology: in Practice</i> , 2017 , 5, 448-456.e2	5.4	15
83	Treatment with rivastigmine or galantamine and risk of urinary incontinence: results from a Dutch database study. <i>Pharmacoepidemiology and Drug Safety</i> , 2015 , 24, 276-85	2.6	15
82	Changes in medication associated with epilepsy-related hospitalisation: a case-crossover study. <i>Pharmacoepidemiology and Drug Safety</i> , 2007 , 16, 189-96	2.6	14

81	Non-steroidal anti-inflammatory drugs and risk of pulmonary embolism. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 635-42	2.6	13
80	Prescription patterns of angiotensin-converting enzyme inhibitors for various indications: A UK population-based study. <i>British Journal of Clinical Pharmacology</i> , 2018 , 84, 2365-2372	3.8	13
79	Differences in VigiBase reporting of aminoglycoside and capreomycin-suspected ototoxicity during tuberculosis treatment. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 1-8	2.6	12
78	Use of rivastigmine or galantamine and risk of adverse cardiac events: a database study from the Netherlands. <i>American Journal of Geriatric Pharmacotherapy</i> , 2012 , 10, 373-80		12
77	Exploring the Association between Monoclonal Antibodies and Depression and Suicidal Ideation and Behavior: A VigiBase Study. <i>Drug Safety</i> , 2019 , 42, 887-895	5.1	12
76	Change in prescription pattern as a potential marker for adverse drug reactions of angiotensin converting enzyme inhibitors. <i>International Journal of Clinical Pharmacy</i> , 2015 , 37, 1095-103	2.3	11
75	Long term patterns of use after initiation of oral antidiabetic drug therapy. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 351-8	2.6	11
74	Drug-related nephrotoxic and ototoxic reactions : a link through a predictive mechanistic commonality. <i>Drug Safety</i> , 2008 , 31, 877-84	5.1	11
73	Non-Publication Is Common among Phase 1, Single-Center, Not Prospectively Registered, or Early Terminated Clinical Drug Trials. <i>PLoS ONE</i> , 2016 , 11, e0167709	3.7	11
72	Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	11
71	Population-based cohort study of anti-infective medication use before and after the onset of type 1 diabetes in children and adolescents. <i>Antimicrobial Agents and Chemotherapy</i> , 2014 , 58, 4666-74	5.9	10
70	Non-adherence to inhaled corticosteroids and the risk of asthma exacerbations in children. <i>Patient Preference and Adherence</i> , 2016 , 10, 531-8	2.4	10
69	Case-only designs for studying the association of antidepressants and hip or femur fracture. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 103-13	2.6	10
68	Risk of myocardial infarction in patients with atrial fibrillation using vitamin K antagonists, aspirin or direct acting oral anticoagulants. <i>British Journal of Clinical Pharmacology</i> , 2017 , 83, 1835-1843	3.8	9
67	Differences in ADHD medication usage patterns in children and adolescents from different cultural backgrounds in the Netherlands. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015 , 50, 1153-62	4.5	9
66	Reduced pre-hospital and in-hospital survival rates after out-of-hospital cardiac arrest of patients with type-2 diabetes mellitus: an observational prospective community-based study. <i>Europace</i> , 2015 , 17, 753-60	3.9	9
65	Identification of exacerbations in obstructive lung disease through biomarkers. <i>Biomarkers</i> , 2009 , 14, 523-8	2.6	9
64	Instrumental variable analysis as a complementary analysis in studies of adverse effects: venous thromboembolism and second-generation versus third-generation oral contraceptives. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25, 317-24	2.6	9

63	Differential effects on out-of-hospital cardiac arrest of dihydropyridines: real-world data from population-based cohorts across two European countries. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2020 , 6, 347-355	6.4	9
62	Primary endpoint discrepancies were found in one in ten clinical drug trials. Results of an inception cohort study. <i>Journal of Clinical Epidemiology</i> , 2017 , 89, 199-208	5.7	8
61	Smoke-free legislation and the incidence of paediatric respiratory infections and wheezing/asthma: interrupted time series analyses in the four UK nations. <i>Scientific Reports</i> , 2015 , 5, 15246	4.9	8
60	Effects of glucocorticoids on the neutrophil count: a cohort study among hospitalized patients. <i>Pulmonary Pharmacology and Therapeutics</i> , 2010 , 23, 129-34	3.5	8
59	Non-selective vs. selective beta-blocker treatment and the risk of thrombo-embolic events in patients with heart failure. <i>European Journal of Heart Failure</i> , 2011 , 13, 220-6	12.3	8
58	Prior outpatient antibacterial therapy as prognostic factor for mortality in hospitalized pneumonia patients. <i>Respiratory Medicine</i> , 2006 , 100, 1342-8	4.6	8
57	The Relationship Between Real-World Inhaled Corticosteroid Adherence and Asthma Outcomes: A Multilevel Approach. <i>Journal of Allergy and Clinical Immunology: in Practice</i> , 2020 , 8, 626-634	5.4	8
56	Safety of clozapine use during pregnancy: Analysis of international pharmacovigilance data. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 725-735	2.6	7
55	Unmeasured confounding in pharmacoepidemiology. <i>Annals of Epidemiology</i> , 2016 , 26, 85-6	6.4	7
54	Recruitment failure and futility were the most common reasons for discontinuation of clinical drug trials. Results of a nationwide inception cohort study in the Netherlands. <i>Journal of Clinical Epidemiology</i> , 2017 , 88, 140-147	5.7	7
53	Patterns of antiplatelet drug use after a first myocardial infarction during a 10-year period. <i>British Journal of Clinical Pharmacology</i> , 2017 , 83, 632-641	3.8	7
52	Exploring the role of low-frequency and rare exonic variants in alcohol and tobacco use. <i>Drug and Alcohol Dependence</i> , 2018 , 188, 94-101	4.9	7
51	Antipsychotic drug use associated with urinary tract infections in older women. <i>Maturitas</i> , 2017 , 98, 46-50		6
50	Association Between Urinary Tract Infections and Antipsychotic Drug Use in Older Adults. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 296-301	1.7	6
49	The impact of serum potassium-influencing antihypertensive drugs on the risk of out-of-hospital cardiac arrest: A case-control study. <i>British Journal of Clinical Pharmacology</i> , 2017 , 83, 2541-2548	3.8	6
48	Age as a determinant of instability of serum lithium concentrations. <i>Therapeutic Drug Monitoring</i> , 2013 , 35, 643-8	3.2	6
47	Statin use and markers of immunity in the Doetinchem cohort study. <i>PLoS ONE</i> , 2013 , 8, e77587	3.7	6
46	Continuation of angiotensin converting enzyme inhibitor therapy, in spite of occurrence of angioedema. <i>International Journal of Cardiology</i> , 2015 , 201, 644-5	3.2	5

45	Do the benefits of statins outweigh their drawbacks? Assessing patients' trade-off preferences with conjoint analysis. <i>International Journal of Cardiology</i> , 2014 , 176, 1220-2	3.2	5
44	The association between prescription change frequency, chronic disease score and hospital admissions: a case control study. <i>BMC Pharmacology & Toxicology</i> , 2013 , 14, 39	2.6	5
43	Occurrence and determinants of selective reporting of clinical drug trials: design of an inception cohort study. <i>BMJ Open</i> , 2015 , 5, e007827	3	5
42	Differences in lithium use patterns in the Netherlands: comparing middle-aged and older patients in a database study. <i>American Journal of Geriatric Pharmacotherapy</i> , 2012 , 10, 193-200		5
41	Testing bias in clinical databases: methodological considerations. <i>Emerging Themes in Epidemiology</i> , 2010 , 7, 2	3.9	5
40	Cardiovascular medication use and cardiovascular disease in children and adolescents with type 1 diabetes: a population-based cohort study. <i>Pediatric Diabetes</i> , 2016 , 17, 433-40	3.6	5
39	Compliance with the guidelines for laboratory monitoring of patients treated with lithium: A retrospective follow-up study among ambulatory patients in the Netherlands. <i>Bipolar Disorders</i> , 2019 , 21, 419-427	3.8	5
38	Risk of Hospitalization for Hypoglycemia in Older Patients with Diabetes Using Antipsychotic Drugs. <i>American Journal of Geriatric Psychiatry</i> , 2015 , 23, 1144-53	6.5	4
37	Medication changes prior to hospitalization for obstructive lung disease: a case-crossover study. <i>Annals of Pharmacotherapy</i> , 2010 , 44, 267-73	2.9	4
36	Measuring exacerbations in obstructive lung disease. <i>Pharmacoepidemiology and Drug Safety</i> , 2010 , 19, 367-74	2.6	4
35	Disease history and medication use as risk factors for the clinical manifestation of type 1 diabetes in children and young adults: an explorative case control study. <i>PLoS ONE</i> , 2014 , 9, e87408	3.7	4
34	Sulfonylurea antidiabetics are associated with lower risk of out-of-hospital cardiac arrest: Real-world data from a population-based study. <i>British Journal of Clinical Pharmacology</i> , 2021 , 87, 3588-3598	3.8	4
33	Major bleeding in users of direct oral anticoagulants in atrial fibrillation: A pooled analysis of results from multiple population-based cohort studies. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 1339-1352	2.6	4
32	Risk of myocardial infarction associated with non-steroidal anti-inflammatory drugs: Impact of additional confounding control for variables collected from self-reported data. <i>Journal of Clinical Pharmacy and Therapeutics</i> , 2019 , 44, 623-631	2.2	3
31	Asthma related medication use and exacerbations in children and adolescents with type 1 diabetes. <i>Pediatric Pulmonology</i> , 2016 , 51, 1113-1121	3.5	3
30	Prescriber adoption of newly approved selective COX-2 inhibitors. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 1168-74	2.6	3
29	The association between receptor binding affinity and metabolic side effect profile of antipsychotics and major cardio- and cerebrovascular events: A case/non-case study using VigiBase. <i>European Neuropsychopharmacology</i> , 2020 , 35, 30-38	1.2	3
28	The Number of Concomitant Drugs and the Safety of Direct Oral Anticoagulants in Routine Care Patients with Atrial Fibrillation. <i>TH Open</i> , 2020 , 4, e417-e426	2.7	3

27	Comparing risk of major bleeding between users of different oral anticoagulants in patients with nonvalvular atrial fibrillation. <i>British Journal of Clinical Pharmacology</i> , 2021 , 87, 988-1000	3.8	3
26	Low-dose oral glucocorticoid therapy and risk of osteoporotic fractures in patients with rheumatoid arthritis: a cohort study using the Clinical Practice Research Datalink. <i>Rheumatology</i> , 2021 ,	3.9	3
25	Out-of-hospital cardiac arrest and differential risk of cardiac and non-cardiac QT-prolonging drugs in 37 000 cases. <i>British Journal of Clinical Pharmacology</i> , 2021 ,	3.8	3
24	Cyclo-oxygenase selectivity and chemical groups of nonsteroidal anti-inflammatory drugs and the frequency of reporting hypersensitivity reactions: a case/noncase study in VigiBase. <i>Fundamental and Clinical Pharmacology</i> , 2019 , 33, 589-600	3.1	2
23	Comparative effectiveness and safety of direct oral anticoagulants versus warfarin in UK patients with atrial fibrillation and type 2 diabetes: A retrospective cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 1293-1320	2.6	2
22	Reporting Patterns of Sialorrhea Comparing Users of Clozapine to Users of Other Antipsychotics: A Disproportionality Analysis Using VigiBase. <i>Journal of Clinical Psychopharmacology</i> , 2020 , 40, 283-286	1.7	2
21	Ventricular fibrillation waveform characteristics in out-of-hospital cardiac arrest and cardiovascular medication use. <i>Resuscitation</i> , 2020 , 151, 173-180	4	2
20	The risk of new fragility fractures in patients with chronic kidney disease and hip fracture-a population-based cohort study in the UK. <i>Osteoporosis International</i> , 2020 , 31, 1487-1497	5.3	2
19	Statins After Myocardial Infarction in the Oldest: A Cohort Study in the Clinical Practice Research Datalink Database. <i>Journal of the American Geriatrics Society</i> , 2020 , 68, 329-336	5.6	2
18	Psychiatric medication use before and after the onset of type 1 diabetes in children and adolescents: A population-based cohort study. <i>Pediatric Diabetes</i> , 2018 , 19, 121-128	3.6	1
17	Non-steroidal anti-inflammatory drugs and the risk of out-of-hospital cardiac arrest: a case-control study. <i>Europace</i> , 2019 , 21, 99-105	3.9	1
16	Rivaroxaban was found to be noninferior to warfarin in routine clinical care: A retrospective noninferiority cohort replication study. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 1263-1272	2.6	1
15	Impact of anticoagulant exposure misclassification on the bleeding risk of direct oral anticoagulants. <i>British Journal of Clinical Pharmacology</i> , 2021 , 87, 3508-3517	3.8	1
14	Prescribers' compliance with summary of product characteristics of dabigatran, rivaroxaban and apixaban-A European comparative drug utilization study. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2021 , 128, 440-454	3.1	1
13	Response to: Correspondence on Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study by Zheng. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	1
12	Hydroxyzine Initiation Following Drug Safety Advisories on Cardiac Arrhythmias in the UK and Canada: A Longitudinal Cohort Study.. <i>Drug Safety</i> , 2022 , 1	5.1	1
11	Mapping the risk of infections in patients with multiple sclerosis: A multi-database study in the United Kingdom Clinical Practice Research Datalink GOLD and Aurum.. <i>Multiple Sclerosis Journal</i> , 2022 , 13524585221094218	5	1
10	Effects of corticosteroid use on readmission in obstructive lung disease. <i>Respiratory Medicine</i> , 2010 , 104, 211-8	4.6	0

9	Statins After Ischemic Stroke in the Oldest: A Cohort Study Using the Clinical Practice Research Datalink Database. <i>Stroke</i> , 2021 , 52, 1244-1252	6.7	o
8	Angiotensin-converting enzyme inhibitors and angiotensin receptor blockers and risk of depression among older people with hypertension.. <i>Journal of Psychopharmacology</i> , 2022 , 2698811221082470	4.6	o
7	Impulse control disorders associated with dopaminergic drugs: A disproportionality analysis using vigibase.. <i>European Neuropsychopharmacology</i> , 2022 , 58, 30-38	1.2	o
6	Use of parenteral glucocorticoids and the risk of new onset type 2 diabetes mellitus: A case-control study. <i>Diabetes Research and Clinical Practice</i> , 2018 , 139, 100-106	7.4	
5	Interpretation of the association between beta-blockers and hip/femur fracture. <i>American Journal of Nephrology</i> , 2012 , 35, 101-2	4.6	
4	Association of beta-blockers and first-registered heart rhythm in out-of-hospital cardiac arrest: real-world data from population-based cohorts across two European countries. <i>Europace</i> , 2020 , 22, 1206-1215	3.9	
3	Response to: Correspondence on Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort studyTby Gong and Zhang and Lin. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	
2	Effect of metabolic genetic variants on long-term disease comorbidity in patients with type 2 diabetes. <i>Scientific Reports</i> , 2021 , 11, 2794	4.9	
1	Characteristics of patients with lung cancer in clinical practice and their potential eligibility for clinical trials evaluating tyrosine kinase inhibitors or immune checkpoint inhibitors.. <i>Cancer Epidemiology</i> , 2022 , 78, 102149	2.8	