Guido Miccinesi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5962192/publications.pdf

Version: 2024-02-01

196 papers 8,455 citations

³⁸⁷⁴² 50 h-index

83 g-index

205 all docs 205 docs citations

205 times ranked 7732 citing authors

#	Article	IF	CITATIONS
1	Integration of oncology and palliative care: a Lancet Oncology Commission. Lancet Oncology, The, 2018, 19, e588-e653.	10.7	452
2	The Impact of Mammographic Screening on Breast Cancer Mortality in Europe: A Review of Observational Studies. Journal of Medical Screening, 2012, 19, 14-25.	2.3	348
3	Overdiagnosis in Mammographic Screening for Breast Cancer in Europe: A Literature Review. Journal of Medical Screening, 2012, 19, 42-56.	2.3	338
4	Are There Differences in the Prevalence of Palliative Care-Related Problems in People Living With Advanced Cancer and Eight Non-Cancer Conditions? A Systematic Review. Journal of Pain and Symptom Management, 2014, 48, 660-677.	1,2	322
5	Actual and preferred place of death of cancer patients. Results from the Italian survey of the dying of cancer (ISDOC). Journal of Epidemiology and Community Health, 2006, 60, 412-416.	3.7	193
6	Which Patients With Cancer Die at Home? A Study of Six European Countries Using Death Certificate Data. Journal of Clinical Oncology, 2010, 28, 2267-2273.	1.6	193
7	Validation of the Consensus-Definition for Cancer Cachexia and evaluation of a classification model—a study based on data from an international multicentre project (EPCRC-CSA). Annals of Oncology, 2014, 25, 1635-1642.	1.2	190
8	Edmonton symptom assessment scale: Italian validation in two palliative care settings. Supportive Care in Cancer, 2006, 14, 30-37.	2.2	177
9	Summary of the Evidence of Breast Cancer Service Screening Outcomes in Europe and First Estimate of the Benefit and Harm Balance Sheet. Journal of Medical Screening, 2012, 19, 5-13.	2.3	172
10	Continuous Deep Sedation: Physicians' Experiences in Six European Countries. Journal of Pain and Symptom Management, 2006, 31, 122-129.	1.2	170
11	International study of the place of death of people with cancer: a population-level comparison of 14 countries across 4 continents using death certificate data. British Journal of Cancer, 2015, 113, 1397-1404.	6.4	158
12	Breast Cancer Mortality in Mammographic Screening in Europe: A Review of Incidence-Based Mortality Studies. Journal of Medical Screening, 2012, 19, 33-41.	2.3	152
13	Place of death in the population dying from diseases indicative of palliative care need: a cross-national population-level study in 14 countries. Journal of Epidemiology and Community Health, 2016, 70, 17-24.	3.7	150
14	Physicians' attitudes towards end-of-life decisions: a comparison between seven countries. Social Science and Medicine, 2005, 60, 1961-1974.	3.8	145
15	Population-based study of dying in hospital in six European countries. Palliative Medicine, 2008, 22, 702-710.	3.1	143
16	Actual and Preferred Place of Death of Home-Dwelling Patients in Four European Countries: Making Sense of Quality Indicators. PLoS ONE, 2014, 9, e93762.	2.5	119
17	Assessment of hospitalised cancer patients' needs by the Needs Evaluation Questionnaire. Annals of Oncology, 2000, 11, 31-38.	1.2	114
18	Quality-of-Life Evaluation. Journal of Pain and Symptom Management, 1998, 15, 151-158.	1.2	108

#	Article	IF	Citations
19	False-Positive Results in Mammographic Screening for Breast Cancer in Europe: A Literature Review and Survey of Service Screening Programmes. Journal of Medical Screening, 2012, 19, 57-66.	2.3	104
20	Home palliative care for terminal cancer patients: a survey on the final week of life. Palliative Medicine, 1999, 13, 233-241.	3.1	94
21	The Impact of Mammographic Screening on Breast Cancer Mortality in Europe: A Review of Trend Studies. Journal of Medical Screening, 2012, 19, 26-32.	2.3	93
22	Physicians' experiences with end-of-life decision-making: Survey in 6 European countries and Australia. BMC Medicine, 2008, 6, 4.	5.5	89
23	Prevalence, distress, management, and relief of pain during the last 3 months of cancer patients' life. Results of an Italian mortality follow-back survey. Annals of Oncology, 2009, 20, 729-735.	1.2	84
24	Terminal Cancer Patients and Timing of Referral to Palliative Care A Multicenter Prospective Cohort Study. Journal of Pain and Symptom Management, 1999, 18, 243-252.	1.2	82
25	Forgoing Treatment at the End of Life in 6 European Countries. Archives of Internal Medicine, 2005, 165, 401.	3.8	81
26	Estimate of overdiagnosis of breast cancer due to mammography after adjustment for lead time. A service screening study in Italy. Breast Cancer Research, 2006, 8, R68.	5.0	79
27	Liverpool Care Pathway for patients with cancer in hospital: a cluster randomised trial. Lancet, The, 2014, 383, 226-237.	13.7	76
28	How People Die in Hospital General Wards: A Descriptive Study. Journal of Pain and Symptom Management, 2005, 30, 33-40.	1.2	75
29	Diagnosis and prognosis disclosure among cancer patients. Results from an Italian mortality follow-back survey. Annals of Oncology, 2006, 17, 853-859.	1.2	75
30	Effectiveness of service screening: a case–control study to assess breast cancer mortality reduction. British Journal of Cancer, 2008, 99, 423-427.	6.4	75
31	Palliative Care Service Use in Four European Countries: A Cross-National Retrospective Study via Representative Networks of General Practitioners. PLoS ONE, 2013, 8, e84440.	2.5	72
32	Visiting policies in Italian intensive care units: aÂnationwide survey. Intensive Care Medicine, 2008, 34, 1256-1262.	8.2	70
33	Four Essential Drugs Needed for Quality Care of the Dying: A Delphi-Study Based International Expert Consensus Opinion. Journal of Palliative Medicine, 2013, 16, 38-43.	1.1	70
34	An estimate of overdiagnosis 15 years after the start of mammographic screening in Florence. European Journal of Cancer, 2009, 45, 3166-3171.	2.8	68
35	Advance care planning in patients with advanced cancer: A 6-country, cluster-randomised clinical trial. PLoS Medicine, 2020, 17, e1003422.	8.4	68
36	The last three months of life of Italian cancer patients. Methods, sample characteristics and response rate of the Italian Survey of the Dying of Cancer (ISDOC). Palliative Medicine, 2005, 19, 628-638.	3.1	67

#	Article	IF	CITATIONS
37	Cancer patients' needs during hospitalisation: a quantitative and qualitative study. BMC Cancer, 2003, 3, 12.	2.6	66
38	Inequity in the provision of and access to palliative care for cancer patients. Results from the Italian survey of the dying of cancer (ISDOC). BMC Public Health, 2007, 7, 66.	2.9	66
39	End-of-Life Communication: A Retrospective Survey of Representative General Practitioner Networks in Four Countries. Journal of Pain and Symptom Management, 2014, 47, 604-619.e3.	1.2	66
40	Partial liberalization of visiting policies and ICU staff: a before-and-after study. Intensive Care Medicine, 2013, 39, 2180-2187.	8.2	65
41	End-of-Life Decisions: A Cross-National Study of Treatment Preference Discussions and Surrogate Decision-Maker Appointments. PLoS ONE, 2013, 8, e57965.	2.5	62
42	Interprofessional spiritual care in oncology: a literature review. ESMO Open, 2019, 4, e000465.	4.5	61
43	End-of-life decision-making in Belgium, Denmark, Sweden and Switzerland: does place of death make a difference?. Journal of Epidemiology and Community Health, 2007, 61, 1062-1068.	3.7	59
44	Basic variables at different positivity thresholds of a quantitative immunochemical test for faecal occult blood. Journal of Medical Screening, 2002, 9, 99-103.	2.3	57
45	Using death certificate data to study place of death in 9 European countries: opportunities and weaknesses. BMC Public Health, 2007, 7, 283.	2.9	57
46	Forgoing Artificial Nutrition or Hydration in Patients Nearing Death in Six European Countries. Journal of Pain and Symptom Management, 2007, 34, 305-314.	1.2	57
47	The measure of psychological distress in cancer patients: the use of Distress Thermometer in the Oncological Rehabilitation Center of Florence. Supportive Care in Cancer, 2009, 17, 771-779.	2.2	55
48	End-of-life care in general practice: A cross-sectional, retrospective survey of â€~cancer', â€~organ failure' and â€~old-age/dementia' patients. Palliative Medicine, 2014, 28, 965-975.	3.1	53
49	Transitions between health care settings in the final three months of life in four EU countries. European Journal of Public Health, 2015, 25, 569-575.	0.3	53
50	What "best practice" could be in Palliative Care: an analysis of statements on practice and ethics expressed by the main Health Organizations. BMC Palliative Care, 2010, 9, 1.	1.8	52
51	Evaluation of Social Support, Quality of Life, and Body Image in Women with Breast Cancer. Breast Care, 2016, 11, 28-32.	1.4	52
52	Differences in place of death between lung cancer and COPD patients: a 14-country study using death certificate data. Npj Primary Care Respiratory Medicine, 2017, 27, 14.	2.6	52
53	End-of-Life Decision Making in Europe and Australia. Archives of Internal Medicine, 2006, 166, 921.	3.8	51
54	Complexity in Non-Pharmacological Caregiving Activities at the End of Life: An International Qualitative Study. PLoS Medicine, 2012, 9, e1001173.	8.4	51

#	Article	IF	Citations
55	Spiritual care in cancer patients: a need or an option?. Current Opinion in Oncology, 2018, 30, 212-218.	2.4	47
56	Parental presence and visiting policies in Italian pediatric intensive care units: A national survey*. Pediatric Critical Care Medicine, 2011, 12, e46-e50.	0.5	46
57	Overdiagnosis in breast cancer: Design and methods of estimation in observational studies. Preventive Medicine, 2011, 53, 131-133.	3.4	44
58	Predicting survival in terminal cancer patients: clinical observation or quality-of-life evaluation?. Palliative Medicine, 2005, 19, 220-227.	3.1	43
59	Advance care planning $\hat{a} \in \hat{a}$ a multi-centre cluster randomised clinical trial: the research protocol of the ACTION study. BMC Cancer, 2016, 16, 264.	2.6	43
60	Accelerated partial breast irradiation using intensity modulated radiotherapy versus whole breast irradiation: Health-related quality of life final analysis from the Florence phase 3 trial. European Journal of Cancer, 2017, 76, 17-26.	2.8	43
61	Exclusive endocrine therapy or partial breast irradiation for women aged ≥70 †years with luminal A-like early stage breast cancer (NCT04134598 †EUROPA): Proof of concept of a randomized controlled trial comparing health related quality of life by patient reported outcome measures. lournal of Geriatric Oncology, 2021, 12, 182-189.	1.0	42
62	Physician discussions with terminally ill patients: a cross-national comparison. Palliative Medicine, 2007, 21, 295-303.	3.1	41
63	The effectiveness of the Liverpool care pathway in improving end of life care for dying cancer patients in hospital. A cluster randomised trial. BMC Health Services Research, 2011, 11, 13.	2.2	40
64	Palliative home care reduces time spent in hospital wards: a population-based study in the Tuscany Region, Italy. Cancer Causes and Control, 2003, 14, 971-977.	1.8	39
65	Nationwide continuous monitoring of end-of-life care via representative networks of general practitioners in Europe. BMC Family Practice, 2013, 14, 73.	2.9	39
66	Drugs Used to Alleviate Symptoms with Life Shortening as a Possible Side Effect: End-of-Life Care in Six European Countries. Journal of Pain and Symptom Management, 2006, 31, 111-121.	1.2	38
67	Patient Dignity Inventory (PDI) Questionnaire: The Validation Study in Italian Patients with Solid and Hematological Cancers on Active Oncological Treatments. Tumori, 2012, 98, 491-500.	1.1	38
68	End-of-life preferences in advanced cancer patients willing to discuss issues surrounding their terminal condition. European Journal of Cancer Care, 2012, 21, 623-633.	1.5	38
69	Balancing harms and benefits of service mammography screening programs: a cohort study. Breast Cancer Research, 2012, 14, R9.	5.0	38
70	Deaths of children occurring at home in six European countries. Child: Care, Health and Development, 2010, 36, 375-384.	1.7	37
71	An application of the two-source capture–recapture method to estimate the completeness of the Tuscany Cancer Registry, Italy. European Journal of Cancer Prevention, 2001, 10, 417-423.	1.3	35
72	Prevalence in the use of Complementary Medicine among Cancer Patients in Tuscany, Italy. Tumori, 2008, 94, 406-410.	1.1	34

#	Article	IF	CITATIONS
73	Hope Herth Index (HHI): A Validation Study in Italian Patients with Solid and Hematological Malignancies on Active Cancer Treatment. Tumori, 2012, 98, 385-392.	1.1	34
74	Prospective observational Italian study on palliative sedation in two hospice settings: differences in casemixes and clinical care. Supportive Care in Cancer, 2012, 20, 2829-2836.	2.2	34
75	Burden on family carers and care-related financial strain at the end of life: a cross-national population-based study. European Journal of Public Health, 2014, 24, 819-826.	0.3	33
76	The Edmonton Symptom Assessment System (ESAS) as a screening tool for depression and anxiety in non-advanced patients with solid or haematological malignancies on cure or follow-up. Supportive Care in Cancer, 2014, 22, 783-793.	2.2	33
77	Suicide mortality is decreasing among cancer patients in Central Italy. European Journal of Cancer, 2004, 40, 1053-1057.	2.8	32
78	Dying of cancer in Italy: impact on family and caregiver. The Italian Survey of Dying of Cancer. Journal of Epidemiology and Community Health, 2007, 61, 547-554.	3.7	32
79	Care provided and care setting transitions in the last three months of life of cancer patients: a nationwide monitoring study in four European countries. BMC Cancer, 2014, 14, 960.	2.6	32
80	Use and Perceived Benefits of Complementary Therapies by Cancer Patients Receiving Conventional Treatment in Italy. Journal of Pain and Symptom Management, 2014, 47, 26-34.	1.2	32
81	Quality of Life Assessment and Outcome of Palliative Care. Journal of Pain and Symptom Management, 2001, 21, 179-188.	1.2	31
82	Does an organised screening programme reduce the inequalities in breast cancer survival?. Annals of Oncology, 2012, 23, 319-323.	1.2	31
83	Prognostic impact of the early detection of metachronous contralateral breast cancer. European Journal of Cancer, 2004, 40, 1496-1501.	2.8	30
84	Mortality Among Discharged Psychiatric Patients in Florence, Italy. Psychiatric Services, 2006, 57, 1474-1481.	2.0	30
85	Discussing prognosis with terminally ill cancer patients and relatives: A survey of physicians' intentions in seven countries. Patient Education and Counseling, 2009, 77, 430-436.	2.2	30
86	Assessment of psychological distress in cancer patients: a pivotal role for clinical interview. Psycho-Oncology, 2010, 19, 1294-1302.	2.3	30
87	Endâ€ofâ€ife care in Italy: personal experience of family caregivers. A content analysis of open questions from the Italian Survey of the Dying of Cancer (ISDOC). Psycho-Oncology, 2008, 17, 1073-1080.	2.3	29
88	Comparing Cancer Care, Outcomes, and Costs Across Health Systems: Charting the Course. Journal of the National Cancer Institute Monographs, 2013, 2013, 124-130.	2.1	28
89	Hospitalisations at the endÂofÂlife in four European countries: a population-based study via epidemiological surveillance networks. Journal of Epidemiology and Community Health, 2016, 70, 430-436.	3.7	28
90	Awareness of General Practitioners concerning cancer patients' preferences for place of death: Evidence from four European countries. European Journal of Cancer, 2013, 49, 1967-1974.	2.8	27

#	Article	IF	CITATIONS
91	Differences in psychophysical well-being and signs of depression in couples undergoing their first consultation for assisted reproduction technology (ART): an Italian pilot study. European Journal of Obstetrics, Gynecology and Reproductive Biology, 2016, 197, 179-185.	1.1	26
92	Is it possible to encourage hope in non-advanced cancer patients? We must try. Annals of Oncology, 2016, 27, 513-519.	1.2	26
93	Italian validation of the Purpose In Life (PIL) test and the Seeking Of Noetic Goals (SONG) test in a population of cancer patients. Supportive Care in Cancer, 2012, 20, 2775-2783.	2.2	25
94	Hope Herth Index (HHI): a validation study in Italian patients with solid and hematological malignancies on active cancer treatment. Tumori, 2012, 98, 385-92.	1.1	25
95	Informed consent and patient participation in the medical encounter: a list of questions for an informed choice about the type of anaesthesia. European Journal of Anaesthesiology, 1999, 16, 160.	1.7	24
96	Intentionally hastening death by withholding or withdrawing treatment. Wiener Klinische Wochenschrift, 2006, 118, 322-326.	1.9	23
97	Responders and non-responders in a study on medical end-of-life decisions in Denmark, the Netherlands, Sweden and Switzerland. International Journal of Public Health, 2006, 51, 24-33.	2.6	23
98	Relationships among unmet needs, depression, and anxiety in non–advanced cancer patients. Tumori, 2019, 105, 144-150.	1.1	23
99	Breast cancer mortality trends in two areas of the province of Florence, Italy, where screening programmes started in the 1970s and 1990s. British Journal of Cancer, 2004, 90, 1780-1783.	6.4	22
100	Eliciting Individual Preferences About Death: Development of the End-of-Life Preferences Interview. Journal of Pain and Symptom Management, 2008, 36, 335-350.	1.2	22
101	Use of the Needs Evaluation Questionnaire with cancer outpatients. Supportive Care in Cancer, 2016, 24, 3507-3515.	2.2	22
102	Assessing unmet needs in patients with cancer: An investigation of differential item functioning of the Needs Evaluation Questionnaire across gender, age and phase of the disease. PLoS ONE, 2017, 12, e0179765.	2.5	21
103	An increasing number of qualitative research papers in oncology and palliative care: does it mean a thorough development of the methodology of research?. Health and Quality of Life Outcomes, 2004, 2, 7.	2.4	20
104	Hospitalizations of cancer patients in the last month of life: quality indicator scores reveal large variation between four European countries in a mortality follow-back study. BMC Palliative Care, 2014, 13, 54.	1.8	20
105	End of life care preferences. Current Opinion in Supportive and Palliative Care, 2008, 2, 54-59.	1.3	19
106	Seeking and experiencing meaning: Exploring the role of meaning in promoting mental adjustment and eudaimonic well-being in cancer patients. Palliative and Supportive Care, 2015, 13, 673-681.	1.0	19
107	Palliative sedation: ethical aspects. Minerva Anestesiologica, 2017, 83, 1317-1323.	1.0	19
108	Cost profiles of colorectal cancer patients in Italy based on individual patterns of care. BMC Cancer, 2013, 13, 329.	2.6	18

#	Article	IF	Citations
109	Evidence for Adapted Physical Activity as an Effective Intervention for Upper Limb Mobility and Quality of Life in Breast Cancer Survivors. Journal of Physical Activity and Health, 2014, 11, 814-822.	2.0	18
110	Psychological Distress and Health-Related Quality of Life among Head and Neck Cancer Patients during the First Year after Treatment. Tumori, 2016, 102, 96-102.	1.1	18
111	Longitudinal assessment of the impact of adapted physical activity on upper limb disability and quality of life in breast cancer survivors from an Italian cohort. Supportive Care in Cancer, 2018, 26, 329-332.	2.2	18
112	Missing not at random in end of life care studies: multiple imputation and sensitivity analysis on data from the ACTION study. BMC Medical Research Methodology, 2021, 21, 13.	3.1	18
113	Patient dignity inventory (PDI) questionnaire: the validation study in Italian patients with solid and hematological cancers on active oncological treatments. Tumori, 2012, 98, 491-500.	1.1	18
114	Life at the end of life: beliefs about individual life after death and "good death" models - a qualitative study. Health and Quality of Life Outcomes, 2003, 1, 65.	2.4	17
115	Communicating the Balance Sheet in Breast Cancer Screening. Journal of Medical Screening, 2012, 19, 67-71.	2.3	17
116	Complementary needs behind complementary therapies in cancer patients. Psycho-Oncology, 2015, 24, 1124-1130.	2.3	17
117	The Path of Cicely Saunders: The "Peculiar Beauty―of Palliative Care. Journal of Palliative Care, 2020, 35, 3-7.	1.0	17
118	Factors Associated with Fulfilling the Preference for Dying at Home among Cancer Patients: The role of General Practitioners. Journal of Palliative Care, 2014, 30, 141-150.	1.0	16
119	How do general end-of-life treatment goals and values relate to specific treatment preferences? A population-based study. Palliative Medicine, 2014, 28, 1206-1212.	3.1	16
120	Physicians' Opinion and Practice With the Continuous Use of Sedatives in the Last Days of Life. Journal of Pain and Symptom Management, 2022, 63, 78-87.	1.2	16
121	Analysis of morphologic patterns of fine-needle aspiration of the breast to reduce false-negative results in breast cytology. Cancer, 2005, 105, 152-157.	4.1	15
122	Talking About End-of-Life Preferences With Advanced Cancer Patients: Factors Influencing Feasibility. Journal of Pain and Symptom Management, 2012, 43, 739-746.	1.2	15
123	How Are Patient Populations Characterized in Studies Investigating Depression in Advanced Cancer? Results From a Systematic Literature Review. Journal of Pain and Symptom Management, 2014, 48, 678-698.	1.2	15
124	Physician-reported practices on continuous deep sedation until death: A descriptive and comparative study. Palliative Medicine, 2014, 28, 491-500.	3.1	14
125	Unmet needs of Italian cancer patients in different stages of the disease and care process. Tumori, 2018, 104, 285-291.	1.1	14
126	Andante moderato: signs of change in visiting policies for Italian ICUs. Intensive Care Medicine, 2011, 37, 1890-1890.	8.2	13

#	Article	IF	Citations
127	Initial Treatment for Newly Diagnosed Elderly Colorectal Cancer Patients: Patterns of Care in Italy and the United States. Journal of the National Cancer Institute Monographs, 2013, 2013, 88-98.	2.1	13
128	A comparison of drugs and procedures of care in the Italian hospice and hospital settings: the final three days of life for cancer patients. BMC Health Services Research, 2014, 14, 496.	2.2	13
129	Physical and Psychological Distress Are Related to Dying Peacefully in Residents With Dementia in Long-Term Care Facilities. Journal of Pain and Symptom Management, 2015, 50, 1-8.	1.2	13
130	Differences in primary palliative care between people with organ failure and people with cancer: An international mortality follow-back study using quality indicators. Palliative Medicine, 2018, 32, 1498-1508.	3.1	13
131	Quality of primary palliative care for older people with mild and severe dementia: an international mortality follow-back study using quality indicators. Age and Ageing, 2018, 47, 824-833.	1.6	13
132	Experience of an information aid for newly diagnosed multiple sclerosis patients: a qualitative study on the SIMSâ€Trial. Health Expectations, 2014, 17, 36-48.	2.6	12
133	Coping strategies of patients with advanced lung or colorectal cancer in six European countries: Insights from the ACTION Study. Psycho-Oncology, 2020, 29, 347-355.	2.3	12
134	Are Three Items Sufficient to Measure Sense of Coherence?. European Journal of Psychological Assessment, 2018, 34, 229-237.	3.0	12
135	Melanoma incidence in central Italy will go on increasing also in the near future: A registry-based, age–period–cohort analysis. European Journal of Cancer Prevention, 2007, 16, 50-54.	1.3	11
136	Evaluation of service mammography screening impact in Italy. The contribution of hazard analysis. European Journal of Cancer, 2008, 44, 858-865.	2.8	10
137	More on Screening Mammography. New England Journal of Medicine, 2011, 364, 281-286.	27.0	10
138	Endâ€ofâ€ife care for people dying with dementia in general practice in Belgium, Italy and Spain: A crossâ€sectional, retrospective study. Geriatrics and Gerontology International, 2017, 17, 1667-1676.	1.5	10
139	Live music intervention for cancer inpatients: The Music Givers format. Palliative and Supportive Care, 2018, 16, 777-784.	1.0	10
140	Psychological well-being and quality of life in visually impaired baseball players: An Italian national survey. PLoS ONE, 2019, 14, e0218124.	2.5	10
141	OUTCOME ASSESSMENT IN CANCER MANAGEMENT. Surgical Clinics of North America, 2000, 80, 471-486.	1.5	9
142	Candidates for liver transplantation for cancer: physical, psychological, and social conditions 1. Transplantation, 2002, 73, 1627-1635.	1.0	9
143	Time trends in cancer incidence and mortality in central Italy. European Journal of Cancer Prevention, 2002, 11, 387-395.	1.3	9
144	Impact of Home Care on Place of Death, Access to Emergency Departments and Opioid Therapy in 350 Terminal Cancer Patients. Tumori, 2008, 94, 87-95.	1.1	9

#	Article	IF	Citations
145	Contesting the Equivalency of Continuous Sedation until Death and Physician-assisted Suicide/Euthanasia: A Commentary on LiPuma. Journal of Medicine and Philosophy, 2015, 40, 529-553.	0.8	9
146	Rapid and sound assessment of well-being within a multi-dimensional approach: The Well-being Numerical Rating Scales (WB-NRSs). PLoS ONE, 2021, 16, e0252709.	2.5	9
147	Does low-dose aspirin use for cardiovascular disease prevention reduce colorectal cancer deaths? A comparison of two cohorts in the Florence district, Italy. European Journal of Cancer Prevention, 2018, 27, 134-139.	1.3	8
148	Primary palliative care for older people in three European countries: a mortality follow-back quality study. BMJ Supportive and Palliative Care, 2020, 10, 462-468.	1.6	8
149	Impact of home care on place of death, access to emergency departments and opioid therapy in 350 terminal cancer patients. Tumori, 2008, 94, 87-95.	1.1	8
150	Parental presence in Italian pediatric intensive care units: a reappraisal of current visiting policies. Intensive Care Medicine, 2017, 43, 458-459.	8.2	7
151	Discussions about palliative sedation in hospice: Frequency, timing and factors associated with patient involvement. European Journal of Cancer Care, 2019, 28, e13019.	1.5	7
152	Italian version of the Edmonton Symptom Assessment System (ESAS)–Total Care (TC): development and psychometric validation in patients undergoing cancer treatment or follow-up. Supportive Care in Cancer, 2022, 30, 1923-1933.	2.2	7
153	Interlaboratory reproducibility of the immunocytochemical assessment of oestrogen and proliferative activity in fine needle aspiration of breast cancer. Cytopathology, 2002, 13, 92-100.	0.7	6
154	Functional impairment, symptom severity, and overall quality of life in patients with advanced lung or colorectal cancer in six European countries: baseline findings from the ACTION study. Supportive Care in Cancer, 2021, 29, 5797-5810.	2.2	6
155	Access to Death Certificates: What Should Research Ethics Committees Require for Approval?. Annals of Epidemiology, 2006, 16, 281-284.	1.9	5
156	Is the Spiritual Life of Cancer Patients a Resource to be taken into Account by Professional Caregivers from the Time of Diagnosis?. Tumori, 2012, 98, 158-161.	1.1	5
157	Final transitions to place of death: patients and families wishes. Journal of Public Health, 2017, 39, e302-e311.	1.8	5
158	Sociodemographic, clinical, and psychological characteristics identify groups of Italian cancer patients with high rates of unmet needs. Tumori, 2019, 105, 288-295.	1.1	5
159	Tailored Sailing Experience to Reduce Psychological Distress and Improve the Quality of Life of Breast Cancer Survivors: A Survey-Based Pilot Study. International Journal of Environmental Research and Public Health, 2020, 17, 4406.	2.6	5
160	Mortality Among Discharged Psychiatric Patients in Florence, Italy. Psychiatric Services, 2006, 57, 1474-1481.	2.0	5
161	Incidence of second cancers among women with in situ carcinoma of the breast. Breast, 2001, 10, 438-441.	2.2	4
162	Incidence and Survival Cancer Trends in Children and Adolescents in the Provinces of Florence and Prato (Central Italy), 1985–1997. Tumori, 2002, 88, 461-466.	1.1	4

#	Article	IF	CITATIONS
163	Long-Term Benefits of Adapted Physical Activity on Upper Limb Performance and Quality of Life in Breast Cancer Survivors. Journal of Functional Morphology and Kinesiology, 2017, 2, 38.	2.4	4
164	Fears, beliefs, and quality of life of patients with cancer vs the general population during the coronavirus disease 2019 (COVID-19) pandemic in Lombardy. Tumori, 2022, 108, 431-438.	1.1	4
165	Temperament and character traits associated with health-related quality of life in cancer patients. Tumori, 2012, 98, 377-84.	1.1	4
166	Intercountry and intracountry variations in opinions of palliative care specialist physicians in Germany, Italy, Japan and UK about continuous use of sedatives: an international cross-sectional survey. BMJ Open, 2022, 12, e060489.	1.9	4
167	Detection of False-Negative Pap Smears using the Papnet System. Tumori, 2000, 86, 455-457.	1.1	3
168	Decreasing incidence of all histological subtypes of oesophagus cancer in Tuscany, Italy. European Journal of Cancer Prevention, 2001, 10, 379-380.	1.3	3
169	Impact of a psycho-oncological rehabilitation intervention on psychological distress: the experience of CeRiOn (Oncological Rehabilitation Centre) Florence 2007–2010. Supportive Care in Cancer, 2013, 21, 2381-2386.	2.2	3
170	Mechanical ventilation withdrawal at end of life: terminal extubation and terminal weaning. Minerva Anestesiologica, 2017, 83, 350-352.	1.0	3
171	Assessing Emotional Functioning with Increased Power: Relative Validity of a Customized EORTC Short Form in the International ACTION Trial. Value in Health, 2019, 22, 92-98.	0.3	3
172	Construction of new personal meanings by cancer patients: a qualitative analysis in an Italian patient population. Supportive Care in Cancer, 2019, 27, 1911-1918.	2.2	3
173	Sexual Satisfaction Assessment in 194 Nonmetastatic Cancer Patients on Treatment or in follow-up. Tumori, 2014, 100, 232-236.	1.1	2
174	Feasibility of assessing quality of care at the end of life in two cluster trials using an after-death approach with multiple assessments. BMC Palliative Care, 2014, 13, 36.	1.8	2
175	Live concerts reduce cancer inpatients' anxiety. European Journal of Cancer Care, 2017, 26, e12590.	1.5	2
176	Quality of life improvement in breast cancer survivors affected by upper limb lymphedema through a novel multiperspective physical activity methodology: a monocentric pilot study. Breast Cancer, 2022, , $1. $	2.9	2
177	Is the spiritual life of cancer patients a resource to be taken into account by professional caregivers from the time of diagnosis?. Tumori, 2012, 98, 158-61.	1.1	2
178	Sexual satisfaction assessment in 194 nonmetastatic cancer patients on treatment or in follow-up. Tumori, 2014, 100, 232-6.	1.1	2
179	An Unconventional Cancer Treatment Lacking Clinical Efficacy Remains Available to Italian Cancer Patients. Tumori, 2008, 94, 830-832.	1.1	1
180	Estimates of cancer burden in Tuscany. Tumori, 2013, 99, 334-341.	1.1	1

#	Article	IF	CITATIONS
181	Estimating all cancers incidence with the MIAMOD model: a new method to include multiple tumors. Tumori, 2012, 98, 296-302.	1.1	1
182	Effects of a specific adapted upper limb exercise on lymphedema in breast cancer survivors: a pilot study. Gazzetta Medica Italiana Archivio Per Le Scienze Mediche, 2018, 177, .	0.1	1
183	Pre-operative prediction of invasive vs intraductal breast cancer type: multivariate analysis of the accuracy of clinical and imaging findings. Breast, 2002, 11, 151-155.	2.2	0
184	Ethical Concerns With Continuous Deep Sedation Until Death in France. Journal of Pain and Symptom Management, 2021, 62, e5-e6.	1.2	0
185	Act of Consoling Helps Those Who Console: A Multicenter Observational Survey Study in Hospice Services. Journal of Palliative Medicine, 2021, 24, 1226-1231.	1.1	0
186	Withholding or Withdrawing Nutritional Support at the End-of-Lifein Six European Countries. , 2011, , 231-242.		0
187	Self-reported active lifestyle behavior and upper limb disability in breast cancer survivors following a structured adapted physical activity intervention. Gazzetta Medica Italiana Archivio Per Le Scienze Mediche, 2019, 178, .	0.1	0
188	Title is missing!. , 2020, 17, e1003422.		0
189	Title is missing!. , 2020, 17, e1003422.		0
190	Title is missing!. , 2020, 17, e1003422.		0
191	Title is missing!. , 2020, 17, e1003422.		0
192	Title is missing!. , 2020, 17, e1003422.		0
193	Title is missing!. , 2020, 17, e1003422.		0
194	Title is missing!. , 2020, 17, e1003422.		0
195	Estimates of cancer burden in Tuscany. Tumori, 2013, 99, 334-41.	1.1	0
196	Do you agree with the doctor's decision to continue treatment? A scenario-based study of hospital nurses in Italy. Annali Dell'Istituto Superiore Di Sanita, 2015, 51, 352-7.	0.4	0