

# Akira Kiyonaga

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5969352/publications.pdf>

Version: 2024-02-01

13  
papers

325  
citations

933447

10  
h-index

1199594

12  
g-index

14  
all docs

14  
docs citations

14  
times ranked

518  
citing authors

#	ARTICLE	IF	CITATIONS
1	Association between echo intensity and attenuation of skeletal muscle in young and older adults: a comparison between ultrasonography and computed tomography. <i>Clinical Interventions in Aging</i> , 2018, Volume 13, 1871-1878.	2.9	39
2	Association between the Frequency of Protein-Rich Food Intakes and Kihon-Checklist Frailty Indices in Older Japanese Adults: The Kyoto-Kameoka Study. <i>Nutrients</i> , 2018, 10, 84.	4.1	17
3	Role of selected polymorphisms in determining muscle fiber composition in Japanese men and women. <i>Journal of Applied Physiology</i> , 2018, 124, 1377-1384.	2.5	22
4	Sex Difference in the Association Between Protein Intake and Frailty: Assessed Using the Kihon Checklist Indexes Among Older Adults. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 801-805.	2.5	26
5	Prevalence of Frailty Assessed by Fried and Kihon Checklist Indexes in a Prospective Cohort Study: Design and Demographics of the Kyoto-Kameoka Longitudinal Study. <i>Journal of the American Medical Directors Association</i> , 2017, 18, 733.e7-733.e15.	2.5	68
6	Lifestyle Intervention Involving Calorie Restriction with or without Aerobic Exercise Training Improves Liver Fat in Adults with Visceral Adiposity. <i>Journal of Obesity</i> , 2014, 2014, 1-8.	2.7	49
7	Aerobic Exercise Attenuates the Loss of Skeletal Muscle during Energy Restriction in Adults with Visceral Adiposity. <i>Obesity Facts</i> , 2014, 7, 26-35.	3.4	36
8	Minute-by-minute stepping rate of daily physical activity in normal and overweight/obese adults. <i>Obesity Research and Clinical Practice</i> , 2011, 5, e151-e156.	1.8	16
9	Relationships between body fat accumulation, aerobic capacity and insulin resistance in Japanese participants. <i>Obesity Research and Clinical Practice</i> , 2011, 5, e143-e150.	1.8	2
10	A 12-week aerobic exercise program without energy restriction improves intrahepatic fat, liver function and atherosclerosis-related factors. <i>Obesity Research and Clinical Practice</i> , 2011, 5, e249-e257.	1.8	14
11	Changes in Serum Concentrations of Taurine and Other Amino Acids in Clinical Antihypertensive Exercise Therapy. <i>Clinical and Experimental Hypertension</i> , 1989, 11, 149-165.	0.3	33
12	Effects of Mild Exercise Therapy on Serum High Density Lipoprotein Subfraction in Patients with Essential Hypertension. <i>The Journal of Japan Atherosclerosis Society</i> , 1986, 14, 1107-1109.	0.0	0
13	Effects of Mild Exercise Therapy on Serum Lipids and Apoproteins. <i>The Journal of Japan Atherosclerosis Society</i> , 1985, 13, 189-193.	0.0	3