

Max Birchwood

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5973085/max-birchwood-publications-by-citations.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

268
papers

15,440
citations

61
h-index

118
g-index

337
ext. papers

17,316
ext. citations

4.7
avg, IF

6.47
L-index

#	Paper	IF	Citations
268	The psychosis high-risk state: a comprehensive state-of-the-art review. <i>JAMA Psychiatry</i> , 2013 , 70, 107-204	4.5	965
267	The Social Functioning Scale. The development and validation of a new scale of social adjustment for use in family intervention programmes with schizophrenic patients. <i>British Journal of Psychiatry</i> , 1990 , 157, 853-9	5.4	788
266	The omnipotence of voices. A cognitive approach to auditory hallucinations. <i>British Journal of Psychiatry</i> , 1994 , 164, 190-201	5.4	615
265	Early intervention in psychosis: The critical period hypothesis. <i>British Journal of Psychiatry</i> , 1998 , 172, 53-59	5.4	522
264	Prediction of psychosis in adolescents and young adults at high risk: results from the prospective European prediction of psychosis study. <i>Archives of General Psychiatry</i> , 2010 , 67, 241-51		491
263	A self-report Insight Scale for psychosis: reliability, validity and sensitivity to change. <i>Acta Psychiatrica Scandinavica</i> , 1994 , 89, 62-7	6.5	483
262	The power and omnipotence of voices: subordination and entrapment by voices and significant others. <i>Psychological Medicine</i> , 2000 , 30, 337-44	6.9	377
261	Cognitive therapy and recovery from acute psychosis: a controlled trial. I. Impact on psychotic symptoms. <i>British Journal of Psychiatry</i> , 1996 , 169, 593-601	5.4	329
260	Designing youth mental health services for the 21st century: examples from Australia, Ireland and the UK. <i>British Journal of Psychiatry</i> , 2013 , 54, s30-5	5.4	295
259	Depression, demoralization and control over psychotic illness: a comparison of depressed and non-depressed patients with a chronic psychosis. <i>Psychological Medicine</i> , 1993 , 23, 387-95	6.9	285
258	Predicting relapse in schizophrenia: the development and implementation of an early signs monitoring system using patients and families as observers, a preliminary investigation. <i>Psychological Medicine</i> , 1989 , 19, 649-56	6.9	265
257	'Theory of mind' skills during an acute episode of psychosis and following recovery. <i>Psychological Medicine</i> , 1998 , 28, 1101-12	6.9	262
256	The revised Beliefs About Voices Questionnaire (BAVQ-R). <i>British Journal of Psychiatry</i> , 2000 , 177, 229-32	5.4	218
255	Pathways to emotional dysfunction in first-episode psychosis. <i>British Journal of Psychiatry</i> , 2003 , 182, 373-375	5.4	216
254	Cognitive therapy for command hallucinations: randomised controlled trial. <i>British Journal of Psychiatry</i> , 2004 , 184, 312-20	5.4	211
253	Early detection and intervention evaluation for people at risk of psychosis: multisite randomised controlled trial. <i>BMJ, The</i> , 2012 , 344, e2233	5.9	209
252	Cognitive approach to depression and suicidal thinking in psychosis. 1. Ontogeny of post-psychotic depression. <i>British Journal of Psychiatry</i> , 2000 , 177, 516-21	5.4	202

251	The omnipotence of voices: testing the validity of a cognitive model. <i>Psychological Medicine</i> , 1997 , 27, 1345-53	6.9	201
250	Interpersonal and role-related schema influence the relationship with the dominant 'voice' in schizophrenia: a comparison of three models. <i>Psychological Medicine</i> , 2004 , 34, 1571-80	6.9	183
249	Psychotic-like experiences and correlation with distress and depressive symptoms in a community sample of adolescents and young adults. <i>Schizophrenia Research</i> , 2010 , 119, 258-65	3.6	174
248	Cognitive approach to depression and suicidal thinking in psychosis: 2. Testing the validity of a social ranking model. <i>British Journal of Psychiatry</i> , 2000 , 177, 522-528	5.4	169
247	Early intervention in psychosis. The critical period hypothesis. <i>The British Journal of Psychiatry Supplement</i> , 1998 , 172, 53-9		166
246	Social anxiety and the shame of psychosis: a study in first episode psychosis. <i>Behaviour Research and Therapy</i> , 2007 , 45, 1025-37	5.2	158
245	A new scale (SES) to measure engagement with community mental health services. <i>Journal of Mental Health</i> , 2002 , 11, 191-8	2.7	155
244	Depression and Schizophrenia: Cause, Consequence, or Trans-diagnostic Issue?. <i>Schizophrenia Bulletin</i> , 2017 , 43, 240-244	1.3	154
243	Cognitive therapy and recovery from acute psychosis: a controlled trial. II. Impact on recovery time. <i>British Journal of Psychiatry</i> , 1996 , 169, 602-7	5.4	150
242	Predicting engagement with services for psychosis: insight, symptoms and recovery style. <i>British Journal of Psychiatry</i> , 2003 , 182, 123-8	5.4	146
241	Specific and non-specific effects of educational intervention with families living with a schizophrenic relative. <i>British Journal of Psychiatry</i> , 1987 , 150, 645-52	5.4	144
240	The omnipotence of voices. II: The Beliefs About Voices Questionnaire (BAVQ). <i>British Journal of Psychiatry</i> , 1995 , 166, 773-6	5.4	129
239	The evolution of depression and suicidality in first episode psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2010 , 122, 211-8	6.5	124
238	Psychological pathways to depression in schizophrenia: studies in acute psychosis, post psychotic depression and auditory hallucinations. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2005 , 255, 202-12	5.1	121
237	An exploration of evolved mental mechanisms for dominant and subordinate behaviour in relation to auditory hallucinations in schizophrenia and critical thoughts in depression. <i>Psychological Medicine</i> , 2001 , 31, 1117-27	6.9	121
236	Adapting to the challenge of psychosis: personal resilience and the use of sealing-over (avoidant) coping strategies. <i>British Journal of Psychiatry</i> , 2004 , 185, 410-5	5.4	120
235	From environment to therapy in psychosis: a real-world momentary assessment approach. <i>Schizophrenia Bulletin</i> , 2011 , 37, 244-7	1.3	114
234	Cognitive therapy and recovery from acute psychosis: a controlled trial. 3. Five-year follow-up. <i>British Journal of Psychiatry</i> , 2000 , 177, 8-14	5.4	113

233	Clinical outcomes of Joint Crisis Plans to reduce compulsory treatment for people with psychosis: a randomised controlled trial. <i>Lancet, The</i> , 2013 , 381, 1634-41	4.0	111
232	Early intervention in schizophrenia. <i>Australian and New Zealand Journal of Psychiatry</i> , 1993 , 27, 374-8	2.6	111
231	Reducing duration of untreated psychosis: care pathways to early intervention in psychosis services. <i>British Journal of Psychiatry</i> , 2013 , 203, 58-64	5.4	108
230	Schizophrenia: early warning signs. <i>Advances in Psychiatric Treatment</i> , 2000 , 6, 93-101		102
229	Disability in people clinically at high risk of psychosis. <i>British Journal of Psychiatry</i> , 2010 , 197, 278-84	5.4	101
228	Cognitive behaviour therapy to prevent harmful compliance with command hallucinations (COMMAND): a randomised controlled trial. <i>Lancet Psychiatry, the</i> , 2014 , 1, 23-33	23.3	100
227	Families coping with schizophrenia: coping styles, their origins and correlates. <i>Psychological Medicine</i> , 1990 , 20, 857-65	6.9	100
226	The European Prediction of Psychosis Study (EPOS): integrating early recognition and intervention in Europe. <i>World Psychiatry</i> , 2005 , 4, 161-7	14.4	97
225	The influence of ethnicity and family structure on relapse in first-episode schizophrenia. A comparison of Asian, Afro-Caribbean, and white patients. <i>British Journal of Psychiatry</i> , 1992 , 161, 783-90	5.4	92
224	Social anxiety disorder in first-episode psychosis: incidence, phenomenology and relationship with paranoia. <i>British Journal of Psychiatry</i> , 2009 , 195, 234-41	5.4	91
223	Relatives and patients as partners in the management of schizophrenia. The development of a service model. <i>British Journal of Psychiatry</i> , 1990 , 156, 654-60	5.4	91
222	Specific and non-specific effects of educational intervention for families living with schizophrenia. A comparison of three methods. <i>British Journal of Psychiatry</i> , 1992 , 160, 806-14	5.4	87
221	Axis I diagnoses and transition to psychosis in clinical high-risk patients EPOS project: prospective follow-up of 245 clinical high-risk outpatients in four countries. <i>Schizophrenia Research</i> , 2012 , 138, 192-7	3.6	85
220	Loss, humiliation and entrapment as appraisals of schizophrenic illness: a prospective study of depressed and non-depressed patients. <i>British Journal of Clinical Psychology</i> , 1998 , 37, 259-68	3.6	77
219	A systematic review and meta-regression analysis of aggression during the first episode of psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2013 , 128, 413-21	6.5	75
218	Whither the attenuated psychosis syndrome?. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1130-4	1.3	74
217	Is early intervention for psychosis a waste of valuable resources?. <i>British Journal of Psychiatry</i> , 2003 , 182, 196-8	5.4	74
216	Expressed emotion as an adaptation to loss: prospective study in first-episode psychosis. <i>British Journal of Psychiatry</i> , 2005 , 48, s59-64	5.4	73

215	Duration of untreated psychosis and pathways to care in first-episode psychosis. Investigation of help-seeking behaviour in primary care. <i>British Journal of Psychiatry</i> , 2002 , 43, s73-7	5.4	73
214	Early attachment experience and recovery from psychosis. <i>British Journal of Clinical Psychology</i> , 1998 , 37, 269-84	3.6	71
213	Pathways to violent behavior during first-episode psychosis: a report from the UK National EDEN Study. <i>JAMA Psychiatry</i> , 2013 , 70, 1287-93	14.5	69
212	Early intervention in psychosis: keeping faith with evidence-based health care. <i>Psychological Medicine</i> , 2010 , 40, 399-404	6.9	68
211	Views of young people in early intervention services for first-episode psychosis in England. <i>Psychiatric Services</i> , 2011 , 62, 882-7	3.3	67
210	Efficacy and safety of pharmacological and psychological interventions for the treatment of psychosis and schizophrenia in children, adolescents and young adults: a systematic review and meta-analysis. <i>PLoS ONE</i> , 2015 , 10, e0117166	3.7	66
209	Cannabis use and age at onset of symptoms in subjects at clinical high risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2012 , 125, 45-53	6.5	61
208	Improving psychological adjustment following a first episode of psychosis: a randomised controlled trial of cognitive therapy to reduce post psychotic trauma symptoms. <i>Behaviour Research and Therapy</i> , 2009 , 47, 454-62	5.2	59
207	REDIRECT: cluster randomised controlled trial of GP training in first-episode psychosis. <i>British Journal of General Practice</i> , 2009 , 59, e183-90	1.6	59
206	The UK national evaluation of the development and impact of Early Intervention Services (the National EDEN studies): study rationale, design and baseline characteristics. <i>Microbial Biotechnology</i> , 2014 , 8, 59-67	3.3	57
205	Drug and alcohol problems amongst individuals with severe mental health problems in an inner city area of the UK. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2001 , 36, 448-55	4.5	57
204	Impact of contact on adolescents' mental health literacy and stigma: the SchoolSpace cluster randomised controlled trial. <i>BMJ Open</i> , 2016 , 6, e009435	3	56
203	Acting on command hallucinations: a cognitive approach. <i>British Journal of Clinical Psychology</i> , 1997 , 36, 139-48	3.6	56
202	Early intervention in psychotic relapse. <i>Clinical Psychology Review</i> , 2001 , 21, 1211-26	10.8	56
201	Early intervention in schizophrenia: theoretical background and clinical strategies. <i>British Journal of Clinical Psychology</i> , 1992 , 31, 257-78	3.6	56
200	Cannabis Use Is Associated With Increased Psychotic Symptoms and Poorer Psychosocial Functioning in First-Episode Psychosis: A Report From the UK National EDEN Study. <i>Schizophrenia Bulletin</i> , 2016 , 42, 619-25	1.3	53
199	Investigating trajectories of social recovery in individuals with first-episode psychosis: a latent class growth analysis. <i>British Journal of Psychiatry</i> , 2015 , 207, 536-43	5.4	53
198	Early detection and intervention evaluation for people at high-risk of psychosis-2 (EDIE-2): trial rationale, design and baseline characteristics. <i>Microbial Biotechnology</i> , 2011 , 5, 24-32	3.3	53

197	The trauma of first episode psychosis: the role of cognitive mediation. <i>Australian and New Zealand Journal of Psychiatry</i> , 2004 , 38, 327-33	2.6	53
196	The subjective experience and phenomenology of depression following first episode psychosis: a qualitative study using photo-elicitation. <i>Journal of Affective Disorders</i> , 2013 , 149, 166-74	6.6	51
195	Comparing time use in individuals at different stages of psychosis and a non-clinical comparison group. <i>Schizophrenia Research</i> , 2015 , 161, 188-93	3.6	51
194	Barriers to shared decision making in mental health care: qualitative study of the Joint Crisis Plan for psychosis. <i>Health Expectations</i> , 2016 , 19, 448-58	3.7	50
193	Management of first-episode psychosis. <i>Advances in Psychiatric Treatment</i> , 2001 , 7, 133-140		49
192	Social recovery therapy in combination with early intervention services for enhancement of social recovery in patients with first-episode psychosis (SUPEREDEN3): a single-blind, randomised controlled trial. <i>Lancet Psychiatry</i> , 2018 , 5, 41-50	23.3	48
191	Evidence for a cognitive model of auditory hallucinations. <i>Journal of Nervous and Mental Disease</i> , 2003 , 191, 542-5	1.8	47
190	The needs of high and low expressed emotion families: a normative approach. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 1993 , 28, 11-6	4.5	43
189	Acting on command hallucinations and dangerous behavior: a critique of the major findings in the last decade. <i>Clinical Psychology Review</i> , 2004 , 24, 513-28	10.8	41
188	Development and implementation of early intervention services for young people with psychosis: case study. <i>British Journal of Psychiatry</i> , 2009 , 194, 446-50	5.4	40
187	Delays in mental health services and duration of untreated psychosis. <i>Psychiatric Bulletin</i> , 2007 , 31, 408-410		39
186	The Critical Period for Early Intervention. <i>Psychiatric Rehabilitation Skills</i> , 2000 , 4, 182-198		39
185	Ethnicity and pathways to care during first episode psychosis: the role of cultural illness attributions. <i>BMC Psychiatry</i> , 2015 , 15, 287	4.2	38
184	A multi-element psychosocial intervention for early psychosis (GET UP PIANO TRIAL) conducted in a catchment area of 10 million inhabitants: study protocol for a pragmatic cluster randomized controlled trial. <i>Trials</i> , 2012 , 13, 73	2.8	38
183	Acting on voices: Omnipotence, sources of threat, and safety-seeking behaviours. <i>British Journal of Clinical Psychology</i> , 2008 , 47, 201-13	3.6	38
182	Early intervention in psychosis: opportunities for secondary prevention. <i>British Journal of Clinical Psychology</i> , 1996 , 35, 487-502	3.6	38
181	Early Intervention in Psychotic Relapse: Cognitive Approaches to Detection and Management. <i>Behaviour Change</i> , 1995 , 12, 2-19	1.1	38
180	Social anxiety disorder and shame cognitions in psychosis. <i>Psychological Medicine</i> , 2013 , 43, 133-42	6.9	37

179	Insight, duration of untreated psychosis and attachment in first-episode psychosis: prospective study of psychiatric recovery over 12-month follow-up. <i>British Journal of Psychiatry</i> , 2014 , 205, 60-7	5.4	36
178	Prediction of psychosis in clinical high-risk patients by the Schizotypal Personality Questionnaire. Results of the EPOS project. <i>European Psychiatry</i> , 2013 , 28, 469-75	6	36
177	Combined severe mental health and substance use problems: What are the training and support needs of staff working with this client group?. <i>Journal of Mental Health</i> , 2001 , 10, 131-140	2.7	36
176	Ethnicity and detention: are Black and minority ethnic (BME) groups disproportionately detained under the Mental Health Act 2007?. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016 , 51, 703-11	4.5	36
175	Layers of listening: qualitative analysis of the impact of early intervention services for first-episode psychosis on carers' experiences. <i>British Journal of Psychiatry</i> , 2015 , 207, 135-42	5.4	35
174	What service users with psychotic disorders want in a mental health crisis or relapse: thematic analysis of joint crisis plans. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014 , 49, 1609-17	4.5	34
173	Early intervention in psychosis. <i>International Clinical Psychopharmacology</i> , 1998 , 13, S31-S40	2.2	34
172	Internalized stigma, emotional dysfunction and unusual experiences in young people at risk of psychosis. <i>Microbial Biotechnology</i> , 2015 , 9, 133-40	3.3	33
171	Service users' views of moving on from early intervention services for psychosis: a longitudinal qualitative study in primary care. <i>British Journal of General Practice</i> , 2012 , 62, e183-90	1.6	32
170	Impact of cognitive therapy on internalised stigma in people with at-risk mental states. <i>British Journal of Psychiatry</i> , 2013 , 203, 140-5	5.4	32
169	Control of auditory hallucinations through occlusion of monaural auditory input. <i>British Journal of Psychiatry</i> , 1986 , 149, 104-7	5.4	32
168	Predicting compliance with command hallucinations: anger, impulsivity and appraisals of voices' power and intent. <i>Schizophrenia Research</i> , 2013 , 147, 163-168	3.6	31
167	A prospective study of dysfunctional thinking and the regulation of negative intrusive memories in bipolar 1 disorder: implications for affect regulation theory. <i>Psychological Medicine</i> , 2007 , 37, 689-98	6.9	31
166	Youth services: meeting the mental health needs of adolescents. <i>Irish Journal of Psychological Medicine</i> , 2015 , 32, 13-19	3	30
165	Transitioning care-leavers with mental health needs: 'they set you up to fail!'. <i>Child and Adolescent Mental Health</i> , 2017 , 22, 138-147	2.2	28
164	Seeking help for first-episode psychosis: a family narrative. <i>Microbial Biotechnology</i> , 2016 , 10, 334-45	3.3	28
163	The course of negative symptom in first episode psychosis and the relationship with social recovery. <i>Schizophrenia Research</i> , 2016 , 174, 165-171	3.6	28
162	Psychosocial outcome in patients at clinical high risk of psychosis: a prospective follow-up. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013 , 48, 303-11	4.5	27

161	Design of the BiRmington Early Detection In untREated psyChosis Trial (REDIRECT): cluster randomised controlled trial of general practitioner education in detection of first episode psychosis [ISRCTN87898421]. <i>BMC Health Services Research</i> , 2005 , 5, 19	2.9	27
160	Impact of early intervention services on duration of untreated psychosis: data from the National EDEN prospective cohort study. <i>Schizophrenia Research</i> , 2014 , 159, 1-6	3.6	26
159	Attachment: developmental pathways to affective dysregulation in young people at ultra-high risk of developing psychosis. <i>British Journal of Clinical Psychology</i> , 2013 , 52, 424-37	3.6	26
158	CRIMSON [Crisis plan Impact: Subjective and Objective coercion and eNgagement] protocol: a randomised controlled trial of joint crisis plans to reduce compulsory treatment of people with psychosis. <i>Trials</i> , 2010 , 11, 102	2.8	26
157	Preventing the entrenchment of high expressed emotion in first episode psychosis: early developmental attachment pathways. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34 Suppl, S191-7	2.6	26
156	Effect of delaying treatment of first-episode psychosis on symptoms and social outcomes: a longitudinal analysis and modelling study. <i>Lancet Psychiatry</i> , 2020 , 7, 602-610	23.3	25
155	A prospective study of PTSD following recovery from first-episode psychosis: the threat from persecutors, voices, and patienthood. <i>British Journal of Clinical Psychology</i> , 2012 , 51, 418-33	3.6	25
154	Cognitive approach to depression and suicidal thinking in psychosis. 2. Testing the validity of a social ranking model. <i>British Journal of Psychiatry</i> , 2000 , 177, 522-8	5.4	25
153	Short-term functional outcome and premorbid adjustment in clinical high-risk patients. Results of the EPOS project. <i>European Psychiatry</i> , 2014 , 29, 371-80	6	24
152	Personal beliefs about illness questionnaire-revised (PBIQ-R): reliability and validation in a first episode sample. <i>British Journal of Clinical Psychology</i> , 2012 , 51, 448-58	3.6	24
151	A preliminary evaluation of integrated treatment for co-existing substance use and severe mental health problems: Impact on teams and service users. <i>Journal of Mental Health</i> , 2006 , 15, 577-591	2.7	24
150	The person in recovery from acute and severe psychosis: the role of dependency, self-criticism, and efficacy. <i>American Journal of Orthopsychiatry</i> , 2004 , 74, 480-488	2.8	24
149	The national policy reforms for mental health services and the story of early intervention services in the United Kingdom. <i>Journal of Psychiatry and Neuroscience</i> , 2005 , 30, 362-5	4.5	24
148	Social anxiety in first-episode psychosis: the role of childhood trauma and adult attachment. <i>Journal of Affective Disorders</i> , 2014 , 163, 102-9	6.6	23
147	The Strauss and Carpenter Prognostic Scale in subjects clinically at high risk of psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2013 , 127, 53-61	6.5	23
146	Improving Therapeutic Relationships: Joint Crisis Planning for Individuals With Psychotic Disorders. <i>Qualitative Health Research</i> , 2015 , 25, 1637-47	3.9	23
145	Informing people with schizophrenia about their illness: The effect of residual symptoms. <i>Journal of Mental Health</i> , 1992 , 1, 61-70	2.7	23
144	Antipsychotic medication versus psychological intervention versus a combination of both in adolescents with first-episode psychosis (MAPS): a multicentre, three-arm, randomised controlled pilot and feasibility study. <i>Lancet Psychiatry</i> , 2020 , 7, 788-800	23.3	23

143	Can the therapeutic relationship predict 18 month outcomes for individuals with psychosis?. <i>Psychiatry Research</i> , 2014 , 220, 585-91	9.9	22
142	Psychotic experience subtypes, poor mental health status and help-seeking behaviour in a community sample of young adults. <i>Microbial Biotechnology</i> , 2012 , 6, 300-8	3.3	22
141	The contribution of shame to post-psychotic trauma. <i>British Journal of Clinical Psychology</i> , 2013 , 52, 162-8	3.2	22
140	Pathways to care in subjects at high risk for psychotic disorders - a European perspective. <i>Schizophrenia Research</i> , 2014 , 152, 400-7	3.6	21
139	Randomised controlled trial of joint crisis plans to reduce compulsory treatment for people with psychosis: economic outcomes. <i>PLoS ONE</i> , 2013 , 8, e74210	3.7	21
138	Autobiographical memory and post-psychotic depression in first episode psychosis. <i>British Journal of Clinical Psychology</i> , 2004 , 43, 97-104	3.6	21
137	Cognitive behavioural therapy group work with voice hearers. Part 1. <i>British Journal of Nursing</i> , 2007 , 16, 248-52	0.7	20
136	Early Intervention in Psychosis. <i>Disease Management and Health Outcomes</i> , 2005 , 13, 53-63		20
135	The development and implementation of an educational intervention on first episode psychosis for primary care. <i>Medical Education</i> , 2005 , 39, 1006-14	3.7	20
134	Negative cognition, affect, metacognition and dimensions of paranoia in people at ultra-high risk of psychosis: a multi-level modelling analysis. <i>Psychological Medicine</i> , 2015 , 45, 2675-84	6.9	19
133	Early intervention and sustaining the management of vulnerability. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34 Suppl, S181-4	2.6	19
132	Views of early psychosis clinicians on discontinuation of antipsychotic medication following symptom remission in first episode psychosis. <i>Microbial Biotechnology</i> , 2016 , 10, 355-61	3.3	19
131	Joint crisis planning in mental health care: the challenge of implementation in randomized trials and in routine care. <i>World Psychiatry</i> , 2015 , 14, 281-3	14.4	18
130	The concept of schizophrenia and phase-specific treatment: cognitive-behavioral treatment in pre-psychosis and in nonresponders. <i>Journal of the American Academy of Psychoanalysis and Dynamic Psychiatry</i> , 2003 , 31, 209-28		18
129	Depression predicts persistence of paranoia in clinical high-risk patients to psychosis: results of the EPOS project. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016 , 51, 247-57	4.5	17
128	Prevalence of psychotic-like experiences in young adults with social anxiety disorder and correlation with affective dysregulation. <i>Journal of Nervous and Mental Disease</i> , 2013 , 201, 1053-9	1.8	17
127	Through the looking glass: self-reassuring meta-cognitive capacity and its relationship with the thematic content of voices. <i>Frontiers in Human Neuroscience</i> , 2013 , 7, 213	3.3	17
126	Cognitive behavioural therapy group work with voice hearers. Part 2. <i>British Journal of Nursing</i> , 2007 , 16, 292-5	0.7	17

125	Development and validation of multivariable prediction models of remission, recovery, and quality of life outcomes in people with first episode psychosis: a machine learning approach. <i>The Lancet Digital Health</i> , 2019 , 1, e261-e270	14.4	16
124	Core outcome sets for use in effectiveness trials involving people with bipolar and schizophrenia in a community-based setting (PARTNERS2): study protocol for the development of two core outcome sets. <i>Trials</i> , 2015 , 16, 47	2.8	16
123	Preservation and compensation: the functional neuroanatomy of insight and working memory in schizophrenia. <i>Schizophrenia Research</i> , 2014 , 152, 201-9	3.6	16
122	A randomised controlled feasibility trial for an educational school-based mental health intervention: study protocol. <i>BMC Psychiatry</i> , 2012 , 12, 23	4.2	16
121	Command Hallucinations: Cognitive Theory, Therapy, and Research. <i>Journal of Cognitive Psychotherapy</i> , 2003 , 17, 67-84	0.9	16
120	Stigma and access to care in first-episode psychosis. <i>Microbial Biotechnology</i> , 2019 , 13, 1208-1213	3.3	16
119	Components of therapy as mechanisms of change in cognitive therapy for people at risk of psychosis: analysis of the EDIE-2 trial. <i>British Journal of Psychiatry</i> , 2015 , 207, 123-9	5.4	15
118	Barriers and facilitators to partnership working between Early Intervention Services and the voluntary and community sector. <i>Health and Social Care in the Community</i> , 2008 , 16, 493-500	2.6	15
117	Pilot randomised trial of a brief intervention for comorbid substance misuse in psychiatric in-patient settings. <i>Acta Psychiatrica Scandinavica</i> , 2016 , 133, 298-309	6.5	15
116	Co-designing a virtual world with young people to deliver social cognition therapy in early psychosis. <i>Microbial Biotechnology</i> , 2020 , 14, 37-43	3.3	15
115	The COMMAND trial of cognitive therapy to prevent harmful compliance with command hallucinations: predictors of outcome and mediators of change. <i>Psychological Medicine</i> , 2018 , 48, 1966-1974	6.9	15
114	A multi-centre, randomised controlled trial of cognitive therapy to prevent harmful compliance with command hallucinations. <i>BMC Psychiatry</i> , 2011 , 11, 155	4.2	14
113	Psychosocial family intervention in schizophrenia: a review of empirical studies. <i>British Journal of Psychiatry</i> , 1992 , 160, 272-5	5.4	14
112	Carers' and service users' experiences of early intervention in psychosis services: implications for care partnerships. <i>Microbial Biotechnology</i> , 2018 , 12, 410-416	3.3	13
111	Don't turn your back on the symptoms of psychosis: the results of a proof-of-principle, quasi-experimental intervention to reduce duration of untreated psychosis. <i>BMC Psychiatry</i> , 2016 , 16, 127	4.2	13
110	Preventing the Entrenchment of High Expressed Emotion in First Episode Psychosis: Early Developmental Attachment Pathways. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34, A191-A197	2.6	13
109	Early Signs Monitoring to Prevent Relapse in Psychosis and Promote Well-Being, Engagement, and Recovery: Protocol for a Feasibility Cluster Randomized Controlled Trial Harnessing Mobile Phone Technology Blended With Peer Support. <i>JMIR Research Protocols</i> , 2020 , 9, e15058	2	13
108	Duration of untreated psychosis and clinical outcomes of first episode psychosis: An observational and an instrumental variables analysis. <i>Microbial Biotechnology</i> , 2019 , 13, 841-847	3.3	13

107	Power and perceived expressed emotion of voices: their impact on depression and suicidal thinking in those who hear voices. <i>Clinical Psychology and Psychotherapy</i> , 2013 , 20, 199-205	2.9	11
106	Metacognitive beliefs as psychological predictors of social functioning: An investigation with young people at risk of psychosis. <i>Psychiatry Research</i> , 2018 , 262, 520-526	9.9	10
105	CBT for psychosis: not a 'quasi-neuroleptic'. <i>British Journal of Psychiatry</i> , 2014 , 204, 488-9	5.4	10
104	Perceived negative attitude of others predicts transition to psychosis in patients at risk of psychosis. <i>European Psychiatry</i> , 2012 , 27, 264-6	6	10
103	Early Intervention and Sustaining the Management of Vulnerability. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34, A181-A184	2.6	10
102	Specialist community teams backed by years of quality research. <i>The Psychiatrist</i> , 2013 , 37, 38-38		10
101	Predictors of engagement in first-episode psychosis. <i>Schizophrenia Research</i> , 2016 , 175, 204-208	3.6	10
100	Vulnerability to psychosocial disability in psychosis. <i>Epidemiology and Psychiatric Sciences</i> , 2019 , 28, 140-145	3.45	10
99	Expressed emotion as a predictor of the first psychotic episode - Results of the European prediction of psychosis study. <i>Schizophrenia Research</i> , 2018 , 199, 346-352	3.6	9
98	Cost-effectiveness of early intervention services for psychosis and fidelity to national policy implementation guidance. <i>Microbial Biotechnology</i> , 2018 , 12, 747-756	3.3	9
97	Usefulness of EQ-5D for evaluation of health-related quality of life in young adults with first-episode psychosis. <i>Quality of Life Research</i> , 2013 , 22, 1055-63	3.7	9
96	Individualisation in crisis planning for people with psychotic disorders. <i>Epidemiology and Psychiatric Sciences</i> , 2014 , 23, 353-9	5.1	9
95	Perceived negative attitude of others as an early sign of psychosis. <i>European Psychiatry</i> , 2009 , 24, 233-8	6	9
94	Service innovations: Coexisting severe mental health and substance use problems: developing integrated services in the UK. <i>Psychiatric Bulletin</i> , 2003 , 27, 183-186		9
93	Psychological and social treatments: course and outcome. <i>Current Opinion in Psychiatry</i> , 1999 , 12, 61-66	4.9	9
92	Short-term outcome of substance-induced psychotic disorder in a large UK first episode psychosis cohort. <i>Acta Psychiatrica Scandinavica</i> , 2016 , 134, 321-8	6.5	9
91	Refining a model of collaborative care for people with a diagnosis of bipolar, schizophrenia or other psychoses in England: a qualitative formative evaluation. <i>BMC Psychiatry</i> , 2019 , 19, 7	4.2	9
90	A Feasibility and Acceptability Trial of Social Cognitive Therapy in Early Psychosis Delivered Through a Virtual World: The VEEP Study. <i>Frontiers in Psychiatry</i> , 2020 , 11, 219	5	9

89	A high-fidelity virtual environment for the study of paranoia. <i>Schizophrenia Research and Treatment</i> , 2013 , 2013, 538185	0.6	8
88	Cognitive Behavioural Relating Therapy (CBRT) for voice hearers: a case study. <i>Behavioural and Cognitive Psychotherapy</i> , 2013 , 41, 626-31	2.1	8
87	Cognitive Therapy for Command Hallucinations: Not a Quasi-Neuroleptic. <i>Journal of Contemporary Psychotherapy</i> , 2006 , 36, 1-7	2.3	8
86	Psychotherapies for Schizophrenia: A Review 2001 , 147-241		8
85	Distress, omnipotence, and responsibility beliefs in command hallucinations. <i>British Journal of Clinical Psychology</i> , 2017 , 56, 303-309	3.6	7
84	The CIRCuITS study (Implementation of cognitive remediation in early intervention services): protocol for a randomised controlled trial. <i>Trials</i> , 2018 , 19, 183	2.8	7
83	Adolescent construction of mental illness: implication for engagement and treatment. <i>Microbial Biotechnology</i> , 2018 , 12, 626-636	3.3	7
82	Childhood adversity predicts persistence of suicidal thoughts differently in females and males at clinical high-risk patients of psychosis. Results of the EPOS project. <i>Microbial Biotechnology</i> , 2019 , 13, 935-942	3.3	7
81	Systematic Review of Cognitive-Behavioural Therapy for Social Anxiety Disorder in Psychosis. <i>Brain Sciences</i> , 2017 , 7,	3.4	7
80	Psychopathology and affect dysregulation across the continuum of psychosis: a multiple comparison group study. <i>Microbial Biotechnology</i> , 2014 , 8, 221-8	3.3	7
79	Muslim faith healers' views on substance misuse and psychosis. <i>Mental Health, Religion and Culture</i> , 2012 , 15, 653-673	1.1	7
78	Preventing Relapse of Psychotic Illness. <i>Disease Management and Health Outcomes</i> , 2002 , 10, 395-407		7
77	Prevalence of treatment resistance and clozapine use in early intervention services. <i>BJPsych Open</i> , 2020 , 6, e107	5	7
76	Cognitive Predictors of Social and Occupational Functioning in Early Psychosis: A Systematic Review and Meta-analysis of Cross-Sectional and Longitudinal Data. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1243-1253 ^{1.3}		7
75	Lived experiences of negative symptoms in first-episode psychosis: A qualitative secondary analysis. <i>Microbial Biotechnology</i> , 2019 , 13, 773-779	3.3	7
74	Informed, advance refusals of treatment by people with severe mental illness in a randomised controlled trial of joint crisis plans: demand, content and correlates. <i>BMC Psychiatry</i> , 2017 , 17, 376	4.2	6
73	Study protocol for a randomised controlled trial of CBT vs antipsychotics vs both in 14-18-year-olds: Managing Adolescent first episode Psychosis: a feasibility study (MAPS). <i>Trials</i> , 2019 , 20, 395	2.8	6
72	A pilot study to assess the feasibility and impact of a brief motivational intervention on problem drug and alcohol use in adult mental health inpatient units: study protocol for a randomized controlled trial. <i>Trials</i> , 2014 , 15, 308	2.8	6

71	Don't turn your back on the symptoms of psychosis: a proof-of-principle, quasi-experimental public health trial to reduce the duration of untreated psychosis in Birmingham, UK. <i>BMC Psychiatry</i> , 2013 , 13, 67	4.2	6
70	Abuse and dysfunctional affiliations in childhood: an exploration of their impact on voice-hearers' appraisals of power and expressed emotion. <i>Psychosis</i> , 2012 , 4, 19-31	1.2	6
69	Collaborating to provide early-intervention services to persons in England with first-episode psychosis. <i>Psychiatric Services</i> , 2009 , 60, 1484-8	3.3	6
68	Controversies and Growing Points in Cognitive-Behavioural interventions for People with Schizophrenia. <i>Behavioural and Cognitive Psychotherapy</i> , 1992 , 20, 305-342	2.1	6
67	Pathways to emotional dysfunction in first-episode psychosis		6
66	Experiences and satisfaction of children, young people and their parents with alternative mental health models to inpatient settings: a systematic review. <i>European Child and Adolescent Psychiatry</i> , 2020 , 29, 1621-1633	5.5	6
65	Urban remediation: a new recovery-oriented strategy to manage urban stress after first-episode psychosis. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2020 , 55, 273-283	4.5	6
64	Evaluating substance misuse interventions in psychosis: The limitations of the RCT with 'patient' as the unit of analysis. <i>Journal of Mental Health</i> , 2001 , 10, 585-587	2.7	5
63	Expressed emotions and first episodes of schizophrenia. <i>British Journal of Psychiatry</i> , 1987 , 151, 859-60	5.4	5
62	The Trauma of First Episode Psychosis: The Role of Cognitive Mediation		5
61	Appraisals: Voices' power and purpose 2010 , 81-102		5
60	NICE v. SIGN on psychosis and schizophrenia: same roots, similar guidelines, different interpretations. <i>British Journal of Psychiatry</i> , 2016 , 208, 316-9	5.4	5
59	After the storm, Solar comes out: A new service model for children and adolescent mental health. <i>Microbial Biotechnology</i> , 2021 , 15, 731-738	3.3	5
58	Derivation of a prediction model for a diagnosis of depression in young adults: a matched case-control study using electronic primary care records. <i>Microbial Biotechnology</i> , 2018 , 12, 444-455	3.3	4
57	New directions in the psychosocial approach to the psychoses. <i>Journal of Mental Health</i> , 1998 , 7, 111-114.	4.7	4
56	TC5D EDEN: EVALUATING THE DEVELOPMENT AND IMPACT OF EARLY INTERVENTION SERVICES IN THE WEST MIDLANDS. <i>Schizophrenia Research</i> , 2006 , 86, S50	3.6	4
55	Early intervention and sustaining the management of vulnerability. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34, S181-S184	2.6	4
54	Using the crisis 1995 , 116-148		4

53	The COMMAND trial of cognitive therapy for harmful compliance with command hallucinations (CTCH): a qualitative study of acceptability and tolerability in the UK. <i>BMJ Open</i> , 2018 , 8, e021657	3	4
52	A Consultation-Liaison Service on Integrated Treatment: A Program Description. <i>Journal of Dual Diagnosis</i> , 2013 , 9, 149-157	2.4	3
51	Physical illnesses, developmental risk factors and psychiatric diagnoses among subjects at risk of psychosis. <i>European Psychiatry</i> , 2013 , 28, 135-40	6	3
50	Frequency and preventative interventions for non-suicidal self-injury and suicidal behaviour in primary school-age children: a scoping review protocol. <i>BMJ Open</i> , 2017 , 7, e017291	3	3
49	Cannabis use and abstinence in first-episode psychosis: the participants' view. <i>Mental Health and Substance Use: Dual Diagnosis</i> , 2013 , 6, 47-58		3
48	Exploring the attitudes of staff working within mental health settings toward clients who use cannabis. <i>Drugs: Education, Prevention and Policy</i> , 2009 , 16, 311-327	1.2	3
47	The impact of training within a comprehensive dual diagnosis strategy: the Combined Psychosis and Substance Use (COMPASS) experience. <i>Mental Health and Substance Use: Dual Diagnosis</i> , 2012 , 5, 206-216		3
46	Commentary on Garety & Freeman. I: 'Cognitive approaches to delusions--a critical review of theories and evidence'. <i>British Journal of Clinical Psychology</i> , 1999 , 38, 315-8	3.6	3
45	Community stakeholders' perspectives on youth mental health in India: Problems, challenges and recommendations. <i>Microbial Biotechnology</i> , 2021 , 15, 716-722	3.3	3
44	Cognitive Therapy for command hallucinations 2010 , 103-122		3
43	Predictors of social and role outcomes in first episode psychosis: A prospective 12-month study of social cognition, neurocognition and symptoms. <i>Microbial Biotechnology</i> , 2021 , 15, 993-1001	3.3	3
42	Measuring adherence in social recovery therapy with people with first episode psychosis. <i>Behavioural and Cognitive Psychotherapy</i> , 2020 , 48, 82-90	2.1	3
41	Interactive highly realistic virtual reality as a tool for understanding the genesis and treatment of psychotic symptoms. <i>Studies in Health Technology and Informatics</i> , 2009 , 144, 138-40	0.5	3
40	Early intervention and identification strategies for young people at risk of developing mental health issues: working in partnership with schools in Birmingham, UK. <i>Microbial Biotechnology</i> , 2017 , 11, 471-479	3.3	2
39	Protocol for a systematic review and meta-analysis of cognitive-behavioural therapy for social anxiety disorder in psychosis. <i>Systematic Reviews</i> , 2014 , 3, 62	3	2
38	Recruiting South Asians into a UK Mental Health Randomised Controlled Trial: Experiences of Field Researchers. <i>Journal of Racial and Ethnic Health Disparities</i> , 2014 , 1, 181-193	3.5	2
37	The impact of emotional acknowledgement and self-esteem on resilience to family dysfunction. <i>European Psychiatry</i> , 2017 , 41, S734-S735	6	2
36	Treating prolactinoma and psychosis: medication and cognitive behavioural therapy. <i>BMJ Case Reports</i> , 2011 , 2011,	0.9	2

35	Emotional and personality dysfunctions in early psychosis	283-302		2
34	FC4D IMPROVING PSYCHOLOGICAL ADJUSTMENT FOLLOWING A FIRST EPISODE OF PSYCHOSIS: A RANDOMISED CONTROLLED TRIAL OF COGNITIVE THERAPY TO REDUCE POST-PSYCHOTIC TRAUMA SYMPTOMS. <i>Schizophrenia Research</i> , 2006 , 86, S39		3.6	2
33	Psychotherapies for Schizophrenia: A Review	2003 , 153-247		2
32	Preventing the entrenchment of high expressed emotion in first episode psychosis: early developmental attachment pathways. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34, S191-S197		2.6	2
31	A case of monosymptomatic musical hallucinations. <i>Irish Journal of Psychological Medicine</i> , 1993 , 10, 89-90			2
30	Predicting and controlling relapse in schizophrenia: early signs monitoring	1992 , 293-308		2
29	Development and Validation of a Nonremission Risk Prediction Model in First-Episode Psychosis: An Analysis of 2 Longitudinal Studies. <i>Schizophrenia Bulletin Open</i> , 2021 , 2, sgab041		2.2	2
28	Knitting up the ravelled sleeve of care: sleep and psychosis. <i>Lancet Psychiatry</i> , 2015 , 2, 950-1		23.3	1
27	A Synthetic Literature Review on the Management of Emerging Treatment Resistance in First Episode Psychosis: Can We Move towards Precision Intervention and Individualised Care?. <i>Medicina (Lithuania)</i> , 2020 , 56,		3.1	1
26	The association between treatment beliefs and engagement in care in first episode psychosis. <i>Schizophrenia Research</i> , 2019 , 204, 409-410		3.6	1
25	'Schizophrenia postdrome': a study of low-level psychotic experience after remission of first-episode schizophrenia. <i>Microbial Biotechnology</i> , 2009 , 3, 296-9		3.3	1
24	Cognitive Theory and Therapy for Command Hallucinations. <i>Journal of Experimental Psychopathology</i> , 2012 , 3, 537-551		2.3	1
23	Staff attitude towards cocaine/crack-cocaine use amongst individuals with severe mental health problems in an inner city area of the UK. <i>Mental Health and Substance Use: Dual Diagnosis</i> , 2008 , 1, 205-215			1
22	Commentary on Tiffin et al.'s 'From commitment to reality - early intervention in psychosis services in England'. Reasons to be cheerful. <i>Microbial Biotechnology</i> , 2007 , 1, 109		3.3	1
21	Applying CBT to promote recovery from acute psychosis. <i>Psychological Medicine</i> , 2005 , 35, 152; author reply 152-3		6.9	1
20	Service innovations: Coexisting severe mental health and substance use problems: developing integrated services in the UK. <i>Psychiatric Bulletin</i> , 2003 , 27, 183-186			1
19	Mental Health Hospital Admissions: a Teachable Moment and Window of Opportunity To Promote Change in Drug and Alcohol Misuse. <i>International Journal of Mental Health and Addiction</i> , 2019 , 17, 22-40		8.8	1
18	Improved social functioning following social recovery therapy in first episode psychosis: Do social cognition and neurocognition change following therapy, and do they predict treatment response?. <i>Schizophrenia Research</i> , 2021 , 228, 249-255		3.6	1

17	Early Intervention Services Versus Generic Community Mental Health Services: A Paradigm Shift 2014 , 169-184		o
16	G493 How can health services effectively improve the mental health of young people leaving public care? the lync study. <i>Archives of Disease in Childhood</i> , 2015 , 100, A212.1-A212	2.2	o
15	Structure and stability of symptoms in first episode psychosis: a longitudinal network approach. <i>Translational Psychiatry</i> , 2021 , 11, 567	8.6	o
14	The impact of manic symptoms in first-episode psychosis: Findings from the UK National EDEN study. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 144, 358-367	6.5	o
13	Warwick-India-Canada (WIC) global mental health group: rationale, design and protocol. <i>BMJ Open</i> , 2021 , 11, e046362	3	o
12	Status of primary and secondary mental healthcare of people with severe mental illness: an epidemiological study from the UK PARTNERS2 programme. <i>BJPsych Open</i> , 2021 , 7, e53	5	o
11	Prognostic models in first-episode psychosis - Authors' reply. <i>The Lancet Digital Health</i> , 2020 , 2, e61	14.4	
10	The impact of the transition from primary school to secondary school on young adolescents. <i>European Psychiatry</i> , 2017 , 41, S179-S180	6	
9	Subjective quality of life and its changes in patients at-risk for psychosis results of the EPOS study. <i>European Psychiatry</i> , 2011 , 26, 2039-2039	6	
8	Course of psychopathology in the at-risk mental state - outcomes beyond transition to psychosis. <i>European Psychiatry</i> , 2011 , 26, 2037-2037	6	
7	Treatment in the at-risk mental state for psychosis - a comparison of four european countries. <i>European Psychiatry</i> , 2011 , 26, 2040-2040	6	
6	0366 SCOTTISH STUDY OF PSYCHOSIS: ENGAGEMENT, ATTACHMENT AND RECOVERY (SSP: EAR)-STUDY OUTLINE. <i>Schizophrenia Research</i> , 2006 , 86, S142-S143	3.6	
5	O-90 Randomised controlled trial of joint crisis plans to reduce compulsory treatment for people with psychosis: Clinical outcomes and implementation. <i>BMJ Supportive and Palliative Care</i> , 2015 , 5, A28.3-A28		
4	Understanding the Role of Emotion in Psychosis: Social Anxiety Disorder in First-Episode Psychosis 2011 , 89-110		
3	Duration of Untreated Psychosis and Pathways to Care9-16		
2	Cognitive Therapy for Reducing Distress and Harmful Compliance with Command Hallucinations13-34		
1	Treatment Targets in the Pre-psychotic Phase75-91		