

# Won Tae Kim

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5976079/publications.pdf>

Version: 2024-02-01

62  
papers

825  
citations

516710

16  
h-index

642732

23  
g-index

63  
all docs

63  
docs citations

63  
times ranked

1434  
citing authors

#	ARTICLE	IF	CITATIONS
1	TOX-expressing terminally exhausted tumor-infiltrating CD8+ T cells are reinvigorated by co-blockade of PD-1 and TIGIT in bladder cancer. <i>Cancer Letters</i> , 2021, 499, 137-147.	7.2	42
2	Urinary MicroRNAs of Prostate Cancer: Virus-Encoded hsv1-miRH18 and hsv2-miR-H9-5p Could Be Valuable Diagnostic Markers. <i>International Neurourology Journal</i> , 2015, 19, 74-84.	1.2	40
3	HSPA6 augments garlic extract-induced inhibition of proliferation, migration, and invasion of bladder cancer EJ cells; Implication for cell cycle dysregulation, signaling pathway alteration, and transcription factor-associated MMP-9 regulation. <i>PLoS ONE</i> , 2017, 12, e0171860.	2.5	39
4	Urinary cell-free microRNA biomarker could discriminate bladder cancer from benign hematuria. <i>International Journal of Cancer</i> , 2019, 144, 380-388.	5.1	30
5	Novel Combination Markers for Predicting Survival in Patients with Muscle Invasive Bladder Cancer: USP18 and DGCR2. <i>Journal of Korean Medical Science</i> , 2014, 29, 351.	2.5	29
6	Expression levels of FGFR3 as a prognostic marker for the progression of primary pT1 bladder cancer and its association with mutation status. <i>Oncology Letters</i> , 2017, 14, 3817-3824.	1.8	29
7	Prostate Size Correlates with Fasting Blood Glucose in Non-Diabetic Benign Prostatic Hyperplasia Patients with Normal Testosterone Levels. <i>Journal of Korean Medical Science</i> , 2011, 26, 1214.	2.5	28
8	MicroRNA-106a suppresses proliferation, migration, and invasion of bladder cancer cells by modulating MAPK signaling cell cycle regulators, and Ets-1-mediated MMP-2 expression. <i>Oncology Reports</i> , 2016, 36, 2421-2429.	2.6	27
9	MicroRNA-892b influences proliferation, migration and invasion of bladder cancer cells by mediating the p19ARF/cyclin D1/CDK6 and Sp-1/MMP-9 pathways. <i>Oncology Reports</i> , 2016, 36, 2313-2320.	2.6	25
10	The age-adjusted Charlson comorbidity index as a predictor of overall survival of surgically treated non-metastatic clear cell renal cell carcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 187-196.	2.5	24
11	p21WAF1 Is Required for Interleukin-16-Induced Migration and Invasion of Vascular Smooth Muscle Cells via the p38MAPK/Sp-1/MMP-9 Pathway. <i>PLoS ONE</i> , 2015, 10, e0142153.	2.5	23
12	Advances in urinary biomarker discovery in urological research. <i>Investigative and Clinical Urology</i> , 2020, 61, S8.	2.0	22
13	Hypertriglyceridemia Is Associated With Increased Risk for Stone Recurrence in Patients With Urolithiasis. <i>Urology</i> , 2014, 84, 766-771.	1.0	21
14	Collagen type $\alpha 1$ and 2 repress the proliferation, migration and invasion of bladder cancer cells. <i>International Journal of Oncology</i> , 2021, 59, .	3.3	21
15	Impact of Young Age at Diagnosis on Survival in Patients with Surgically Treated Renal Cell Carcinoma: a Multicenter Study. <i>Journal of Korean Medical Science</i> , 2016, 31, 1976.	2.5	20
16	Overexpression of caldesmon is associated with tumor progression in patients with primary non-muscle-invasive bladder cancer. <i>Oncotarget</i> , 2015, 6, 40370-40384.	1.8	20
17	<i>CDC6</i> mRNA Expression Is Associated with the Aggressiveness of Prostate Cancer. <i>Journal of Korean Medical Science</i> , 2018, 33, e303.	2.5	19
18	A Low Geriatric Nutritional Risk Index is Associated with Aggressive Pathologic Characteristics and Poor Survival after Nephrectomy in Clear Renal Cell Carcinoma: A Multicenter Retrospective Study. <i>Nutrition and Cancer</i> , 2020, 72, 88-97.	2.0	19

#	ARTICLE	IF	CITATIONS
19	Prognostic Impact of Nutritional Status Assessed by the Controlling Nutritional Status (CONUT) Score in Patients with Surgically Treated Renal Cell Carcinoma. <i>Nutrition and Cancer</i> , 2018, 70, 886-894.	2.0	18
20	Methylation Signature for Prediction of Progression Free Survival in Surgically Treated Clear Cell Renal Cell Carcinoma. <i>Journal of Korean Medical Science</i> , 2019, 34, e144.	2.5	17
21	Urinary microRNA-1913 to microRNA-3659 expression ratio as a non-invasive diagnostic biomarker for prostate cancer. <i>Investigative and Clinical Urology</i> , 2021, 62, 340.	2.0	14
22	Identification of C16orf74 as a Marker of Progression in Primary Non-Muscle Invasive Bladder Cancer. <i>PLoS ONE</i> , 2010, 5, e15260.	2.5	13
23	Role of 1,25-Dihydroxy Vitamin D <sub>3</sub> and Parathyroid Hormone in Urinary Calcium Excretion in Calcium Stone Formers. <i>Yonsei Medical Journal</i> , 2014, 55, 1326.	2.2	13
24	Effect of Renal Insufficiency on Stone Recurrence in Patients with Urolithiasis. <i>Journal of Korean Medical Science</i> , 2014, 29, 1132.	2.5	13
25	Urinary Cell-Free DNA IQGAP3/BMP4 Ratio as a Prognostic Marker for Non-Muscle-Invasive Bladder Cancer. <i>Clinical Genitourinary Cancer</i> , 2019, 17, e704-e711.	1.9	12
26	A prognostic immune predictor, HLA-DRA, plays diverse roles in non-muscle invasive and muscle invasive bladder cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021, 39, 237.e21-237.e29.	1.6	12
27	Increased Expression of Herpes Virus-Encoded hsv1-miR-H18 and hsv2-miR-H9-5p in Cancer-Containing Prostate Tissue Compared to That in Benign Prostate Hyperplasia Tissue. <i>International Neurourology Journal</i> , 2016, 20, 122-130.	1.2	12
28	<i>GSTT1</i> as a Prognosticator for Recurrence and Progression in Patients with Non-Muscle-Invasive Bladder Cancer. <i>Disease Markers</i> , 2010, 29, 81-87.	1.3	11
29	The association of benign prostatic hyperplasia with lower urinary tract stones in adult men: A retrospective multicenter study. <i>Asian Journal of Urology</i> , 2018, 5, 118-121.	1.2	11
30	Diagnostic value of combined IQGAP3/BMP4 and IQGAP3/FAM107A expression ratios in urinary cell-free DNA for discriminating bladder cancer from hematuria. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019, 37, 86-96.	1.6	11
31	Impact of the ASA Physical Status Score on Adjuvant Chemotherapy Eligibility and Survival of Upper Tract Urothelial Carcinoma Patients: a Multicenter Study. <i>Journal of Korean Medical Science</i> , 2017, 32, 335.	2.5	10
32	The predictive value of GSTT1 polymorphisms in predicting the early response to induction BCG therapy in patients with non-muscle invasive bladder cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 458-465.	1.6	9
33	Clinical, prognostic, and therapeutic significance of heat shock protein 27 in bladder cancer. <i>Oncotarget</i> , 2018, 9, 7961-7974.	1.8	9
34	A novel tumor suppressing gene, ARHGAP9, is an independent prognostic biomarker for bladder cancer. <i>Oncology Letters</i> , 2020, 19, 476-486.	1.8	9
35	Parathyroid hormone is not involved in prostate growth in patients with benign prostatic hyperplasia. <i>Prostate</i> , 2011, 71, 1210-1215.	2.3	8
36	FAM70B as a Novel Prognostic Marker for Cancer Progression and Cancer-Specific Death in Muscle-Invasive Bladder Cancer. <i>Korean Journal of Urology</i> , 2012, 53, 598.	1.2	8

#	ARTICLE	IF	CITATIONS
37	Decreased <i>DBC1</i> Expression Is Associated With Poor Prognosis in Patients With Non-Muscle-Invasive Bladder Cancer. <i>Korean Journal of Urology</i> , 2013, 54, 631.	1.2	8
38	Lower Levels of Human MOB3B Are Associated with Prostate Cancer Susceptibility and Aggressive Clinicopathological Characteristics. <i>Journal of Korean Medical Science</i> , 2015, 30, 937.	2.5	8
39	Clinical Implications and Prognostic Values of <i>Prostate Cancer Susceptibility Candidate</i> Methylation in Primary Nonmuscle Invasive Bladder Cancer. <i>Disease Markers</i> , 2015, 2015, 1-6.	1.3	8
40	Angiopoietin-like protein 4 potentiates DMS-induced inhibition of proliferation, migration, and invasion of bladder cancer EJ cells; involvement of G <sub>2</sub> /M-phase cell cycle arrest, signaling pathways, and transcription factors-mediated MMP-9 expression. <i>Food and Nutrition Research</i> , 2017, 61, 1338918.	2.6	8
41	Molecular Progression Risk Score for Prediction of Muscle Invasion in Primary T1 High-Grade Bladder Cancer. <i>Clinical Genitourinary Cancer</i> , 2018, 16, 274-280.	1.9	8
42	Twenty-four-hour urine osmolality as a representative index of adequate hydration and a predictor of recurrence in patients with urolithiasis. <i>International Urology and Nephrology</i> , 2019, 51, 1129-1135.	1.4	8
43	Metabolic Characteristics and Risks Associated with Stone Recurrence in Korean Young Adult Stone Patients. <i>Journal of Endourology</i> , 2017, 31, 806-811.	2.1	7
44	Chronological Trends in Clinical and Urinary Metabolic Features over 20 Years in Korean Urolithiasis Patients. <i>Journal of Korean Medical Science</i> , 2017, 32, 1496.	2.5	7
45	Effects of poloxamer-based thermo-sensitive sol-gel agent on urethral stricture after transurethral resection of the prostate for benign prostatic hyperplasia: a multicentre, single-blinded, randomised controlled trial. <i>BJU International</i> , 2020, 125, 160-167.	2.5	7
46	A novel urinary mRNA signature using the droplet digital polymerase chain reaction platform improves discrimination between prostate cancer and benign prostatic hyperplasia within the prostate-specific antigen gray zone. <i>Investigative and Clinical Urology</i> , 2020, 61, 411.	2.0	7
47	Prognostic Value of BUB1 for Predicting Non-Muscle-Invasive Bladder Cancer Progression. <i>International Journal of Molecular Sciences</i> , 2021, 22, 12756.	4.1	7
48	Long-term validation of a molecular progression-associated gene classifier for prediction of muscle invasion in primary non-muscle-invasive bladder cancer. <i>Oncology Letters</i> , 2017, 14, 2468-2474.	1.8	6
49	For Physicians Managing Voiding Dysfunction, Improving the Detection Rate of Early Prostate Cancer and Discrimination From Benign Prostatic Hyperplasia, in a Molecular Biomarker Aspects. <i>International Neurourology Journal</i> , 2019, 23, 5-12.	1.2	6
50	ZNF492 and GPR149 methylation patterns as prognostic markers for clear cell renal cell carcinoma: Array-based DNA methylation profiling. <i>Oncology Reports</i> , 2019, 42, 453-460.	2.6	6
51	The prognostic value of the pretreatment serum albumin to globulin ratio for predicting adverse pathology in patients undergoing radical prostatectomy for prostate cancer. <i>Investigative and Clinical Urology</i> , 2021, 62, 545.	2.0	6
52	Expression of hsv1-miR-H18 and hsv2-miR-H9 as a field defect marker for detecting prostate cancer. <i>Prostate International</i> , 2022, 10, 1-6.	2.3	5
53	Distinct Metabolic Characteristics and Risk of Stone Recurrence in Patients With Multiple Stones at the First-time Presentation. <i>Urology</i> , 2014, 84, 274-278.	1.0	4
54	A high basal metabolic rate is an independent predictor of stone recurrence in obese patients. <i>Investigative and Clinical Urology</i> , 2021, 62, 195.	2.0	4

#	ARTICLE	IF	CITATIONS
55	Impact of Transobturator Tape Treatment on Overactive Bladder Symptoms, Particularly Nocturia, in Patients With Mixed Urinary Incontinence. <i>Korean Journal of Urology</i> , 2014, 55, 520.	1.2	3
56	Urinary hsv2-miR-H9 to hsa-miR-3659 ratio is an effective marker for discriminating prostate cancer from benign prostate hyperplasia in patients within the prostate-specific antigen grey zone. <i>Investigative and Clinical Urology</i> , 2022, 63, 238.	2.0	3
57	Change in Prostate Specific Antigen Concentration in Men with Prostate Specific Antigen Less than 2.5 ng/ml Taking Low Dose Finasteride or Dutasteride for Male Androgenetic Alopecia. <i>Journal of Urology</i> , 2017, 198, 1340-1345.	0.4	2
58	Expression of phosphorylated p21-activated kinase 4 is associated with aggressive histologic characteristics and poor prognosis in patients with surgically treated renal cell carcinoma. <i>Investigative and Clinical Urology</i> , 2021, 62, 399.	2.0	2
59	Nutritional status assessed by the Controlling Nutritional Status (CONUT) score as a predictor of recurrence of urolithiasis. <i>Investigative and Clinical Urology</i> , 2021, 62, 553.	2.0	2
60	Trends in clinical, operative, and pathologic characteristics of surgically treated renal mass in a Korean center: A surgical series from 1988 through 2015. <i>Investigative and Clinical Urology</i> , 2019, 60, 184.	2.0	2
61	Expression of RPL9 predicts the recurrence of non-muscle invasive bladder cancer with BCG therapy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2022, , .	1.6	2
62	Effect of pre-operative internal obturator muscle mass index in MRI on biochemical recurrence of prostate cancer patients after radical prostatectomy: a multi-center study. <i>BMC Urology</i> , 2021, 21, 85.	1.4	1