

Fiona Crawford

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5983784/publications.pdf>

Version: 2024-02-01

19
papers

510
citations

840776

11
h-index

888059

17
g-index

19
all docs

19
docs citations

19
times ranked

733
citing authors

#	ARTICLE	IF	CITATIONS
1	Chronic Repetitive Mild Traumatic Brain Injury Results in Reduced Cerebral Blood Flow, Axonal Injury, Gliosis, and Increased T-Tau and Tau Oligomers. <i>Journal of Neuropathology and Experimental Neurology</i> , 2016, 75, 636-655.	1.7	104
2	Gulf War Agent Exposure Causes Impairment of Long-Term Memory Formation and Neuropathological Changes in a Mouse Model of Gulf War Illness. <i>PLoS ONE</i> , 2015, 10, e0119579.	2.5	68
3	Translational potential of long-term decreases in mitochondrial lipids in a mouse model of Gulf War Illness. <i>Toxicology</i> , 2016, 372, 22-33.	4.2	50
4	Primum non nocere: a call for balance when reporting on CTE. <i>Lancet Neurology</i> , The, 2019, 18, 231-233.	10.2	48
5	Spleen tyrosine kinase (SYK) blocks autophagic Tau degradation in vitro and in vivo. <i>Journal of Biological Chemistry</i> , 2019, 294, 13378-13395.	3.4	31
6	A permethrin metabolite is associated with adaptive immune responses in Gulf War Illness. <i>Brain, Behavior, and Immunity</i> , 2019, 81, 545-559.	4.1	31
7	APOE Genotype Specific Effects on the Early Neurodegenerative Sequelae Following Chronic Repeated Mild Traumatic Brain Injury. <i>Neuroscience</i> , 2019, 404, 297-313.	2.3	25
8	Targeting sirtuin activity with nicotinamide riboside reduces neuroinflammation in a GWI mouse model. <i>NeuroToxicology</i> , 2020, 79, 84-94.	3.0	23
9	Dopamine DRD2/Cys311 is not associated with chronic schizophrenia. , 1996, 67, 483-484.		22
10	The Vasoactivity of A β Peptides. <i>Annals of the New York Academy of Sciences</i> , 1997, 826, 35-46.	3.8	18
11	Occurrence of the Cys311 DRD2 variant in a pedigree multiply affected with panic disorder. <i>American Journal of Medical Genetics Part A</i> , 1995, 60, 332-334.	2.4	17
12	No interaction between the APOE and the alpha-1-antichymotrypsin genes on risk for Alzheimer's disease. , 1997, 74, 192-194.		17
13	Repeated unpredictable stress and social isolation induce chronic HPA axis dysfunction and persistent abnormal fear memory. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 104, 110035.	4.8	16
14	Apolipoprotein E isoforms differentially regulate matrix metalloproteinase 9 function in Alzheimer's disease. <i>Neurobiology of Aging</i> , 2020, 95, 56-68.	3.1	13
15	β -amyloid Peptides and Enhancement of Vasoconstriction by Endothelin-1. <i>Annals of the New York Academy of Sciences</i> , 1997, 826, 461-462.	3.8	9
16	Exogenous lipase administration alters gut microbiota composition and ameliorates Alzheimer's disease-like pathology in APP/PS1 mice. <i>Scientific Reports</i> , 2022, 12, 4797.	3.3	6
17	A fast, miniaturised <i>in-vitro</i> assay developed for quantification of lipase enzyme activity. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2019, 34, 1474-1480.	5.2	5
18	Alzheimer's disease is not associated with the hypertension genetic risk factors PLA2 or G protein β 3, either independently or interactively with apolipoprotein e. <i>American Journal of Medical Genetics Part A</i> , 1999, 88, 465-468.	2.4	4

#	ARTICLE	IF	CITATIONS
19	Candida rugosa lipase alters the gastrointestinal environment in wild-type mice. Biomedicine and Pharmacotherapy, 2020, 130, 110579.	5.6	3