

# Yuan Yuan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5984235/publications.pdf>

Version: 2024-02-01

52  
papers

1,548  
citations

257450

24  
h-index

330143

37  
g-index

52  
all docs

52  
docs citations

52  
times ranked

2033  
citing authors

#	ARTICLE	IF	CITATIONS
1	By restoring autophagic flux and improving mitochondrial function, corosolic acid protects against Dox-induced cardiotoxicity. <i>Cell Biology and Toxicology</i> , 2022, 38, 451-467.	5.3	16
2	Alteration of Autonomic Nervous System Is Associated With Severity and Outcomes in Patients With COVID-19. <i>Frontiers in Physiology</i> , 2021, 12, 630038.	2.8	50
3	High-Mobility Group A1 Promotes Cardiac Fibrosis by Upregulating FOXO1 in Fibroblasts. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 666422.	3.7	8
4	Long non-coding RNA Pvt1 modulates the pathological cardiac hypertrophy via miR-196b-mediated OSMR regulation. <i>Cellular Signalling</i> , 2021, 86, 110077.	3.6	7
5	Nucleotide-Binding Oligomerization Domain-Like Receptor 3 Deficiency Attenuated Isoproterenol-Induced Cardiac Fibrosis via Reactive Oxygen Species/High Mobility Group Box 1 Protein Axis. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 713.	3.7	8
6	The effect of HMGA1 in LPS-induced Myocardial Inflammation. <i>International Journal of Biological Sciences</i> , 2020, 16, 1798-1810.	6.4	26
7	High-mobility group AT-hook 1 promotes cardiac dysfunction in diabetic cardiomyopathy via autophagy inhibition. <i>Cell Death and Disease</i> , 2020, 11, 160.	6.3	32
8	Leukocyte immunoglobulin-like receptor B4 protects against cardiac hypertrophy via SHP-2-dependent inhibition of the NF- $\kappa$ B pathway. <i>Journal of Molecular Medicine</i> , 2020, 98, 691-705.	3.9	11
9	Corosolic acid attenuates cardiac fibrosis following myocardial infarction in mice. <i>International Journal of Molecular Medicine</i> , 2020, 45, 1425-1435.	4.0	7
10	Zingerone attenuates aortic banding-induced cardiac remodeling via activating the eNOS/Nrf2 pathway. <i>Journal of Cellular and Molecular Medicine</i> , 2019, 23, 6466-6478.	3.6	19
11	Protective role of berberine in isoprenaline-induced cardiac fibrosis in rats. <i>BMC Cardiovascular Disorders</i> , 2019, 19, 219.	1.7	13
12	Indigo Fruits Ingredient, Aucubin, Protects against LPS-Induced Cardiac Dysfunction in Mice. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2019, 371, 348-359.	2.5	20
13	The 5-Lipoxygenase Inhibitor Zileuton Protects Pressure Overload-Induced Cardiac Remodeling via Activating PPAR $\alpha$ . <i>Oxidative Medicine and Cellular Longevity</i> , 2019, 2019, 1-17.	4.0	15
14	The protective effect of high mobility group protein HMGA2 in pressure overload-induced cardiac remodeling. <i>Journal of Molecular and Cellular Cardiology</i> , 2019, 128, 160-178.	1.9	20
15	Long non-coding RNA cytoskeleton regulator RNA (CYTOR) modulates pathological cardiac hypertrophy through miR-155-mediated IKK $\alpha$ signaling. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2019, 1865, 1421-1427.	3.8	40
16	Corosolic acid ameliorates cardiac hypertrophy via regulating autophagy. <i>Bioscience Reports</i> , 2019, 39, .	2.4	14
17	TAX1BP1 overexpression attenuates cardiac dysfunction and remodeling in STZ-induced diabetic cardiomyopathy in mice by regulating autophagy. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018, 1864, 1728-1743.	3.8	51
18	Aucubin protects against pressure overload-induced cardiac remodeling via the $\beta$ 3-adrenoceptor-neuronal NOS cascades. <i>British Journal of Pharmacology</i> , 2018, 175, 1548-1566.	5.4	36

#	ARTICLE	IF	CITATIONS
19	Aucubin Protects against Myocardial Infarction-Induced Cardiac Remodeling via nNOS/NO-Regulated Oxidative Stress. <i>Oxidative Medicine and Cellular Longevity</i> , 2018, 2018, 1-15.	4.0	26
20	Role of autophagy in a model of obesity: A long-term high fat diet induces cardiac dysfunction. <i>Molecular Medicine Reports</i> , 2018, 18, 3251-3261.	2.4	20
21	Activating transcription factor 3 in cardiovascular diseases: a potential therapeutic target. <i>Basic Research in Cardiology</i> , 2018, 113, 37.	5.9	87
22	Syringin prevents cardiac hypertrophy induced by pressure overload through the attenuation of autophagy. <i>International Journal of Molecular Medicine</i> , 2017, 39, 199-207.	4.0	10
23	Cucurbitacin B Protects Against Pressure Overload Induced Cardiac Hypertrophy. <i>Journal of Cellular Biochemistry</i> , 2017, 118, 3899-3910.	2.6	43
24	Evodiamine attenuates TGF- $\beta$ 1-induced fibroblast activation and endothelial to mesenchymal transition. <i>Molecular and Cellular Biochemistry</i> , 2017, 430, 81-90.	3.1	28
25	Evodiamine Prevents Isoproterenol-Induced Cardiac Fibrosis by Regulating Endothelial-to-Mesenchymal Transition. <i>Planta Medica</i> , 2017, 83, 761-769.	1.3	26
26	Mechanisms contributing to cardiac remodelling. <i>Clinical Science</i> , 2017, 131, 2319-2345.	4.3	132
27	Sesamin prevents apoptosis and inflammation after experimental myocardial infarction by JNK and NF- $\kappa$ B pathways. <i>Food and Function</i> , 2017, 8, 2875-2885.	4.6	58
28	Acacetin protects against cardiac remodeling after myocardial infarction by mediating MAPK and PI3K/Akt signal pathway. <i>Journal of Pharmacological Sciences</i> , 2017, 135, 156-163.	2.5	32
29	Sesamin Protects Against Cardiac Remodeling Via Sirt3/ROS Pathway. <i>Cellular Physiology and Biochemistry</i> , 2017, 44, 2212-2227.	1.6	35
30	Baicalein protects against endothelial cell injury by inhibiting the TLR4/NF- $\kappa$ B signaling pathway. <i>Molecular Medicine Reports</i> , 2017, 17, 3085-3091.	2.4	11
31	Puerarin Protects against Cardiac Fibrosis Associated with the Inhibition of TGF- $\beta$ 1/Smad2-Mediated Endothelial-to-Mesenchymal Transition. <i>PPAR Research</i> , 2017, 2017, 1-14.	2.4	27
32	Mnk1 (Mitogen-Activated Protein Kinase-Interacting Kinase 1) Deficiency Aggravates Cardiac Remodeling in Mice. <i>Hypertension</i> , 2016, 68, 1393-1399.	2.7	30
33	Achievement of a target dose of bisoprolol may not be a preferred option for attenuating pressure overload-induced cardiac hypertrophy and fibrosis. <i>Experimental and Therapeutic Medicine</i> , 2016, 12, 2027-2038.	1.8	11
34	Puerarin attenuates the inflammatory response and apoptosis in LPS-stimulated cardiomyocytes. <i>Experimental and Therapeutic Medicine</i> , 2016, 11, 415-420.	1.8	38
35	OX40 regulates pressure overload-induced cardiac hypertrophy and remodelling via CD4+ T-cells. <i>Clinical Science</i> , 2016, 130, 2061-2071.	4.3	35
36	Protection against cardiac hypertrophy by geniposide involves the GLP-1 receptor / AMPK signalling pathway. <i>British Journal of Pharmacology</i> , 2016, 173, 1502-1516.	5.4	94

#	ARTICLE	IF	CITATIONS
37	Pleiotropic and puzzling effects of ATF3 in maladaptive cardiac remodeling. <i>International Journal of Cardiology</i> , 2016, 206, 87-88.	1.7	5
38	Sanguinarine inhibits angiotensin II-induced apoptosis in H9c2 cardiac cells via restoring reactive oxygen species-mediated decreases in the mitochondrial membrane potential. <i>Molecular Medicine Reports</i> , 2015, 12, 3400-3408.	2.4	20
39	3,3'-Diindolylmethane attenuates cardiac H9c2 cell hypertrophy through 5'-adenosine monophosphate-activated protein kinase- $\beta$ . <i>Molecular Medicine Reports</i> , 2015, 12, 1247-1252.	2.4	13
40	Pachymic acid protects H9c2 cardiomyocytes from lipopolysaccharide-induced inflammation and apoptosis by inhibiting the extracellular signal-regulated kinase 1/2 and p38 pathways. <i>Molecular Medicine Reports</i> , 2015, 12, 2807-2813.	2.4	25
41	Icariin protects H9c2 cardiomyocytes from lipopolysaccharide-induced injury via inhibition of the reactive oxygen species-dependent c-Jun N-terminal kinases/nuclear factor- $\kappa$ B pathway. <i>Molecular Medicine Reports</i> , 2015, 11, 4327-4332.	2.4	23
42	Naringenin attenuates pressure overload-induced cardiac hypertrophy. <i>Experimental and Therapeutic Medicine</i> , 2015, 10, 2206-2212.	1.8	34
43	Cathepsin B deficiency attenuates cardiac remodeling in response to pressure overload via TNF- $\alpha$ /ASK1/JNK pathway. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2015, 308, H1143-H1154.	3.2	71
44	Oleanolic acid alleviated pressure overload-induced cardiac remodeling. <i>Molecular and Cellular Biochemistry</i> , 2015, 409, 145-154.	3.1	23
45	Soluble ST2 may possess special superiority as a risk predictor in heart failure patients. <i>International Journal of Cardiology</i> , 2015, 186, 146-147.	1.7	5
46	DIM attenuates TGF- $\beta$ 1-induced myofibroblast differentiation in neonatal rat cardiac fibroblasts. <i>International Journal of Clinical and Experimental Pathology</i> , 2015, 8, 5121-8.	0.5	8
47	Icariin attenuates angiotensin II-induced hypertrophy and apoptosis in H9c2 cardiomyocytes by inhibiting reactive oxygen species-dependent JNK and p38 pathways. <i>Experimental and Therapeutic Medicine</i> , 2014, 7, 1116-1122.	1.8	33
48	Hesperetin attenuates mitochondria-dependent apoptosis in lipopolysaccharide-induced H9C2 cardiomyocytes. <i>Molecular Medicine Reports</i> , 2014, 9, 1941-1946.	2.4	36
49	ATF3 regulates multiple targets and may play a dual role in cardiac hypertrophy and injury. <i>International Journal of Cardiology</i> , 2014, 174, 838-839.	1.7	35
50	Puerarin attenuates pressure overload-induced cardiac hypertrophy. <i>Journal of Cardiology</i> , 2014, 63, 73-81.	1.9	73
51	Evidence for a Novel Autosomal Dominant Retinitis Pigmentosa Linked to Chromosome 1p22.1-q12 in a Chinese Family. <i>Current Eye Research</i> , 2011, 36, 154-167.	1.5	5
52	Molecular genetic analysis of a new form of spinocerebellar ataxia in a Chinese Han family. <i>Neuroscience Letters</i> , 2010, 479, 321-326.	2.1	3