

# Ulf Elbelt

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5989597/publications.pdf>

Version: 2024-02-01

35  
papers

1,572  
citations

304743

22  
h-index

315739

38  
g-index

38  
all docs

38  
docs citations

38  
times ranked

2535  
citing authors

#	ARTICLE	IF	CITATIONS
1	Circulating levels of irisin in patients with anorexia nervosa and different stages of obesity â€“ Correlation with body mass index. <i>Peptides</i> , 2013, 39, 125-130.	2.4	341
2	Irisin as a muscle-derived hormone stimulating thermogenesis â€“ A critical update. <i>Peptides</i> , 2014, 54, 89-100.	2.4	126
3	Diagnosis of Primary Hypophysitis in Germany. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 3841-3849.	3.6	98
4	Treatment of Primary Hypophysitis in Germany. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 3460-3469.	3.6	88
5	Molecular and Clinical Evidence for an <i>ARMC5</i> Tumor Syndrome: Concurrent Inactivating Germline and Somatic Mutations Are Associated With Both Primary Macronodular Adrenal Hyperplasia and Meningioma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, E119-E128.	3.6	85
6	Plasma bile acids show a positive correlation with body mass index and are negatively associated with cognitive restraint of eating in obese patients. <i>Frontiers in Neuroscience</i> , 2015, 9, 199.	2.8	79
7	Ghrelin and NUCB2/nesfatin-1 are expressed in the same gastric cell and differentially correlated with body mass index in obese subjects. <i>Histochemistry and Cell Biology</i> , 2013, 139, 909-918.	1.7	68
8	NUCB2/nesfatin-1 is associated with elevated scores of anxiety in female obese patients. <i>Psychoneuroendocrinology</i> , 2013, 38, 2502-2510.	2.7	57
9	The ghrelin activating enzyme ghrelin-O-acyltransferase (GOAT) is present in human plasma and expressed dependent on body mass index. <i>Peptides</i> , 2013, 43, 13-19.	2.4	51
10	Sex-specific regulation of NUCB2/nesfatin-1: Differential implication in anxiety in obese men and women. <i>Psychoneuroendocrinology</i> , 2015, 60, 130-137.	2.7	50
11	NUCB2/nesfatin-1 Is Associated with Elevated Levels of Anxiety in Anorexia Nervosa. <i>PLoS ONE</i> , 2015, 10, e0132058.	2.5	45
12	Differences of energy expenditure and physical activity patterns in subjects with various degrees of obesity. <i>Clinical Nutrition</i> , 2010, 29, 766-772.	5.0	44
13	Irisin Levels are Not Affected by Physical Activity in Patients with Anorexia Nervosa. <i>Frontiers in Endocrinology</i> , 2014, 4, 202.	3.5	40
14	Plasma kisspeptin and ghrelin levels are independently correlated with physical activity in patients with anorexia nervosa. <i>Appetite</i> , 2017, 108, 141-150.	3.7	38
15	Determinants of Weight Loss following Laparoscopic Sleeve Gastrectomy: The Role of Psychological Burden, Coping Style, and Motivation to Undergo Surgery. <i>Journal of Obesity</i> , 2015, 2015, 1-10.	2.7	34
16	Phoenixin is negatively associated with anxiety in obese men. <i>Peptides</i> , 2017, 88, 32-36.	2.4	34
17	Efficacy of Temozolomide Therapy in Patients With Aggressive Pituitary Adenomas and Carcinomasâ€“A German Survey. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e660-e675.	3.6	34
18	Altered insulin requirement in patients with type 1 diabetes and primary adrenal insufficiency receiving standard glucocorticoid replacement therapy. <i>European Journal of Endocrinology</i> , 2009, 160, 919-924.	3.7	31

#	ARTICLE	IF	CITATIONS
19	Heterozygous inactivating CaSR mutations causing neonatal hyperparathyroidism: function, inheritance and phenotype. <i>European Journal of Endocrinology</i> , 2016, 175, 421-431.	3.7	31
20	Irisin. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2013, 16, 541-547.	2.5	29
21	Determinants of Successful Weight Loss After Using a Commercial Web-Based Weight Reduction Program for Six Months: Cohort Study. <i>Journal of Medical Internet Research</i> , 2013, 15, e219.	4.3	28
22	Intraoperative indocyanine green videoangiography for identification of pituitary adenomas using a microscopic transsphenoidal approach. <i>Pituitary</i> , 2015, 18, 613-620.	2.9	23
23	Alterations of circulating NUCB2/nesfatin-1 during short term therapeutic improvement of anxiety in obese inpatients. <i>Psychoneuroendocrinology</i> , 2017, 79, 107-115.	2.7	20
24	Abdominal fat distribution differently affects muscle strength of the upper and lower extremities in women. <i>European Journal of Clinical Nutrition</i> , 2017, 71, 372-376.	2.9	17
25	Leptin and Physical Activity in Adult Patients with Anorexia Nervosa: Failure to Demonstrate a Simple Linear Association. <i>Nutrients</i> , 2017, 9, 1210.	4.1	14
26	Palliative treatment of uncontrollable hypercalcemia due to parathyrotoxicosis: denosumab as rescue therapy. <i>Endocrinology, Diabetes and Metabolism Case Reports</i> , 2015, 2015, 150082.	0.5	13
27	The Role of Objectively Measured, Altered Physical Activity Patterns for Body Mass Index Change during Inpatient Treatment in Female Patients with Anorexia Nervosa. <i>Journal of Clinical Medicine</i> , 2018, 7, 289.	2.4	11
28	Surgically and Conservatively Treated Obese Patients Differ in Psychological Factors, Regardless of Body Mass Index or Obesity-Related Co-Morbidities: A Comparison between Groups and an Analysis of Predictors. <i>PLoS ONE</i> , 2015, 10, e0117460.	2.5	9
29	Evaluation of a Portable Armband Device to Assess Resting Energy Expenditure in Patients With Anorexia Nervosa. <i>Nutrition in Clinical Practice</i> , 2016, 31, 362-367.	2.4	6
30	Essential Polyunsaturated Fatty Acids in Blood from Patients with and without Catheter-Proven Coronary Artery Disease. <i>International Journal of Molecular Sciences</i> , 2022, 23, 766.	4.1	6
31	Associations of physical activity with depressiveness and coping in subjects with high-grade obesity aiming at bariatric surgery: a cross-sectional study. <i>BioPsychoSocial Medicine</i> , 2015, 9, 16.	2.1	5
32	Association of Interleukin-1 Beta and Interleukin-1 Receptor Antagonist Gene Polymorphisms and Plasma Levels with Diabetic Nephropathy. <i>BioMed Research International</i> , 2022, 2022, 1-7.	1.9	5
33	Self-Directed Weight Loss Strategies: Energy Expenditure Due to Physical Activity Is Not Increased to Achieve Intended Weight Loss. <i>Nutrients</i> , 2015, 7, 5868-5888.	4.1	4
34	NUCB2/nesfatin-1 is associated with severity of eating disorder symptoms in female patients with obesity. <i>Psychoneuroendocrinology</i> , 2022, 143, 105842.	2.7	2
35	Estimating Resting Energy Expenditure With a Portable Armband Device in an Ambulatory Setting. <i>Nutrition in Clinical Practice</i> , 2012, 27, 825-831.	2.4	1