

Andre Henrique Freiria-Oliveira

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5989874/publications.pdf>

Version: 2024-02-01

15
papers

127
citations

1307594

7
h-index

1281871

11
g-index

15
all docs

15
docs citations

15
times ranked

174
citing authors

#	ARTICLE	IF	CITATIONS
1	Medullary Noradrenergic Neurons Mediate Hemodynamic Responses to Osmotic and Volume Challenges. <i>Frontiers in Physiology</i> , 2021, 12, 649535.	2.8	3
2	Forced internal desynchrony induces cardiometabolic alterations in adult rats. <i>Journal of Endocrinology</i> , 2019, 242, 25-36.	2.6	7
3	Role of the Carotid Bodies in the Hypertensive and Natriuretic Responses to NaCl Load in Conscious Rats. <i>Frontiers in Physiology</i> , 2018, 9, 1690.	2.8	2
4	Clinical data and risk factors for diabetic nephropathy in Brazilian central population. <i>Data in Brief</i> , 2018, 21, 1315-1320.	1.0	4
5	Median Preoptic Nucleus Mediates the Cardiovascular Recovery Induced by Hypertonic Saline in Hemorrhagic Shock. <i>Scientific World Journal</i> , The, 2014, 2014, 1-9.	2.1	7
6	Involvement of the median preoptic nucleus in blood pressure control. <i>Neuroscience Letters</i> , 2014, 558, 91-96.	2.1	9
7	Increased Expression of Angiotensin II Type 2 Receptors in the Solitary Vagal Complex Blunts Renovascular Hypertension. <i>Hypertension</i> , 2014, 64, 777-783.	2.7	35
8	Macrophage migration inhibitory factor in the nucleus of solitary tract decreases blood pressure in SHR. <i>Cardiovascular Research</i> , 2013, 97, 153-160.	3.8	16
9	A1 Noradrenergic Neurons Lesions Reduce Natriuresis and Hypertensive Responses to Hypernatremia in Rats. <i>PLoS ONE</i> , 2013, 8, e73187.	2.5	11
10	A2 Noradrenergic Lesions Prevent Renal Sympathoinhibition Induced by Hypernatremia in Rats. <i>PLoS ONE</i> , 2012, 7, e37587.	2.5	18
11	Angiotensin type 2 receptors (AT2R) over expression in the nucleus of the solitary tract (NTS) attenuate renovascular hypertension. <i>FASEB Journal</i> , 2012, 26, 1091.15.	0.5	0
12	Role of central angiotensinergic mechanisms on the facilitation of the recovery of hemorrhage-induced hypotension by noradrenergic A2 lesions. <i>FASEB Journal</i> , 2010, 24, 794.8.	0.5	1
13	Lesions of medullary catecholaminergic neurons increase salt intake in rats. <i>Brain Research Bulletin</i> , 2008, 76, 572-578.	3.0	13
14	A2 noradrenergic neurons inhibit osmoreceptor-induced pressor responses. <i>FASEB Journal</i> , 2008, 22, .	0.5	1
15	Pressor responses produced by peripheral osmoreceptor activation in commissural nucleus of the solitary tract-lesioned rats. <i>FASEB Journal</i> , 2008, 22, 738.2.	0.5	0