

Marcos Iglesias Lozano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5995750/publications.pdf>

Version: 2024-02-01

17
papers

235
citations

1040056

9
h-index

996975

15
g-index

21
all docs

21
docs citations

21
times ranked

484
citing authors

#	ARTICLE	IF	CITATIONS
1	Gluconate-Lactobionate-Dextran Perfusion Solutions Attenuate Ischemic Injury and Improve Function in a Murine Cardiac Transplant Model. <i>Cells</i> , 2022, 11, 1653.	4.1	1
2	A short course of tofacitinib sustains the immunoregulatory effect of CTLA4-Ig in the presence of inflammatory cytokines and promotes long-term survival of murine cardiac allografts. <i>American Journal of Transplantation</i> , 2021, 21, 2675-2687.	4.7	5
3	Noninvasive evaluation of intragraft immune responses in upper extremity transplantation. <i>Transplant International</i> , 2021, 34, 894-905.	1.6	2
4	Multiphase Assembly of Small Molecule Microcrystalline Peptide Hydrogel Allows Immunomodulatory Combination Therapy for Long-Term Heart Transplant Survival. <i>Small</i> , 2020, 16, e2002791.	10.0	15
5	Therapeutic Effects of Anti-Bone Morphogenetic Protein and Activin Membrane-Bound Inhibitor Treatment in Psoriasis and Arthritis. <i>Arthritis and Rheumatology</i> , 2020, 72, 1547-1558.	5.6	3
6	Type-I Interferons Inhibit Interleukin-10 Signaling and Favor Type 1 Diabetes Development in Nonobese Diabetic Mice. <i>Frontiers in Immunology</i> , 2018, 9, 1565.	4.8	13
7	Anti-IL17 treatment ameliorates Down syndrome phenotypes in mice. <i>Brain, Behavior, and Immunity</i> , 2018, 73, 235-251.	4.1	21
8	Context-dependent regulation of Th17-associated genes and IFN γ expression by the transcription factor NFAT5. <i>Immunology and Cell Biology</i> , 2017, 95, 56-67.	2.3	27
9	Bone Morphogenetic Protein and Activin Membrane-Bound Inhibitor, a Transforming Growth Factor β 2 Rheostat That Controls Murine Treg Cell/Th17 Cell Differentiation and the Development of Autoimmune Arthritis by Reducing Interleukin-2 Signaling. <i>Arthritis and Rheumatology</i> , 2016, 68, 1551-1562.	5.6	17
10	Selective Impairment of TH17-Differentiation and Protection against Autoimmune Arthritis after Overexpression of BCL2A1 in T Lymphocytes. <i>PLoS ONE</i> , 2016, 11, e0159714.	2.5	9
11	Distinct serum proteome profiles associated with collagen-induced arthritis and complete Freund's adjuvant-induced inflammation in CD38 ^{+/+} mice: The discriminative power of protein species or proteoforms. <i>Proteomics</i> , 2015, 15, 3382-3393.	2.2	6
12	p27 ^{Kip1} inhibits systemic autoimmunity through the control of Treg cell activity and differentiation. <i>Arthritis and Rheumatism</i> , 2013, 65, 343-354.	6.7	12
13	Mice Deficient in CD38 Develop an Attenuated Form of Collagen Type II-Induced Arthritis. <i>PLoS ONE</i> , 2012, 7, e33534.	2.5	36
14	Exacerbation of type II collagen-induced arthritis in apolipoprotein E-deficient mice in association with the expansion of Th1 and Th17 cells. <i>Arthritis and Rheumatism</i> , 2011, 63, 971-980.	6.7	31
15	GITR contributes to the systemic adjuvanticity of the <i>Escherichia coli</i> heat-labile enterotoxin. <i>European Journal of Immunology</i> , 2010, 40, 754-763.	2.9	3
16	B-cell overexpression of Bcl-2 cooperates with p21 deficiency for the induction of autoimmunity and lymphomas. <i>Journal of Autoimmunity</i> , 2010, 35, 316-324.	6.5	8
17	Increased Goodpasture Antigen-Binding Protein Expression Induces Type IV Collagen Disorganization and Deposit of Immunoglobulin A in Glomerular Basement Membrane. <i>American Journal of Pathology</i> , 2007, 171, 1419-1430.	3.8	25