
Stefanie A De Boer

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5999907/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	[18F]FDG Uptake in Adipose Tissue Is Not Related to Inflammation in Type 2 Diabetes Mellitus. Molecular Imaging and Biology, 2021, 23, 117-126.	2.6	8
2	Plasma Nitrate Levels Are Related to Metabolic Syndrome and Are Not Altered by Treatment with DPP-4 Inhibitor Linagliptin: A Randomised, Placebo-Controlled Trial in Patients with Early Type 2 Diabetes Mellitus. Antioxidants, 2021, 10, 1548.	5.1	2
3	Angiogenic T cells are decreased in people with type 2 diabetes mellitus and recruited by the dipeptidyl peptidaseâ€4 inhibitor Linagliptin: A subanalysis from a randomized, placeboâ€controlled trial (RELEASE) Tj ETQq1	. 4 .0. 7843	5 1 ⁄4 rgBT /⊖∖
4	Performance Evaluation of a Semi-automated Method for [18F]FDG Uptake in Abdominal Visceral Adipose Tissue. Molecular Imaging and Biology, 2019, 21, 159-167.	2.6	3
5	Visceral adipose tissue volume is associated with premature atherosclerosis in early type 2 diabetes mellitus independent of traditional risk factors. Atherosclerosis, 2019, 290, 87-93.	0.8	20
6	Effect of Linagliptin onÂArterial 18 F-Fluorodeoxyglucose Positron Emission Tomography Uptake. Journal of the American College of Cardiology, 2017, 69, 1097-1098.	2.8	8
7	Effect of linagliptin on pulse wave velocity in early type 2 diabetes: <scp>A</scp> randomized, doubleâ€blind, controlled 26â€week trial (<scp>RELEASE</scp>). Diabetes, Obesity and Metabolism, 2017, 19, 1147-1154.	4.4	33
8	Arterial Stiffness Is Positively Associated With 18F-fluorodeoxyglucose Positron Emission Tomography–Assessed Subclinical Vascular Inflammation in People With Early Type 2 Diabetes. Diabetes Care, 2016, 39, 1440-1447.	8.6	34

2