

Borje Haraldsson

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6002115/borje-haraldsson-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

153
papers

6,073
citations

40
h-index

72
g-index

163
ext. papers

6,697
ext. citations

7.1
avg. IF

5.68
L-index

#	Paper	IF	Citations
153	The Glomerular Endothelium Restricts Albumin Filtration.. <i>Frontiers in Medicine</i> , 2021 , 8, 766689	4.9	2
152	Proteoglycans contribute to the functional integrity of the glomerular endothelial cell surface layer and are regulated in diabetic kidney disease. <i>Scientific Reports</i> , 2021 , 11, 8487	4.9	3
151	Single-dwell treatment with a low-sodium solution in hypertensive peritoneal dialysis patients. <i>Peritoneal Dialysis International</i> , 2020 , 40, 446-454	2.8	1
150	Endothelin receptor-A mediates degradation of the glomerular endothelial surface layer via pathologic crosstalk between activated podocytes and glomerular endothelial cells. <i>Kidney International</i> , 2019 , 96, 957-970	9.9	27
149	Small-molecule factor B inhibitor for the treatment of complement-mediated diseases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 7926-7931	11.5	45
148	Patients' Perceptions and Factors Affecting Dialysis Modality Decisions. <i>Peritoneal Dialysis International</i> , 2018 , 38, 334-342	2.8	6
147	Amplification of the Melanocortin-1 Receptor in Nephrotic Syndrome Identifies a Target for Podocyte Cytoskeleton Stabilization. <i>Scientific Reports</i> , 2018 , 8, 15731	4.9	6
146	Glomerular Endothelial Mitochondrial Dysfunction Is Essential and Characteristic of Diabetic Kidney Disease Susceptibility. <i>Diabetes</i> , 2017 , 66, 763-778	0.9	106
145	Glycaemic control and excess risk of major coronary events in persons with type 1 diabetes. <i>Heart</i> , 2017 , 103, 1687-1695	5.1	27
144	Transcriptomic and Proteomic Profiling Provides Insight into Mesangial Cell Function in IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2017 , 28, 2961-2972	12.7	29
143	Long-term clinical outcome for patients poisoned by the fungal nephrotoxin orellanine. <i>BMC Nephrology</i> , 2017 , 18, 121	2.7	6
142	Orellanine specifically targets renal clear cell carcinoma. <i>Oncotarget</i> , 2017 , 8, 91085-91098	3.3	11
141	Melanocortin 1 receptor agonist protects podocytes through catalase and RhoA activation. <i>American Journal of Physiology - Renal Physiology</i> , 2016 , 310, F846-56	4.3	27
140	Mesangial cells from patients with IgA nephropathy have increased susceptibility to galactose-deficient IgA1. <i>BMC Nephrology</i> , 2016 , 17, 40	2.7	16
139	Treatment pattern in patients with idiopathic membranous nephropathy-practices in Sweden at the start of the millennium. <i>CKJ: Clinical Kidney Journal</i> , 2016 , 9, 227-33	4.5	2
138	Assessing the Accuracy of Continuous Glucose Monitoring (CGM) Calibrated With Capillary Values Using Capillary or Venous Glucose Levels as a Reference. <i>Journal of Diabetes Science and Technology</i> , 2016 , 10, 876-84	4.1	19
137	Continuous Glucose Monitoring in 2015. <i>Diabetes Technology and Therapeutics</i> , 2016 , 18 Suppl 1, S10-21	8.1	0

136	Perinatal DDT Exposure Induces Hypertension and Cardiac Hypertrophy in Adult Mice. <i>Environmental Health Perspectives</i> , 2016 , 124, 1722-1727	8.4	12
135	Biokinetic Modeling and Dosimetry for Optimizing Intraperitoneal Radioimmunotherapy of Ovarian Cancer Microtumors. <i>Journal of Nuclear Medicine</i> , 2016 , 57, 594-600	8.9	16
134	The relationship between three eGFR formulas and hospitalization for heart failure in 54 486 individuals with type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2016 , 32, 730-735	7.5	9
133	Decreased eGFR as a Risk Factor for Heart Failure in 13 781 Individuals With Type 1 Diabetes. <i>Journal of Diabetes Science and Technology</i> , 2015 , 10, 131-6	4.1	6
132	Predicting the effectiveness of insulin pump therapy on glycemic control in clinical practice: a retrospective study of patients with type 1 diabetes from 10 outpatient diabetes clinics in Sweden over 5 years. <i>Diabetes Technology and Therapeutics</i> , 2015 , 17, 21-8	8.1	17
131	Urinary albumin excretion in healthy adults: a cross sectional study of 24-hour versus timed overnight samples and impact of GFR and other personal characteristics. <i>BMC Nephrology</i> , 2015 , 16, 8	2.7	6
130	Glomerular IgG subclasses in idiopathic and malignancy-associated membranous nephropathy. <i>CKJ: Clinical Kidney Journal</i> , 2015 , 8, 433-9	4.5	25
129	The incidence of albuminuria after bariatric surgery and usual care in Swedish Obese Subjects (SOS): a prospective controlled intervention trial. <i>International Journal of Obesity</i> , 2015 , 39, 169-75	5.5	54
128	A clinical trial of the accuracy and treatment experience of the Dexcom G4 sensor (Dexcom G4 system) and Enlite sensor (guardian REAL-time system) tested simultaneously in ambulatory patients with type 1 diabetes. <i>Diabetes Technology and Therapeutics</i> , 2014 , 16, 759-67	8.1	66
127	Endothelial mitochondrial oxidative stress determines podocyte depletion in segmental glomerulosclerosis. <i>Journal of Clinical Investigation</i> , 2014 , 124, 1608-21	15.9	175
126	The endothelium as part of the integrative glomerular barrier complex. <i>Kidney International</i> , 2014 , 85, 8-11	9.9	24
125	Rapid increase of interleukin-10 plasma levels after combined auxiliary liver-kidney transplantation in presensitized patients. <i>Transplantation</i> , 2014 , 98, 208-15	1.8	12
124	Effects of melanocortin 1 receptor agonists in experimental nephropathies. <i>PLoS ONE</i> , 2014 , 9, e87816	3.7	15
123	A Retrospective Study in 5,989 Patients with Type 1 Diabetes in 10 Outpatient Diabetes Clinics in Sweden of the Frequency of Measuring HbA1c in Clinical Practice. <i>Journal of Diabetes & Metabolism</i> , 2014 , 05,	0	3
122	Modulation of microvascular permeability in the preovulatory rat ovary by an ovulatory gonadotropin stimulus. <i>Fertility and Sterility</i> , 2013 , 99, 903-9	4.8	7
121	A meta-analysis of expression signatures in glomerular disease. <i>Kidney International</i> , 2013 , 84, 591-9	9.9	3
120	Reactive oxygen species modulate the barrier function of the human glomerular endothelial glycocalyx. <i>PLoS ONE</i> , 2013 , 8, e55852	3.7	102
119	Analysis of the mushroom nephrotoxin orellanine and its glucosides. <i>Journal of Natural Products</i> , 2012 , 75, 1690-6	4.9	18

118	Hemodiafiltration improves plasma 25-hepcidin levels: a prospective, randomized, blinded, cross-over study comparing hemodialysis and hemodiafiltration. <i>Nephron Extra</i> , 2012 , 2, 55-65		18
117	Evaluation of effects on the peritoneum after intraperitoneal β -radioimmunotherapy with (211)At. <i>Cancer Biotherapy and Radiopharmaceuticals</i> , 2012 , 27, 353-64	3.9	10
116	The glomerular endothelium: new insights on function and structure. <i>Current Opinion in Nephrology and Hypertension</i> , 2012 , 21, 258-63	3.5	63
115	The glomerular endothelial cell coat is essential for glomerular filtration. <i>Kidney International</i> , 2011 , 79, 1322-30	9.9	89
114	Postischemic inflammatory response in an auxiliary liver graft predicts renal graft outcome in sensitized patients. <i>Transplantation</i> , 2011 , 91, 888-94	1.8	20
113	High glucose causes dysfunction of the human glomerular endothelial glycocalyx. <i>American Journal of Physiology - Renal Physiology</i> , 2011 , 300, F40-8	4.3	92
112	Dosing of erythropoiesis-stimulating agents can be reduced by a new administration regimen. <i>Nephron Extra</i> , 2011 , 1, 45-54		
111	Acute oxidative stress following intravenous iron injection in patients on chronic hemodialysis: a comparison of iron-sucrose and iron-dextran. <i>Nephron Clinical Practice</i> , 2011 , 118, c249-56		21
110	Role of glomerular proteoglycans in IgA nephropathy. <i>PLoS ONE</i> , 2011 , 6, e18575	3.7	29
109	Melanocortin 1 receptor agonists reduce proteinuria. <i>Journal of the American Society of Nephrology: JASN</i> , 2010 , 21, 1290-8	12.7	84
108	Tubular reabsorption of albumin: it's all about cubilin. <i>Journal of the American Society of Nephrology: JASN</i> , 2010 , 21, 1810-2	12.7	8
107	Impaired glomerular and tubular antioxidative defense mechanisms in nephrotic syndrome. <i>American Journal of Physiology - Renal Physiology</i> , 2010 , 299, F898-904	4.3	23
106	Permeability, ultrastructural changes, and distribution of novel proteins in the glomerular barrier in early puromycin aminonucleoside nephrosis. <i>Nephron Experimental Nephrology</i> , 2010 , 116, e42-52		9
105	Glomerular filtration rate after alpha-radioimmunotherapy with 211At-MX35-F(ab) γ 2: a long-term study of renal function in nude mice. <i>Cancer Biotherapy and Radiopharmaceuticals</i> , 2009 , 24, 649-58	3.9	24
104	Adriamycin alters glomerular endothelium to induce proteinuria. <i>Journal of the American Society of Nephrology: JASN</i> , 2009 , 20, 114-22	12.7	120
103	Glomerular filtration barrier. <i>Current Opinion in Nephrology and Hypertension</i> , 2009 , 18, 331-5	3.5	57
102	Is indoleamine 2,3-dioxygenase important for graft acceptance in highly sensitized patients after combined auxiliary liver-kidney transplantation?. <i>Transplantation</i> , 2009 , 88, 911-9	1.8	18
101	Citrate supplementation of PD fluid: effects on net ultrafiltration and clearance of small solutes in single dwells. <i>Nephrology Dialysis Transplantation</i> , 2009 , 24, 286-92	4.3	8

100	The fungal nephrotoxin orellanine simultaneously increases oxidative stress and down-regulates cellular defenses. <i>Free Radical Biology and Medicine</i> , 2008 , 44, 1562-9	7.8	21
99	Ascorbyl free radical reflects catalytically active iron after intravenous iron saccharate injection. <i>Free Radical Biology and Medicine</i> , 2008 , 45, 1302-7	7.8	6
98	Positive effects of protein restriction in patients with chronic kidney disease. <i>Journal of Renal Nutrition</i> , 2008 , 18, 269-80	3	14
97	Properties of the glomerular barrier and mechanisms of proteinuria. <i>Physiological Reviews</i> , 2008 , 88, 451-87	47.9	594
96	Resolved: normal glomeruli filter nephrotic levels of albumin. <i>Journal of the American Society of Nephrology: JASN</i> , 2008 , 19, 427-32	12.7	74
95	VEGF inhibition and renal thrombotic microangiopathy. <i>New England Journal of Medicine</i> , 2008 , 359, 205-6; author reply 206-7	59.2	20
94	In Vivo Peritoneal Surface Area Measurement in Rats by Micro-Computed Tomography (µCT). <i>Peritoneal Dialysis International</i> , 2008 , 28, 188-194	2.8	9
93	Changes in muscarinic receptors in the toad urothelial cell line TBM-54 following acrolein treatment. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2008 , 35, 217-22	3	3
92	Methodological issues on the use of urinary alpha-1-microglobuline in epidemiological studies. <i>Nephrology Dialysis Transplantation</i> , 2008 , 23, 1252-6	4.3	21
91	Mild renal ischemia-reperfusion reduces charge and size selectivity of the glomerular barrier. <i>American Journal of Physiology - Renal Physiology</i> , 2007 , 292, F1802-9	4.3	37
90	Functional and molecular alterations of the glomerular barrier in long-term diabetes in mice. <i>Diabetologia</i> , 2006 , 49, 2200-9	10.3	44
89	Beneficial effects of orosomucoid on the glomerular barrier in puromycin aminonucleoside-induced nephrosis. <i>Nephrology Dialysis Transplantation</i> , 2006 , 21, 1223-30	4.3	15
88	Morphological and functional evidence for an important role of the endothelial cell glycocalyx in the glomerular barrier. <i>American Journal of Physiology - Renal Physiology</i> , 2006 , 290, F111-6	4.3	149
87	Podocyte proteoglycan synthesis is involved in the development of nephrotic syndrome. <i>American Journal of Physiology - Renal Physiology</i> , 2006 , 291, F722-30	4.3	36
86	N-acetylcysteine attenuates kidney injury in rats subjected to renal ischaemia-reperfusion. <i>Nephrology Dialysis Transplantation</i> , 2006 , 21, 1240-7	4.3	83
85	Physiological Properties of the Peritoneum in an Adult Peritoneal Dialysis Population over a Three-Year Period. <i>Peritoneal Dialysis International</i> , 2006 , 26, 482-489	2.8	15
84	Prenatal exposure to interleukin-6 results in hypertension and alterations in the renin-angiotensin system of the rat. <i>Journal of Physiology</i> , 2006 , 575, 855-67	3.9	26
83	A new method for monitoring nitric oxide production using Teflon membrane microdialysis. <i>Free Radical Biology and Medicine</i> , 2005 , 39, 249-56	7.8	8

82	The Influence of Peritoneal Surface Area on Dialysis Adequacy. <i>Peritoneal Dialysis International</i> , 2005 , 25, 137-140	2.8	11
81	Primary human glomerular endothelial cells produce proteoglycans, and puromycin affects their posttranslational modification. <i>American Journal of Physiology - Renal Physiology</i> , 2005 , 288, F748-56	4.3	37
80	Measurement by magnetic resonance imaging of the peritoneal membrane in contact with dialysate in rats. <i>Advances in Peritoneal Dialysis Conference on Peritoneal Dialysis</i> , 2005 , 21, 17-20		7
79	Effect of peritoneal dialysis fluid composition on peritoneal area available for exchange in children. <i>Nephrology Dialysis Transplantation</i> , 2004 , 19, 925-32	4.3	43
78	Electron microscopic evaluation of the endothelial surface layer of glomerular capillaries. <i>Microvascular Research</i> , 2004 , 67, 9-17	3.7	77
77	Effects of norepinephrine or prostaglandin E2 on extracellular acidification rate of MCG 101, or K1735-M2 tumor cells. <i>Life Sciences</i> , 2004 , 75, 1747-59	6.8	
76	Why do we not all have proteinuria? An update of our current understanding of the glomerular barrier. <i>Physiology</i> , 2004 , 19, 7-10	9.8	47
75	Glomerular size and charge selectivity in the mouse after exposure to glucosaminoglycan-degrading enzymes. <i>Journal of the American Society of Nephrology: JASN</i> , 2003 , 14, 1756-65	12.7	105
74	Dynamic alterations of glomerular charge density in fixed rat kidneys suggest involvement of endothelial cell coat. <i>American Journal of Physiology - Renal Physiology</i> , 2003 , 285, F722-30	4.3	13
73	Synthesis of sulfated proteoglycans by bovine glomerular endothelial cells in culture. <i>American Journal of Physiology - Renal Physiology</i> , 2003 , 284, F373-80	4.3	32
72	The peritoneal membrane: a dynamic dialysis membrane in children. <i>Advances in Peritoneal Dialysis Conference on Peritoneal Dialysis</i> , 2003 , 19, 265-8		10
71	Effects of pH-neutral, bicarbonate-buffered dialysis fluid on peritoneal transport kinetics in children. <i>Kidney International</i> , 2002 , 61, 1527-36	9.9	58
70	Long-term clinical effects of a peritoneal dialysis fluid with less glucose degradation products. <i>Kidney International</i> , 2001 , 59, 348-57	9.9	199
69	Altered striatal amino acid neurotransmitter release monitored using microdialysis in R6/1 Huntington transgenic mice. <i>European Journal of Neuroscience</i> , 2001 , 13, 206-10	3.5	73
68	Effects of filtration rate on the glomerular barrier and clearance of four differently shaped molecules. <i>American Journal of Physiology - Renal Physiology</i> , 2001 , 281, F103-13	4.3	72
67	A gel-membrane model of glomerular charge and size selectivity in series. <i>American Journal of Physiology - Renal Physiology</i> , 2001 , 280, F396-405	4.3	111
66	Optimization of Peritoneal Dialysis Prescription Using Computer Models of Peritoneal Transport. <i>Peritoneal Dialysis International</i> , 2001 , 21, 148-151	2.8	53
65	A quantitative analysis of the glomerular charge barrier in the rat. <i>American Journal of Physiology - Renal Physiology</i> , 2001 , 280, F646-56	4.3	38

64	Puromycin aminonucleoside damages the glomerular size barrier with minimal effects on charge density. <i>American Journal of Physiology - Renal Physiology</i> , 2001 , 281, F503-12	4.3	24
63	Limitations in anthropometric calculations of total body water in patients on peritoneal dialysis. <i>Journal of the American Society of Nephrology: JASN</i> , 2001 , 12, 568-573	12.7	28
62	Dynamic changes of the total pore area available for peritoneal exchange in children. <i>Journal of the American Society of Nephrology: JASN</i> , 2001 , 12, 1524-1529	12.7	46
61	Unrestricted pore area (A0/Delta α) is a better indicator of peritoneal membrane function than PET. <i>Kidney International</i> , 2000 , 58, 1773-9	9.9	10
60	Dyslipidemia in Peritoneal Dialysis [Relation to Dialytic Variables. <i>Peritoneal Dialysis International</i> , 2000 , 20, 306-314	2.8	36
59	Optimal Volume Prescription for Children on Peritoneal Dialysis. <i>Peritoneal Dialysis International</i> , 2000 , 20, 603-606	2.8	40
58	Orosomuroid has a cAMP-dependent effect on human endothelial cells and inhibits the action of histamine. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2000 , 278, H1725-31	5.2	18
57	Glomerular size and charge selectivity in the rat as revealed by FITC-ficoll and albumin. <i>American Journal of Physiology - Renal Physiology</i> , 2000 , 279, F84-91	4.3	76
56	Human endothelial cells produce orosomuroid, an important component of the capillary barrier. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 1999 , 276, H530-4	5.2	43
55	Physiological and morphological effects of perfusing isolated rat kidneys with hyperosmolal mannitol solutions. <i>Acta Physiologica Scandinavica</i> , 1999 , 166, 231-8		16
54	Are fractional clearances overestimated?. <i>Kidney International</i> , 1999 , 56, 2309-10	9.9	
53	Three-pore model applied to automated peritoneal dialysis. <i>Contributions To Nephrology</i> , 1999 , 129, 35-43		3
52	Improved clearance of iohexol with longer haemodialysis despite similar Kt/V for urea. <i>Nephrology Dialysis Transplantation</i> , 1999 , 14, 2407-12	4.3	10
51	Body composition in renal failure and the effect of dialysis. <i>Applied Radiation and Isotopes</i> , 1998 , 49, 665-67		
50	Estimation of peritoneal mass transport by three-pore model in children. <i>Kidney International</i> , 1998 , 54, 1372-9	9.9	38
49	Glomerular charge selectivity for horseradish peroxidase and albumin at low and normal ionic strengths. <i>Acta Physiologica Scandinavica</i> , 1998 , 163, 83-91		46
48	Glomerular charge selectivity for proteins larger than serum albumin as revealed by lactate dehydrogenase isoforms. <i>Acta Physiologica Scandinavica</i> , 1998 , 162, 481-8		28
47	How to evaluate and optimize peritoneal dialysis treatment. <i>Nephrology Dialysis Transplantation</i> , 1998 , 13 Suppl 6, 112-6	4.3	8

46	Body composition in patients treated with peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , 1998 , 13, 1511-7	4.3	33
45	Capd in Patients with Autosomal Dominant Polycystic Kidney Disease. <i>Peritoneal Dialysis International</i> , 1998 , 18, 429-432	2.8	35
44	Creatinine generation rate and lean body mass: a critical analysis in peritoneal dialysis patients. <i>Kidney International</i> , 1997 , 51, 855-9	9.9	30
43	High glomerular permeability of bikunin despite similarity in charge and hydrodynamic size to serum albumin. <i>Kidney International</i> , 1997 , 51, 1053-8	9.9	36
42	Acute effects of C-peptide on the microvasculature of isolated perfused skeletal muscles and kidneys in rat. <i>Acta Physiologica Scandinavica</i> , 1996 , 156, 19-25		26
41	Impaired glomerular permselectivity for albumin in chemically medullectomized WKY rats. <i>Acta Physiologica Scandinavica</i> , 1996 , 156, 61-7		
40	Physiological and histological characterisation of a pig kidney in vitro perfusion model for xenotransplantation studies. <i>Scandinavian Journal of Urology and Nephrology</i> , 1996 , 30, 213-21		19
39	Assessing the peritoneal dialysis capacities of individual patients. <i>Kidney International</i> , 1995 , 47, 1187-98	9.9	125
38	Transport of macromolecules across microvascular walls: the two-pore theory. <i>Physiological Reviews</i> , 1994 , 74, 163-219	47.9	461
37	Analysis of the pressure-flow characteristics of isolated perfused rat kidneys with inhibited tubular reabsorption. <i>Acta Physiologica Scandinavica</i> , 1994 , 150, 189-99		18
36	Reduced permselectivity in isolated perfused rat kidneys following small elevations of glomerular capillary pressure. <i>Acta Physiologica Scandinavica</i> , 1994 , 150, 201-9		17
35	Addition of purified orosomuroid preserves the glomerular permeability for albumin in isolated perfused rat kidneys. <i>Acta Physiologica Scandinavica</i> , 1993 , 147, 1-8		45
34	Capillary diffusion capacity for Cr-EDTA and cyanocobalamine in spontaneously beating rat hearts. <i>Acta Physiologica Scandinavica</i> , 1993 , 147, 37-47		5
33	Changes in myocardial capillary diffusion capacity during infusion of vasoactive drugs. <i>Acta Physiologica Scandinavica</i> , 1993 , 147, 49-58		2
32	Intraperitoneal fluid fluxes analyzed according to the Three-pore model in individual patients on CAPD. <i>Blood Purification</i> , 1992 , 10, 203-8	3.1	1
31	Glomerular permselectivity is dependent on adequate serum concentrations of orosomuroid. <i>Kidney International</i> , 1992 , 41, 310-6	9.9	58
30	An isolated perfused rat kidney preparation designed for assessment of glomerular permeability characteristics. <i>Acta Physiologica Scandinavica</i> , 1992 , 144, 65-73		25
29	Computer simulations of peritoneal fluid transport in CAPD. <i>Kidney International</i> , 1991 , 40, 315-25	9.9	237

28	A note on the errors of using venous congestion in intact rats for determinations of microvascular permeability. <i>Acta Physiologica Scandinavica</i> , 1991 , 143, 233-8	4
27	Upper and lower bounds on capillary permeability ratios of Cr-EDTA to cyanocobalamin in rat hindquarters. <i>Acta Physiologica Scandinavica</i> , 1991 , 143, 239-41	1
26	Understanding the Kinetics of Peritoneal Transport 1991 , 1563-1572	8
25	Morphology of the isolated rat hindquarter preparation: tissue preservation, perfusion heterogeneity and a note on the effect of fixative osmolality. <i>Acta Physiologica Scandinavica</i> , 1988 , 132, 391-400	2
24	Transcapillary clearance of albumin in rat skeletal muscle monitored by external detection. Effects of alterations in capillary surface area. <i>Acta Physiologica Scandinavica</i> , 1988 , 132, 495-504	9
23	Diffusional transport of albumin from interstitium to blood across small pores in the capillary walls of rat skeletal muscle. <i>Acta Physiologica Scandinavica</i> , 1988 , 133, 63-71	17
22	On the steady-state relationship between the microvascular hydrostatic pressure and the transvascular filtration rate. Effects of heteroporosity. <i>Acta Physiologica Scandinavica</i> , 1987 , 129, 441-2	3
21	Orosomuroid as one of the serum components contributing to normal capillary permselectivity in rat skeletal muscle. <i>Acta Physiologica Scandinavica</i> , 1987 , 129, 127-35	102
20	Influence of perfusate oncotic pressure on the transcapillary clearance of albumin in maximally vasodilated rat skeletal muscle. <i>Acta Physiologica Scandinavica</i> , 1987 , 130, 219-28	5
19	Blood-to-tissue transport of albumin in rat fibrosarcomas at two different implantation sites. <i>Acta Physiologica Scandinavica</i> , 1987 , 131, 93-101	4
18	Fluid and protein fluxes across small and large pores in the microvasculature. Application of two-pore equations. <i>Acta Physiologica Scandinavica</i> , 1987 , 131, 411-28	130
17	A technique for assessing capillary permeability from transvascular protein flux data obtained at low filtration rates. <i>Acta Physiologica Scandinavica</i> , 1986 , 127, 263-5	9
16	Capillary permeability in rat hindquarters as determined by estimations of capillary reflection coefficients. <i>Acta Physiologica Scandinavica</i> , 1986 , 127, 289-303	32
15	Restricted diffusion of CrEDTA and cyanocobalamine across the exchange vessels in rat hindquarters. <i>Acta Physiologica Scandinavica</i> , 1986 , 127, 359-72	25
14	Calcium dependence of histamine-induced increases in capillary permeability in isolated perfused rat hindquarters. <i>Acta Physiologica Scandinavica</i> , 1986 , 128, 247-58	40
13	On-line computer evaluation of dye indicator dilution curves for determinations of capillary diffusion capacities. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 1986 , 46, 325-33	2 7
12	Serum factors other than albumin are needed for the maintenance of normal capillary permselectivity in rat hindlimb muscle. <i>Acta Physiologica Scandinavica</i> , 1985 , 123, 427-36	41
11	Changes in transcapillary exchange induced by perfusion fixation with glutaraldehyde, followed by measurements of capillary filtration coefficient, diffusion capacity and albumin clearance. <i>Acta Physiologica Scandinavica</i> , 1985 , 124, 99-106	29

10	Evaluation of the stretched pore phenomenon in isolated rat hindquarters. <i>Acta Physiologica Scandinavica</i> , 1985 , 125, 453-9		13
9	Effects of noradrenaline on the transcapillary passage of albumin, fluid and CrEDTA in the perfused rat hindlimb. <i>Acta Physiologica Scandinavica</i> , 1985 , 125, 561-71		14
8	Solvent drag component of unidirectional albumin out-flux. <i>Microvascular Research</i> , 1985 , 30, 246-8	3.7	5
7	Higher albumin clearance in rat hindquarters perfused with pure albumin solution than with serum as perfusate. <i>Acta Physiologica Scandinavica</i> , 1984 , 122, 93-5		8
6	Transcapillary passage of albumin in mammary tumours and in normal lactating mammary glands of the rat. <i>Acta Physiologica Scandinavica</i> , 1984 , 122, 497-505		23
5	Importance of molecular charge for the passage of endogenous macromolecules across continuous capillary walls, studied by serum clearance of lactate dehydrogenase (LDH) isoenzymes. <i>Acta Physiologica Scandinavica</i> , 1983 , 117, 123-30		44
4	Permeability of fenestrated capillaries in the isolated pig pancreas, with effects of bradykinin and histamine, as studied by simultaneous registration of filtration and diffusion capacities. <i>Acta Physiologica Scandinavica</i> , 1982 , 114, 67-74		28
3	Capillary permeability of sulphate-substituted and neutral dextran fractions in the rat hindquarter vascular bed. <i>Acta Physiologica Scandinavica</i> , 1982 , 115, 397-404		29
2	Extent of structurally reduced venous distensibility in rats. <i>Clinical Science</i> , 1981 , 61 Suppl 7, 125s-128s		5
1	Structurally reduced distensibility of cardiovascular low-pressure compartments in primary hypertension, as studied in spontaneously hypertensive rats (SHR). <i>Acta Physiologica Scandinavica</i> , 1981 , 112, 473-80		15