

SÃ©bastien Andre

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6005151/publications.pdf>

Version: 2024-02-01

21
papers

1,219
citations

471509

17
h-index

677142

22
g-index

22
all docs

22
docs citations

22
times ranked

2082
citing authors

#	ARTICLE	IF	CITATIONS
1	Impairment of gut microbial biotin metabolism and host biotin status in severe obesity: effect of biotin and prebiotic supplementation on improved metabolism. <i>Gut</i> , 2022, 71, 2463-2480.	12.1	53
2	Senescence-associated β -galactosidase in subcutaneous adipose tissue associates with altered glycaemic status and truncal fat in severe obesity. <i>Diabetologia</i> , 2021, 64, 240-254.	6.3	45
3	Imidazole propionate is increased in diabetes and associated with dietary patterns and altered microbial ecology. <i>Nature Communications</i> , 2020, 11, 5881.	12.8	122
4	AhR activation defends gut barrier integrity against damage occurring in obesity. <i>Molecular Metabolism</i> , 2020, 39, 101007.	6.5	42
5	Synergistic convergence of microbiota-specific systemic IgG and secretory IgA. <i>Journal of Allergy and Clinical Immunology</i> , 2019, 143, 1575-1585.e4.	2.9	86
6	Mucosal-associated invariant T (MAIT) cells are depleted and prone to apoptosis in cardiometabolic disorders. <i>FASEB Journal</i> , 2018, 32, 5078-5089.	0.5	37
7	Immune cell-derived cytokines contribute to obesity-related inflammation, fibrogenesis and metabolic deregulation in human adipose tissue. <i>Scientific Reports</i> , 2017, 7, 3000.	3.3	106
8	T Cell Populations and Functions Are Altered in Human Obesity and Type 2 Diabetes. <i>Current Diabetes Reports</i> , 2017, 17, 81.	4.2	71
9	Adipose tissue adaptive response to <i>trans-10, cis-12</i> -conjugated linoleic acid engages alternatively activated M2 macrophages. <i>FASEB Journal</i> , 2016, 30, 241-251.	0.5	12
10	Jejunal T Cell Inflammation in Human Obesity Correlates with Decreased Enterocyte Insulin Signaling. <i>Cell Metabolism</i> , 2015, 22, 113-124.	16.2	130
11	Cryptic polyreactivity of IgG expressed by splenic marginal zone B-cell lymphoma. <i>Molecular Immunology</i> , 2014, 60, 54-61.	2.2	9
12	T Cell-Derived IL-22 Amplifies IL-1-Driven Inflammation in Human Adipose Tissue: Relevance to Obesity and Type 2 Diabetes. <i>Diabetes</i> , 2014, 63, 1966-1977.	0.6	197
13	Proteolytic antibodies activate factor IX in patients with acquired hemophilia. <i>Blood</i> , 2011, 117, 2257-2264.	1.4	38
14	Discontinuous epitopes on the C2 domain of coagulation Factor VIII mapped by computer-designed synthetic peptides. <i>British Journal of Haematology</i> , 2011, 155, 487-497.	2.5	16
15	A Cellular Viewpoint of Anti-FVIII Immune Response in Hemophilia A. <i>Clinical Reviews in Allergy and Immunology</i> , 2009, 37, 105-113.	6.5	24
16	Kinetics and thermodynamics of interaction of coagulation factor VIII with a pathogenic human antibody. <i>Molecular Immunology</i> , 2009, 47, 290-297.	2.2	6
17	Functional variability of antibodies upon oxidative processes. <i>Autoimmunity Reviews</i> , 2008, 7, 574-578.	5.8	18
18	Factor VIII bypasses CD91/LRP for endocytosis by dendritic cells leading to T-cell activation. <i>Haematologica</i> , 2008, 93, 83-89.	3.5	34

#	ARTICLE	IF	CITATIONS
19	Auditing Protein Therapeutics Management by Professional APCs: Toward Prevention of Immune Responses against Therapeutic Proteins. <i>Journal of Immunology</i> , 2008, 181, 1609-1615.	0.8	18
20	Comparison of the immunogenicity of different therapeutic preparations of human factor VIII in the murine model of hemophilia A. <i>Haematologica</i> , 2007, 92, 1423-1426.	3.5	40
21	A role for exposed mannosylations in presentation of human therapeutic self-proteins to CD4+ T lymphocytes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 8965-8970.	7.1	110