

# Eli Song

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6006803/publications.pdf>

Version: 2024-02-01

18  
papers

403  
citations

1040056

9  
h-index

888059

17  
g-index

20  
all docs

20  
docs citations

20  
times ranked

718  
citing authors

#	ARTICLE	IF	CITATIONS
1	Islet organoid as a promising model for diabetes. <i>Protein and Cell</i> , 2022, 13, 239-257.	11.0	27
2	Metaxins are core components of mitochondrial transport adaptor complexes. <i>Nature Communications</i> , 2021, 12, 83.	12.8	48
3	Rab7-harboring vesicles are carriers of the transferrin receptor through the biosynthetic secretory pathway. <i>Science Advances</i> , 2021, 7, .	10.3	11
4	WFS1 functions in ER export of vesicular cargo proteins in pancreatic Î²-cells. <i>Nature Communications</i> , 2021, 12, 6996.	12.8	24
5	ALKBH1 deficiency leads to loss of homeostasis in human diploid somatic cells. <i>Protein and Cell</i> , 2020, 11, 688-695.	11.0	8
6	Dynamics of Auxilin 1 and GAK in clathrin-mediated traffic. <i>Journal of Cell Biology</i> , 2020, 219, .	5.2	37
7	Extracellular RNA: mechanisms of itâ€™s transporting into target cells. <i>ExRNA</i> , 2019, 1, .	1.0	8
8	Melatonin therapy protects against renal injury before and after release of bilateral ureteral obstruction in rats. <i>Life Sciences</i> , 2019, 229, 104-115.	4.3	6
9	I-BAR protein IRSp53 regulates clathrin-independent endocytosis in a biphasic manner. <i>Science Bulletin</i> , 2018, 63, 149-151.	9.0	0
10	hTAC internalizes via both clathrin-dependent and clathrin-independent endocytosis in mammalian cells. <i>Protein and Cell</i> , 2018, 9, 896-901.	11.0	4
11	AIM interneurons mediate feeding suppression through the TYRA-2 receptor in <i>C. elegans</i> . <i>Biophysics Reports</i> , 2018, 4, 17-24.	0.8	4
12	Semaphorin-3A and Netrin-1 predict the development of kidney injury in children with congenital hydronephrosis. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018, 78, 55-61.	1.2	6
13	Proteomic analysis of insulin secretory granules in INS-1 cells by protein correlation profiling. <i>Biophysics Reports</i> , 2018, 4, 329-338.	0.8	11
14	Dynamics of phosphoinositide conversion in clathrin-mediated endocytic traffic. <i>Nature</i> , 2017, 552, 410-414.	27.8	119
15	HID-1 is required for homotypic fusion of immature secretory granules during maturation. <i>ELife</i> , 2016, 5, .	6.0	35
16	Targeted disruption of Rab10 causes early embryonic lethality. <i>Protein and Cell</i> , 2015, 6, 463-467.	11.0	32
17	An efficient two-step subcellular fractionation method for the enrichment of insulin granules from INS-1 cells. <i>Biophysics Reports</i> , 2015, 1, 34-40.	0.8	5
18	HID-1 is a peripheral membrane protein primarily associated with the medial- and trans- Golgi apparatus. <i>Protein and Cell</i> , 2011, 2, 74-85.	11.0	16