

Lutz Fehrmann

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6013332/publications.pdf>

Version: 2024-02-01

27
papers

710
citations

687363

13
h-index

580821

25
g-index

27
all docs

27
docs citations

27
times ranked

1297
citing authors

#	ARTICLE	IF	CITATIONS
1	The Horizontal Distribution of Branch Biomass in European Beech: A Model Based on Measurements and TLS Based Proxies. <i>Remote Sensing</i> , 2021, 13, 1041.	4.0	4
2	Flooding and land use change in Jambi Province, Sumatra: integrating local knowledge and scientific inquiry. <i>Ecology and Society</i> , 2020, 25, .	2.3	19
3	Trade-offs between multifunctionality and profit in tropical smallholder landscapes. <i>Nature Communications</i> , 2020, 11, 1186.	12.8	156
4	Comparison of estimators of variance for forest inventories with systematic sampling - results from artificial populations. <i>Forest Ecosystems</i> , 2020, 7, .	3.1	15
5	Improving precision of field inventory estimation of aboveground biomass through an alternative view on plot biomass. <i>Forest Ecosystems</i> , 2020, 7, .	3.1	9
6	In search of a variance estimator for systematic sampling. <i>Scandinavian Journal of Forest Research</i> , 2019, 34, 300-312.	1.4	4
7	A unified framework for land cover monitoring based on a discrete global sampling grid (GSG). <i>Environmental Monitoring and Assessment</i> , 2019, 191, 46.	2.7	5
8	A generalized algebraic difference approach allows an improved estimation of aboveground biomass dynamics of <i>Cunninghamia lanceolata</i> and <i>Castanopsis sclerophylla</i> forests. <i>Annals of Forest Science</i> , 2017, 74, 1.	2.0	9
9	Development of a Compatible Taper Function and Stand-Level Merchantable Volume Model for Chinese Fir Plantations. <i>PLoS ONE</i> , 2016, 11, e0147610.	2.5	29
10	Two neighborhood-free plot designs for adaptive sampling of forests. <i>Environmental and Ecological Statistics</i> , 2016, 23, 279-299.	3.5	4
11	Development of stand density management diagrams for Chinese fir plantations. <i>Forestry</i> , 2016, 89, 36-45.	2.3	16
12	Estimation of stand-level aboveground biomass dynamics using tree ring analysis in a Chinese fir plantation in Shitai County, Anhui Province, China. <i>New Forests</i> , 2016, 47, 319-332.	1.7	18
13	ABOVEGROUND AND BELOWGROUND BIOMASS AND CARBON ESTIMATES FOR CLONAL EUCALYPTUS TREES IN SOUTHEAST BRAZIL. <i>Revista Arvore</i> , 2015, 39, 353-363.	0.5	22
14	On the site-level suitability of biomass models. <i>Environmental Modelling and Software</i> , 2015, 73, 14-26.	4.5	14
15	Local Parameter Estimation of Topographic Normalization for Forest Type Classification. <i>IEEE Geoscience and Remote Sensing Letters</i> , 2015, 12, 1998-2002.	3.1	4
16	Moving window-based topographic normalization of optical satellite imagery for forest mapping in mountainous terrain. , 2014, , .		0
17	Sampling for landscape elementsâ€”a case study from Lower Saxony, Germany. <i>Environmental Monitoring and Assessment</i> , 2014, 186, 1421-1430.	2.7	5
18	Sampling for landscape elementsâ€”a case study from Lower Saxony, Germany. , 2014, 186, 1421.		1

#	ARTICLE	IF	CITATIONS
19	Using terrestrial laser scanning to support biomass estimation in densely stocked young tree plantations. <i>International Journal of Remote Sensing</i> , 2013, 34, 8699-8709.	2.9	11
20	The relationship between tree species richness, canopy space exploration and productivity in a temperate broad-leaf mixed forest. <i>Forest Ecology and Management</i> , 2013, 310, 366-374.	3.2	70
21	The potential of terrestrial laser scanning for the estimation of understory biomass in coppice-with-standard systems. <i>Biomass and Bioenergy</i> , 2012, 47, 20-25.	5.7	12
22	Triangulation based inclusion probabilities: a design-unbiased sampling approach. <i>Environmental and Ecological Statistics</i> , 2012, 19, 107-123.	3.5	14
23	Above- and belowground biomass in a Brazilian Cerrado. <i>Forest Ecology and Management</i> , 2011, 262, 491-499.	3.2	86
24	A national level forest resource assessment for Burkina Faso – A field based forest inventory in a semiarid environment combining small sample size with large observation plots. <i>Forest Ecology and Management</i> , 2011, 262, 1532-1540.	3.2	29
25	A new design for sampling with adaptive sample plots. <i>Environmental and Ecological Statistics</i> , 2011, 18, 223-237.	3.5	14
26	Comparison of linear and mixed-effect regression models and a <i>k</i> -nearest neighbour approach for estimation of single-tree biomass. <i>Canadian Journal of Forest Research</i> , 2008, 38, 1-9.	1.7	57
27	General considerations about the use of allometric equations for biomass estimation on the example of Norway spruce in central Europe. <i>Forest Ecology and Management</i> , 2006, 236, 412-421.	3.2	83