

# Heeyoung Chae

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6014248/publications.pdf>

Version: 2024-02-01

7  
papers

134  
citations

1683354

5  
h-index

1719596

7  
g-index

7  
all docs

7  
docs citations

7  
times ranked

266  
citing authors

#	ARTICLE	IF	CITATIONS
1	NNT reverse mode of operation mediates glucose control of mitochondrial NADPH and glutathione redox state in mouse pancreatic $\hat{I}^2$ -cells. <i>Molecular Metabolism</i> , 2017, 6, 535-547.	3.0	35
2	Somatostatin Is Only Partly Required for the Glucagonostatic Effect of Glucose but Is Necessary for the Glucagonostatic Effect of KATP Channel Blockers. <i>Diabetes</i> , 2018, 67, 2239-2253.	0.3	33
3	SGLT2 is not expressed in pancreatic $\hat{I}^{\pm}$ - and $\hat{I}^2$ -cells, and its inhibition does not directly affect glucagon and insulin secretion in rodents and humans. <i>Molecular Metabolism</i> , 2020, 42, 101071.	3.0	26
4	Metallothionein 1 negatively regulates glucose-stimulated insulin secretion and is differentially expressed in conditions of beta cell compensation and failure in mice and humans. <i>Diabetologia</i> , 2019, 62, 2273-2286.	2.9	16
5	KATP channel blockers control glucagon secretion by distinct mechanisms: A direct stimulation of $\hat{I}^{\pm}$ -cells involving a $[Ca^{2+}]_c$ rise and an indirect inhibition mediated by somatostatin. <i>Molecular Metabolism</i> , 2021, 53, 101268.	3.0	13
6	The lack of functional nicotinamide nucleotide transhydrogenase only moderately contributes to the impairment of glucose tolerance and glucose-stimulated insulin secretion in C57BL/6J vs C57BL/6N mice. <i>Diabetologia</i> , 2021, 64, 2550-2561.	2.9	7
7	Can Tea Extracts Exert a Protective Effect Against Diabetes by Reducing Oxidative Stress and Decreasing Glucotoxicity in Pancreatic $\hat{I}^2$ -Cells?. <i>Diabetes and Metabolism Journal</i> , 2015, 39, 27.	1.8	4