

# Fumikazu Ikemori

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6017464/publications.pdf>

Version: 2024-02-01

36  
papers

1,110  
citations

430874

18  
h-index

395702

33  
g-index

36  
all docs

36  
docs citations

36  
times ranked

1369  
citing authors

#	ARTICLE	IF	CITATIONS
1	Formation of secondary organic aerosol tracers from anthropogenic and biogenic volatile organic compounds under varied NO and oxidant conditions. <i>Atmospheric Environment: X</i> , 2022, , 100169.	1.4	0
2	Time-resolved characterization of organic compounds in PM <sub>2.5</sub> collected at Oki Island, Japan, affected by transboundary pollution of biomass and non-biomass burning from Northeast China. <i>Science of the Total Environment</i> , 2021, 750, 142183.	8.0	16
3	Source apportionment in PM <sub>2.5</sub> in central Japan using positive matrix factorization focusing on small-scale local biomass burning. <i>Atmospheric Pollution Research</i> , 2021, 12, 162-172.	3.8	19
4	Organic Molecular Tracers in PM <sub>2.5</sub> at Urban Sites during Spring and Summer in Japan: Impact of Secondary Organic Aerosols on Water-Soluble Organic Carbon. <i>Atmosphere</i> , 2021, 12, 579.	2.3	9
5	Trends in PM <sub>2.5</sub> Concentration in Nagoya, Japan, from 2003 to 2018 and Impacts of PM <sub>2.5</sub> Countermeasures. <i>Atmosphere</i> , 2021, 12, 590.	2.3	13
6	Site-specific variation in mass concentration and chemical components in ambient nanoparticles (PM <sub>0.1</sub> ) in North Sumatra Province-Indonesia. <i>Atmospheric Pollution Research</i> , 2021, 12, 101062.	3.8	18
7	Size-Segregated Particulate Matter Down to PM <sub>0.1</sub> and Carbon Content during the Rainy and Dry Seasons in Sumatra Island, Indonesia. <i>Atmosphere</i> , 2021, 12, 1441.	2.3	12
8	Four- and Five-Carbon Dicarboxylic Acids Present in Secondary Organic Aerosol Produced from Anthropogenic and Biogenic Volatile Organic Compounds. <i>Atmosphere</i> , 2021, 12, 1703.	2.3	9
9	Health effects of PM <sub>2.5</sub> sources on children's allergic and respiratory symptoms in Fukuoka, Japan. <i>Science of the Total Environment</i> , 2020, 709, 136023.	8.0	50
10	A twenty-year deposition record of elemental carbon in Northern Japan retrieved from archived filters. <i>Scientific Reports</i> , 2020, 10, 4520.	3.3	1
11	Estimating Mass Concentration Using a Low-cost Portable Particle Counter Based on Full-year Observations: Issues to Obtain Reliable Atmospheric PM <sub>2.5</sub> Data. <i>Asian Journal of Atmospheric Environment</i> , 2020, 14, 155-169.	1.1	2
12	Traffic source impacts on chlorinated polycyclic aromatic hydrocarbons in PM <sub>2.5</sub> by short-range transport. <i>Atmospheric Environment</i> , 2019, 216, 116944.	4.1	18
13	The influence of the open burning of agricultural biomass and forest fires in Thailand on the carbonaceous components in size-fractionated particles. <i>Environmental Pollution</i> , 2019, 247, 238-247.	7.5	86
14	Characterization and possible sources of nitrated mono- and di-aromatic hydrocarbons containing hydroxyl and/or carboxyl functional groups in ambient particles in Nagoya, Japan. <i>Atmospheric Environment</i> , 2019, 211, 91-102.	4.1	24
15	Evaluation of the genotoxicity of PM <sub>2.5</sub> collected by a high-volume air sampler with impactor. <i>Genes and Environment</i> , 2019, 41, 7.	2.1	11
16	Distributions and multiple sources of chlorinated polycyclic aromatic hydrocarbons in the air over Japan. <i>Science of the Total Environment</i> , 2019, 649, 364-371.	8.0	29
17	Decreasing trend of elemental carbon concentration with changes in major sources at Mega city Nagoya, Central Japan. <i>Atmospheric Environment</i> , 2019, 199, 155-163.	4.1	18
18	Impact of field biomass burning on local pollution and long-range transport of PM <sub>2.5</sub> in Northeast Asia. <i>Environmental Pollution</i> , 2019, 244, 414-422.	7.5	46

#	ARTICLE	IF	CITATIONS
19	A Cascade Air Sampler with Multi-nozzle Inertial Filters for PM <sub>0.1</sub> . Aerosol and Air Quality Research, 2019, 19, 1666-1677.	2.1	9
20	Structural and Light-Absorption Characteristics of Complex Water-Insoluble Organic Mixtures in Urban Submicrometer Aerosols. Environmental Science & Technology, 2017, 51, 8293-8303.	10.0	49
21	Weak size dependence of resuspended radiocesium adsorbed on soil particles collected after the Fukushima nuclear accident. Journal of Environmental Radioactivity, 2017, 172, 122-129.	1.7	15
22	Identification of biased sectors in emission data using a combination of chemical transport model and receptor model. Atmospheric Environment, 2017, 166, 166-181.	4.1	21
23	Source apportionment of chlorinated polycyclic aromatic hydrocarbons associated with ambient particles in a Japanese megacity. Scientific Reports, 2016, 6, 38358.	3.3	18
24	Light Absorption and Excitation-Induced Emission Fluorescence of Urban Organic Aerosol Components and Their Relationship to Chemical Structure. Environmental Science & Technology, 2016, 50, 10859-10868.	10.0	138
25	Associations Between Fine Particulate Matter Components and Daily Mortality in Nagoya, Japan. Journal of Epidemiology, 2016, 26, 249-257.	2.4	41
26	Comparison of Air Pollution in Metropolises in China (Beijing) and Japan (Osaka and Nagoya) on the Basis of the Levels of Contaminants and Mutagenicity. Biological and Pharmaceutical Bulletin, 2016, 39, 415-422.	1.4	6
27	Spatial correlativity of atmospheric particulate components simultaneously collected in Japan. Environmental Monitoring and Assessment, 2016, 188, 85.	2.7	0
28	Local and seasonal variations in concentrations of chlorinated polycyclic aromatic hydrocarbons associated with particles in a Japanese megacity. Journal of Hazardous Materials, 2016, 312, 254-261.	12.4	24
29	Chemical Structural Characteristics of HULIS and Other Fractionated Organic Matter in Urban Aerosols: Results from Mass Spectral and FT-IR Analysis. Environmental Science & Technology, 2016, 50, 1721-1730.	10.0	92
30	Development of an Analytical Method for Strong Mutagens/Carcinogens, 3,9-Dinitrofluoranthene and Dinitropyrene Isomers, in the Environment and Their Particle-Size Distribution in Airborne Particles. Chromatographia, 2015, 78, 55-63.	1.3	5
31	Optimisation of pre-treatment and ionisation for GC/MS analysis for the determination of chlorinated PAHs in atmospheric particulate samples. International Journal of Environmental Analytical Chemistry, 2015, 95, 1157-1168.	3.3	18
32	Influence of contemporary carbon originating from the 2003 Siberian forest fire on organic carbon in PM 2.5 in Nagoya, Japan. Science of the Total Environment, 2015, 530-531, 403-410.	8.0	19
33	Air Pollution with Particulate Matter and Mutagens: Relevance of Asian Dust to Mutagenicity of Airborne Particles in Japan. Genes and Environment, 2014, 36, 120-136.	2.1	8
34	Properties of light-absorbing aerosols in the Nagoya urban area, Japan, in August 2011 and January 2012: Contributions of brown carbon and lensing effect. Journal of Geophysical Research D: Atmospheres, 2014, 119, 12,721.	3.3	57
35	Sulfate Aerosol as a Potential Transport Medium of Radiocesium from the Fukushima Nuclear Accident. Environmental Science & Technology, 2012, 46, 5720-5726.	10.0	208
36	Analysis of Halogenated Polycyclic Aromatic Hydrocarbons in the Air. , 0, , .		1