

Fumikazu Ikemori

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6017464/publications.pdf>

Version: 2024-02-01

36
papers

1,110
citations

430843

18
h-index

395678

33
g-index

36
all docs

36
docs citations

36
times ranked

1369
citing authors

#	ARTICLE	IF	CITATIONS
1	Formation of secondary organic aerosol tracers from anthropogenic and biogenic volatile organic compounds under varied NO and oxidant conditions. <i>Atmospheric Environment: X</i> , 2022, , 100169.	1.4	0
2	Time-resolved characterization of organic compounds in PM _{2.5} collected at Oki Island, Japan, affected by transboundary pollution of biomass and non-biomass burning from Northeast China. <i>Science of the Total Environment</i> , 2021, 750, 142183.	8.0	16
3	Source apportionment in PM _{2.5} in central Japan using positive matrix factorization focusing on small-scale local biomass burning. <i>Atmospheric Pollution Research</i> , 2021, 12, 162-172.	3.8	19
4	Organic Molecular Tracers in PM _{2.5} at Urban Sites during Spring and Summer in Japan: Impact of Secondary Organic Aerosols on Water-Soluble Organic Carbon. <i>Atmosphere</i> , 2021, 12, 579.	2.3	9
5	Trends in PM _{2.5} Concentration in Nagoya, Japan, from 2003 to 2018 and Impacts of PM _{2.5} Countermeasures. <i>Atmosphere</i> , 2021, 12, 590.	2.3	13
6	Site-specific variation in mass concentration and chemical components in ambient nanoparticles (PM _{0.1}) in North Sumatra Province-Indonesia. <i>Atmospheric Pollution Research</i> , 2021, 12, 101062.	3.8	18
7	Size-Segregated Particulate Matter Down to PM _{0.1} and Carbon Content during the Rainy and Dry Seasons in Sumatra Island, Indonesia. <i>Atmosphere</i> , 2021, 12, 1441.	2.3	12
8	Four- and Five-Carbon Dicarboxylic Acids Present in Secondary Organic Aerosol Produced from Anthropogenic and Biogenic Volatile Organic Compounds. <i>Atmosphere</i> , 2021, 12, 1703.	2.3	9
9	Health effects of PM _{2.5} sources on children's allergic and respiratory symptoms in Fukuoka, Japan. <i>Science of the Total Environment</i> , 2020, 709, 136023.	8.0	50
10	A twenty-year deposition record of elemental carbon in Northern Japan retrieved from archived filters. <i>Scientific Reports</i> , 2020, 10, 4520.	3.3	1
11	Estimating Mass Concentration Using a Low-cost Portable Particle Counter Based on Full-year Observations: Issues to Obtain Reliable Atmospheric PM _{2.5} Data. <i>Asian Journal of Atmospheric Environment</i> , 2020, 14, 155-169.	1.1	2
12	Traffic source impacts on chlorinated polycyclic aromatic hydrocarbons in PM _{2.5} by short-range transport. <i>Atmospheric Environment</i> , 2019, 216, 116944.	4.1	18
13	The influence of the open burning of agricultural biomass and forest fires in Thailand on the carbonaceous components in size-fractionated particles. <i>Environmental Pollution</i> , 2019, 247, 238-247.	7.5	86
14	Characterization and possible sources of nitrated mono- and di-aromatic hydrocarbons containing hydroxyl and/or carboxyl functional groups in ambient particles in Nagoya, Japan. <i>Atmospheric Environment</i> , 2019, 211, 91-102.	4.1	24
15	Evaluation of the genotoxicity of PM _{2.5} collected by a high-volume air sampler with impactor. <i>Genes and Environment</i> , 2019, 41, 7.	2.1	11
16	Distributions and multiple sources of chlorinated polycyclic aromatic hydrocarbons in the air over Japan. <i>Science of the Total Environment</i> , 2019, 649, 364-371.	8.0	29
17	Decreasing trend of elemental carbon concentration with changes in major sources at Mega city Nagoya, Central Japan. <i>Atmospheric Environment</i> , 2019, 199, 155-163.	4.1	18
18	Impact of field biomass burning on local pollution and long-range transport of PM _{2.5} in Northeast Asia. <i>Environmental Pollution</i> , 2019, 244, 414-422.	7.5	46

#	ARTICLE	IF	CITATIONS
19	A Cascade Air Sampler with Multi-nozzle Inertial Filters for PM _{0.1} . <i>Aerosol and Air Quality Research</i> , 2019, 19, 1666-1677.	2.1	9
20	Structural and Light-Absorption Characteristics of Complex Water-Insoluble Organic Mixtures in Urban Submicrometer Aerosols. <i>Environmental Science & Technology</i> , 2017, 51, 8293-8303.	10.0	49
21	Weak size dependence of resuspended radiocesium adsorbed on soil particles collected after the Fukushima nuclear accident. <i>Journal of Environmental Radioactivity</i> , 2017, 172, 122-129.	1.7	15
22	Identification of biased sectors in emission data using a combination of chemical transport model and receptor model. <i>Atmospheric Environment</i> , 2017, 166, 166-181.	4.1	21
23	Source apportionment of chlorinated polycyclic aromatic hydrocarbons associated with ambient particles in a Japanese megacity. <i>Scientific Reports</i> , 2016, 6, 38358.	3.3	18
24	Light Absorption and Excitation-Induced Emission Fluorescence of Urban Organic Aerosol Components and Their Relationship to Chemical Structure. <i>Environmental Science & Technology</i> , 2016, 50, 10859-10868.	10.0	138
25	Associations Between Fine Particulate Matter Components and Daily Mortality in Nagoya, Japan. <i>Journal of Epidemiology</i> , 2016, 26, 249-257.	2.4	41
26	Comparison of Air Pollution in Metropolises in China (Beijing) and Japan (Osaka and Nagoya) on the Basis of the Levels of Contaminants and Mutagenicity. <i>Biological and Pharmaceutical Bulletin</i> , 2016, 39, 415-422.	1.4	6
27	Spatial correlativity of atmospheric particulate components simultaneously collected in Japan. <i>Environmental Monitoring and Assessment</i> , 2016, 188, 85.	2.7	0
28	Local and seasonal variations in concentrations of chlorinated polycyclic aromatic hydrocarbons associated with particles in a Japanese megacity. <i>Journal of Hazardous Materials</i> , 2016, 312, 254-261.	12.4	24
29	Chemical Structural Characteristics of HULIS and Other Fractionated Organic Matter in Urban Aerosols: Results from Mass Spectral and FT-IR Analysis. <i>Environmental Science & Technology</i> , 2016, 50, 1721-1730.	10.0	92
30	Development of an Analytical Method for Strong Mutagens/Carcinogens, 3,9-Dinitrofluoranthene and Dinitropyrene Isomers, in the Environment and Their Particle-Size Distribution in Airborne Particles. <i>Chromatographia</i> , 2015, 78, 55-63.	1.3	5
31	Optimisation of pre-treatment and ionisation for GC/MS analysis for the determination of chlorinated PAHs in atmospheric particulate samples. <i>International Journal of Environmental Analytical Chemistry</i> , 2015, 95, 1157-1168.	3.3	18
32	Influence of contemporary carbon originating from the 2003 Siberian forest fire on organic carbon in PM 2.5 in Nagoya, Japan. <i>Science of the Total Environment</i> , 2015, 530-531, 403-410.	8.0	19
33	Air Pollution with Particulate Matter and Mutagens: Relevance of Asian Dust to Mutagenicity of Airborne Particles in Japan. <i>Genes and Environment</i> , 2014, 36, 120-136.	2.1	8
34	Properties of light-absorbing aerosols in the Nagoya urban area, Japan, in August 2011 and January 2012: Contributions of brown carbon and lensing effect. <i>Journal of Geophysical Research D: Atmospheres</i> , 2014, 119, 12,721.	3.3	57
35	Sulfate Aerosol as a Potential Transport Medium of Radiocesium from the Fukushima Nuclear Accident. <i>Environmental Science & Technology</i> , 2012, 46, 5720-5726.	10.0	208
36	Analysis of Halogenated Polycyclic Aromatic Hydrocarbons in the Air. , 0, , .		1