

Giampietro L Vairo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6020620/publications.pdf>

Version: 2024-02-01

8
papers

156
citations

1937685

4
h-index

1588992

8
g-index

8
all docs

8
docs citations

8
times ranked

176
citing authors

#	ARTICLE	IF	CITATIONS
1	Neuromuscular and biomechanical landing performance subsequent to ipsilateral semitendinosus and gracilis autograft anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2008, 16, 2-14.	4.2	84
2	Knee Flexor Strength and Endurance Profiles After Ipsilateral Hamstring Tendons Anterior Cruciate Ligament Reconstruction. <i>Archives of Physical Medicine and Rehabilitation</i> , 2014, 95, 552-561.	0.9	28
3	Tibial Tubercle-Trochlear Groove Distance Is a Reliable and Accurate Indicator of Patellofemoral Instability. <i>Clinical Orthopaedics and Related Research</i> , 2019, 477, 1450-1458.	1.5	23
4	The TT-TG Distance/Trochlear Dysplasia Index Quotient Is the Most Accurate Indicator for Determining Patellofemoral Instability Risk. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2021, , .	2.7	10
5	Effects of Blood Flow Restriction Training on Clinical Outcomes for Patients With ACL Reconstruction: A Systematic Review. <i>Sports Health</i> , 2023, 15, 260-273.	2.7	5
6	Bilateral NIRS measurements of muscle mitochondrial capacity: Feasibility and repeatability. <i>Physiological Reports</i> , 2021, 9, e14826.	1.7	4
7	Standing Flexion Deficits Predict Self-Reported Outcomes in Women After Ipsilateral Hamstring Anterior Cruciate Ligament Reconstruction. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2013, 29, 1796-1803.	2.7	1
8	Association Between Self-Reported Kinesiophobia and Single-Leg Hop for Distance in Patients With ACL Reconstruction: A Systematic Review. <i>Sports Health</i> , 2022, 14, 674-680.	2.7	1