

Holly A Swartz

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6038122/holly-a-swartz-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

101
papers

4,228
citations

32
h-index

64
g-index

127
ext. papers

4,832
ext. citations

4.6
avg, IF

5.57
L-index

#	Paper	IF	Citations
101	The Relationship Between Default Mode and Dorsal Attention Networks Is Associated With Depressive Disorder Diagnosis and the Strength of Memory Representations Acquired Prior to the Resting State Scan.. <i>Frontiers in Human Neuroscience</i> , 2022 , 16, 749767	3.3	1
100	Social Rhythm Disruption is Associated with Greater Depressive Symptoms in People with Mood Disorders: Findings from a Multinational Online Survey During COVID-19.. <i>Canadian Journal of Psychiatry</i> , 2022 , 7067437221097905	4.8	
99	The effects of mood disorders and childhood trauma on fear of positive and negative evaluation.. <i>Acta Psychologica</i> , 2022 , 227, 103603	1.7	
98	Behavioral and neuroimaging evidence prodromal to major depressive disorder onset in a young adult without personal or family history of psychiatric disorder: Case Report 2022 , 100014		
97	Protocol for a machine learning algorithm predicting depressive disorders using the T1w/T2w ratio.. <i>MethodsX</i> , 2021 , 8, 101595	1.9	1
96	Neurobiological and behavioral mechanisms of circadian rhythm disruption in bipolar disorder: A critical multi-disciplinary literature review and agenda for future research from the ISBD task force on chronobiology. <i>Bipolar Disorders</i> , 2021 ,	3.8	2
95	Changing gears in the treatment of bipolar disorder through integrated use of electronic devices. <i>Bipolar Disorders</i> , 2021 , 23, 418-419	3.8	0
94	White matter abnormalities in adults with bipolar disorder type-II and unipolar depression. <i>Scientific Reports</i> , 2021 , 11, 7541	4.9	2
93	Telehealth Social Rhythm Therapy to Reduce Mood Symptoms and Suicide Risk Among Adolescents and Young Adults With Bipolar Disorder. <i>American Journal of Psychotherapy</i> , 2021 , apppsychotherapy20210011	2.3	1
92	Maintaining Daily Routines to Stabilize Mood: Theory, Data, and Potential Intervention for Circadian Consequences of COVID-19. <i>Canadian Journal of Psychiatry</i> , 2021 , 66, 9-13	4.8	12
91	Adrift in time: the subjective experience of circadian challenge during COVID-19 amongst people with mood disorders. <i>Chronobiology International</i> , 2021 , 1-11	3.6	2
90	A randomized pilot study of Rhythms And You (RAY): An internet-based program for bipolar disorder administered with and without clinical helper support in primary care. <i>Journal of Affective Disorders</i> , 2021 , 295, 183-191	6.6	1
89	Aberrant levels of cortical myelin distinguish individuals with depressive disorders from healthy controls. <i>NeuroImage: Clinical</i> , 2021 , 32, 102790	5.3	2
88	Prefrontal cortical activation during working memory task anticipation contributes to discrimination between bipolar and unipolar depression. <i>Neuropsychopharmacology</i> , 2020 , 45, 956-963	8.7	7
87	Social rhythm therapy-A potentially translatable psychosocial intervention for bipolar disorder. <i>Bipolar Disorders</i> , 2020 , 22, 121-127	3.8	12
86	Psychotherapy in the Digital Age: What We Can Learn From Interpersonal Psychotherapy. <i>American Journal of Psychotherapy</i> , 2020 , 73, 15-21	1.3	2
85	Psychosocial Interventions for Bipolar II Disorder. <i>American Journal of Psychotherapy</i> , 2019 , 72, 47-57	1.3	1

84	Changes in inflammation with treatment for bipolar II depression: Pilot trial data on differential effects of psychotherapy and medication. <i>Neurology Psychiatry and Brain Research</i> , 2019 , 33, 112-118	2.1	3
83	The role of the right prefrontal cortex in recognition of facial emotional expressions in depressed individuals: fNIRS study. <i>Journal of Affective Disorders</i> , 2019 , 258, 151-158	6.6	14
82	Evidence-Based Psychotherapies for Bipolar Disorder. <i>Focus (American Psychiatric Publishing)</i> , 2019 , 17, 238-248	1.1	7
81	The chronotherapeutic treatment of bipolar disorders: A systematic review and practice recommendations from the ISBD task force on chronotherapy and chronobiology. <i>Bipolar Disorders</i> , 2019 , 21, 741-773	3.8	75
80	Mass Shootings and Psychotherapy. <i>American Journal of Psychotherapy</i> , 2019 , 72, 58	1.3	
79	Addressing Unmet Needs in Bipolar Disorder. <i>Psychiatric Annals</i> , 2019 , 49, 377-378	0.5	0
78	Interpersonal Psychotherapy: A Scoping Review and Historical Perspective (1974-2017). <i>Harvard Review of Psychiatry</i> , 2019 , 27, 165-180	4.1	10
77	How obstetric settings can help address gaps in psychiatric care for pregnant and postpartum women with bipolar disorder. <i>Archives of Women's Mental Health</i> , 2018 , 21, 543-551	5	8
76	Access to Pharmacotherapy Amongst Women with Bipolar Disorder during Pregnancy: a Preliminary Study. <i>Psychiatric Quarterly</i> , 2018 , 89, 183-190	4.1	8
75	Moderators and mediators of a maternal depression treatment study: Impact of maternal trauma and parenting on child outcomes. <i>Comprehensive Psychiatry</i> , 2018 , 86, 123-130	7.3	7
74	Psychotherapy Alone and Combined With Medication as Treatments for Bipolar II Depression: A Randomized Controlled Trial. <i>Journal of Clinical Psychiatry</i> , 2018 , 79,	4.6	19
73	Revising , Fifth Edition, criteria for the bipolar disorders: Phase I of the AREDOC project. <i>Australian and New Zealand Journal of Psychiatry</i> , 2018 , 52, 1173-1182	2.6	12
72	History of childhood emotional abuse predicts lower resting-state high-frequency heart rate variability in depressed women. <i>Psychiatry Research</i> , 2018 , 269, 681-687	9.9	11
71	Depression impacts the physiological responsiveness of mother-daughter dyads during social interaction. <i>Depression and Anxiety</i> , 2017 , 34, 118-126	8.4	24
70	Prospective assessment of social network quality among depressed mothers treated with brief psychotherapy: The Social Network Quality (SNQ) scales. <i>Comprehensive Psychiatry</i> , 2017 , 78, 98-106	7.3	2
69	Prediction of near-term increases in suicidal ideation in recently depressed patients with bipolar II disorder using intensive longitudinal data. <i>Journal of Affective Disorders</i> , 2017 , 208, 363-368	6.6	8
68	Therapist Use of Specific and Nonspecific Strategies Across Two Affect-Focused Psychotherapies for Depression: Role of Adherence Monitoring. <i>Journal of Psychotherapy Integration</i> , 2017 , 27, 381-394	6.9	5
67	PRogram In Support of Moms (PRISM): Development and Beta Testing. <i>Psychiatric Services</i> , 2016 , 67, 824-6	3.3	11

66	An approach to revealing clinically relevant subgroups across the mood spectrum. <i>Journal of Affective Disorders</i> , 2016 , 203, 265-274	6.6	2
65	Telephone-administered psychotherapy in combination with antidepressant medication for the acute treatment of major depressive disorder. <i>Journal of Affective Disorders</i> , 2016 , 190, 6-11	6.6	11
64	Brief Psychotherapy for Maternal Depression: Impact on Mothers and Children. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2016 , 55, 495-503.e2	7.2	26
63	Readiness to Implement an Evidence-Based Psychotherapy: Perspectives of Community Mental Health Clinicians and Administrators. <i>Psychiatric Services</i> , 2015 , 66, 1109-12	3.3	5
62	Maternal response to child affect: Role of maternal depression and relationship quality. <i>Journal of Affective Disorders</i> , 2015 , 187, 106-13	6.6	9
61	Psychosocial Treatment of Bipolar Disorder: Clinician Knowledge, Common Approaches, and Barriers to Effective Treatment. <i>Psychiatric Services</i> , 2015 , 66, 1361-4	3.3	4
60	IOM REPORT ON PSYCHOSOCIAL INTERVENTIONS FOR MENTAL AND SUBSTANCE USE DISORDERS: THE INTERPERSONAL PSYCHOTHERAPY PERSPECTIVE. <i>Depression and Anxiety</i> , 2015 , 32, 793-5	8.4	2
59	Implementing a Web-Based Intervention to Train Community Clinicians in an Evidence-Based Psychotherapy: A Pilot Study. <i>Psychiatric Services</i> , 2015 , 66, 988-91	3.3	19
58	Are adults with bipolar disorder active? Objectively measured physical activity and sedentary behavior using accelerometry. <i>Journal of Affective Disorders</i> , 2014 , 152-154, 498-504	6.6	76
57	Treating co-occurring depression and anxiety: modeling the dynamics of psychopathology and psychotherapy using the time-varying effect model. <i>Journal of Consulting and Clinical Psychology</i> , 2014 , 82, 839-53	6.5	17
56	Brief Interpersonal Psychotherapy (IPT-B): Overview and Review of Evidence. <i>American Journal of Psychotherapy</i> , 2014 , 68, 443-62	1.3	15
55	Psychotherapy for Bipolar Disorder in Adults: A Review of the Evidence. <i>Focus (American Psychiatric Publishing)</i> , 2014 , 12, 251-266	1.1	29
54	Caregiver depression screening in a specialty mental health clinic. <i>Psychiatric Services</i> , 2014 , 65, 400	3.3	4
53	A critical appraisal of neuroimaging studies of bipolar disorder: toward a new conceptualization of underlying neural circuitry and a road map for future research. <i>American Journal of Psychiatry</i> , 2014 , 171, 829-43	11.9	373
52	Role of maternal childhood trauma on parenting among depressed mothers of psychiatrically ill children. <i>Depression and Anxiety</i> , 2013 , 30, 792-9	8.4	19
51	Exposure to maternal pre- and postnatal depression and anxiety symptoms: risk for major depression, anxiety disorders, and conduct disorder in adolescent offspring. <i>Development and Psychopathology</i> , 2013 , 25, 1045-63	4.3	49
50	Facilitators and barriers associated with implementation of evidence-based psychotherapy in community settings. <i>Psychiatric Services</i> , 2013 , 64, 1263-6	3.3	19
49	A randomized pilot study of psychotherapy and quetiapine for the acute treatment of bipolar II depression. <i>Bipolar Disorders</i> , 2012 , 14, 211-6	3.8	37

48	Impact of childhood trauma on the outcomes of a perinatal depression trial. <i>Depression and Anxiety</i> , 2012 , 29, 563-73	8.4	36
47	Psychotherapy for Bipolar II Disorder: The Role of Interpersonal and Social Rhythm Therapy. <i>Professional Psychology: Research and Practice</i> , 2012 , 43, 145-153	2.2	22
46	Cardiovascular disease and bipolar disorder: risk and clinical implications. <i>Journal of Clinical Psychiatry</i> , 2012 , 73, 1563-5	4.6	38
45	Pharmacotherapy for the treatment of acute bipolar II depression: current evidence. <i>Journal of Clinical Psychiatry</i> , 2011 , 72, 356-66	4.6	48
44	Implementing interpersonal and social rhythm therapy for mood disorders across a continuum of care. <i>Psychiatric Services</i> , 2011 , 62, 1377-80	3.3	20
43	Cardiac vagal control in nonmedicated depressed women and nondepressed controls: impact of depression status, lifetime trauma history, and respiratory factors. <i>Psychosomatic Medicine</i> , 2011 , 73, 336-43	3.7	31
42	Thinking about a close relationship differentially impacts cardiovascular stress responses among depressed and nondepressed women. <i>Health Psychology</i> , 2011 , 30, 276-84	5	4
41	Suicide attempts in bipolar I and bipolar II disorder: a review and meta-analysis of the evidence. <i>Bipolar Disorders</i> , 2010 , 12, 1-9	3.8	203
40	Interpersonal and Social Rhythm Therapy for Bipolar Disorder 2010 , 430-442		
39	Comparative outcomes among the problem areas of interpersonal psychotherapy for depression. <i>Depression and Anxiety</i> , 2010 , 27, 434-40	8.4	7
38	A Randomized Controlled Trial of Culturally Relevant, Brief Interpersonal Psychotherapy for Perinatal Depression. <i>Psychiatric Services</i> , 2009 , 60, 313-321	3.3	180
37	Emotional and cardiovascular reactivity to a child-focused interpersonal stressor among depressed mothers of psychiatrically ill children. <i>Depression and Anxiety</i> , 2009 , 26, 110-6	8.4	9
36	Psychotherapy as monotherapy for the treatment of bipolar II depression: a proof of concept study. <i>Bipolar Disorders</i> , 2009 , 11, 89-94	3.8	35
35	Bipolar Disorder Center for Pennsylvanians: Implementing an Effectiveness Trial to Improve Treatment for At-Risk Patients. <i>Psychiatric Services</i> , 2009 , 60, 888-897	3.3	4
34	A randomized controlled trial of culturally relevant, brief interpersonal psychotherapy for perinatal depression. <i>Psychiatric Services</i> , 2009 , 60, 313-21	3.3	98
33	Bipolar disorder center for Pennsylvanians: implementing an effectiveness trial to improve treatment for at-risk patients. <i>Psychiatric Services</i> , 2009 , 60, 888-97	3.3	6
32	Treatment-emergent mania/hypomania in unipolar patients. <i>Bipolar Disorders</i> , 2008 , 10, 726-32	3.8	21
31	The role of interpersonal and social rhythm therapy in improving occupational functioning in patients with bipolar I disorder. <i>American Journal of Psychiatry</i> , 2008 , 165, 1559-65	11.9	118

30	Brief interpersonal psychotherapy for depressed mothers whose children are receiving psychiatric treatment. <i>American Journal of Psychiatry</i> , 2008 , 165, 1155-62	11.9	123
29	Enhancing Interpersonal Psychotherapy for Mothers and Expectant Mothers on Low Incomes: Adaptations and Additions. <i>Journal of Contemporary Psychotherapy</i> , 2008 , 38, 23-33	2.3	39
28	The role of psychosocial interventions in bipolar II disorder: what do we know?. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, e15	4.6	15
27	The use of antidepressants in bipolar disorder. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 1307-18	4.6	44
26	Engaging women who are depressed and economically disadvantaged in mental health treatment. <i>Social Work</i> , 2007 , 52, 295-308	1.4	112
25	Randomized trial of weekly, twice-monthly, and monthly interpersonal psychotherapy as maintenance treatment for women with recurrent depression. <i>American Journal of Psychiatry</i> , 2007 , 164, 761-7	11.9	68
24	Engaging depressed patients in psychotherapy: Integrating techniques from motivational interviewing and ethnographic interviewing to improve treatment participation.. <i>Professional Psychology: Research and Practice</i> , 2007 , 38, 430-439	2.2	34
23	Interpersonal and social rhythm therapy: an intervention addressing rhythm dysregulation in bipolar disorder. <i>Dialogues in Clinical Neuroscience</i> , 2007 , 9, 325-32	5.7	59
22	An open-label trial of enhanced brief interpersonal psychotherapy in depressed mothers whose children are receiving psychiatric treatment. <i>Depression and Anxiety</i> , 2006 , 23, 398-404	8.4	30
21	Hypothyroidism in patients with bipolar I disorder treated primarily with lithium. <i>Epidemiologia E Psichiatria Sociale</i> , 2006 , 15, 123-7		35
20	Impact of maternal mental health status on child mental health treatment outcome. <i>Community Mental Health Journal</i> , 2006 , 42, 1-12	2.1	23
19	Acute treatment outcomes in patients with bipolar I disorder and co-morbid borderline personality disorder receiving medication and psychotherapy. <i>Bipolar Disorders</i> , 2005 , 7, 192-7	3.8	32
18	Interpersonal psychotherapy for depression with panic spectrum symptoms: a pilot study. <i>Depression and Anxiety</i> , 2005 , 21, 140-2	8.4	14
17	Recognition and treatment of depression. <i>AMA Journal of Ethics</i> , 2005 , 7, 430	1.4	1
16	Two-year outcomes for interpersonal and social rhythm therapy in individuals with bipolar I disorder. <i>Archives of General Psychiatry</i> , 2005 , 62, 996-1004		517
15	Depression and anxiety among mothers who bring their children to a pediatric mental health clinic. <i>Psychiatric Services</i> , 2005 , 56, 1077-83	3.3	53
14	A pilot study of brief interpersonal psychotherapy for depression among women. <i>Psychiatric Services</i> , 2004 , 55, 448-50	3.3	38
13	Culturally Relevant Psychotherapy for Perinatal Depression in Low-Income Ob/Gyn Patients. <i>Clinical Social Work Journal</i> , 2004 , 32, 327-347	1.7	17

12	Feasibility of Providing Culturally Relevant, Brief Interpersonal Psychotherapy for Antenatal Depression in an Obstetrics Clinic: A Pilot Study. <i>Research on Social Work Practice</i> , 2004 , 14, 397-407	1.4	44
11	Personality pathology and outcome in recurrently depressed women over 2 years of maintenance interpersonal psychotherapy. <i>Psychological Medicine</i> , 2004 , 34, 659-69	6.9	47
10	Managing the global burden of depression: lessons from the developing world. <i>World Psychiatry</i> , 2003 , 2, 162-3	14.4	9
9	Suicide attempts in patients with bipolar I disorder during acute and maintenance phases of intensive treatment with pharmacotherapy and adjunctive psychotherapy. <i>American Journal of Psychiatry</i> , 2002 , 159, 1160-4	11.9	75
8	A pilot study of community mental health care for depression in a supermarket setting. <i>Psychiatric Services</i> , 2002 , 53, 1132-7	3.3	16
7	Prevalence of obesity and weight change during treatment in patients with bipolar I disorder. <i>Journal of Clinical Psychiatry</i> , 2002 , 63, 528-33	4.6	138
6	Psychotherapy for bipolar depression: a phase-specific treatment strategy?. <i>Bipolar Disorders</i> , 2001 , 3, 11-22	3.8	41
5	Diagnosis of nonpsychotic patients in community clinics. <i>American Journal of Psychiatry</i> , 2000 , 157, 581-7	11.9	238
4	Interpersonal and social rhythm therapy: managing the chaos of bipolar disorder. <i>Biological Psychiatry</i> , 2000 , 48, 593-604	7.9	370
3	Adjunctive psychotherapy for bipolar disorder: Effects of changing treatment modality.. <i>Journal of Abnormal Psychology</i> , 1999 , 108, 579-587	7	93
2	Psychosocial characteristics of pregnant and nonpregnant HIV-seropositive women. <i>Psychiatric Services</i> , 1998 , 49, 1612-4	3.3	12
1	Interpersonal and Social Rhythm Therapy 275-293		