

# Miriam Ojeda Ojeda

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6041978/publications.pdf>

Version: 2024-02-01

11  
papers

690  
citations

1162889

8  
h-index

1281743

11  
g-index

11  
all docs

11  
docs citations

11  
times ranked

1122  
citing authors

#	ARTICLE	IF	CITATIONS
1	Circulating markers of oxidative stress and polycystic ovary syndrome (PCOS): a systematic review and meta-analysis. <i>Human Reproduction Update</i> , 2013, 19, 268-288.	5.2	399
2	Mediators of Low-Grade Chronic Inflammation in Polycystic Ovary Syndrome (PCOS). <i>Current Pharmaceutical Design</i> , 2013, 19, 5775-5791.	0.9	69
3	Enhancement of the inhibitory effect of an IL-15 antagonist peptide by alanine scanning. <i>Journal of Peptide Science</i> , 2012, 18, 25-29.	0.8	9
4	Imbalanced Low-Grade Inflammation and Endothelial Activation in Patients with Type 2 Diabetes Mellitus and Erectile Dysfunction. <i>Journal of Sexual Medicine</i> , 2011, 8, 2017-2030.	0.3	61
5	Temporal trends of circulating nitric oxide and pro-inflammatory cytokine responses ex vivo in intra-abdominal sepsis: results from a cohort study. <i>Inflammation Research</i> , 2011, 60, 289-297.	1.6	7
6	Dialyzable Leukocyte Extract differentially regulates the production of TNF $\alpha$ , IL-6, and IL-8 in bacterial component-activated leukocytes and endothelial cells. <i>Inflammation Research</i> , 2005, 54, 74-81.	1.6	28
7	Prophylactic Use of Epidermal Growth Factor Reduces Ischemia/Reperfusion Intestinal Damage. <i>American Journal of Pathology</i> , 2002, 161, 373-379.	1.9	73
8	TNF $\alpha$ Production in Whole Blood Cultures from Healthy Individuals. <i>Biochemical and Biophysical Research Communications</i> , 2002, 292, 538-541.	1.0	21
9	Dialyzable Leukocyte Extract Suppresses the Activity of Essential Transcription Factors for HIV-1 Gene Expression in Unstimulated MT-4 Cells. <i>Biochemical and Biophysical Research Communications</i> , 2000, 273, 1099-1103.	1.0	11
10	Inhibition of in vitro HIV infection by dialysable leucocyte extracts. <i>Biotherapy (Dordrecht, Neth)</i> , 1996, 9, 163-170.	0.7	5
11	Dialysable leucocyte extract (DLE) reduces lipopolysaccharide- induced tumour necrosis factor secretion in human leucocytes. <i>Biotherapy (Dordrecht, Netherlands)</i> , 1996, 9, 163-170.	0.7	7